


MARTIN V. ZOMBECK

HANDBOOK OF SPACE ASTRONOMY & ASTROPHYSICS

SECOND
EDITION



Handbook of space astronomy and astrophysics

K.K 2010

IK 191


C 34
H 22

Handbook of space astronomy and astrophysics

MARTIN V. ZOMBECK

Harvard-Smithsonian Center for Astrophysics

SECOND EDITION



CAMBRIDGE UNIVERSITY PRESS

Cambridge

New York Port Chester Melbourne Sydney

7.96
93

01

Contents

	<i>page</i>
<i>Preface</i>	vii
1 General data	1
2 Astronomy and astrophysics	23
3 Radio astronomy	141
4 Infrared astronomy	159
5 Ultraviolet astronomy	167
6 X-ray astronomy	183
7 Gamma-ray astronomy	211
8 Cosmic rays	219
9 Earth's atmosphere and environment	231
10 Relativity	245
11 Atomic physics	255
12 Electromagnetic radiation	267
13 Plasma physics	283
14 Experimental astronomy and astrophysics	289
15 Aeronautics and astronautics	379
16 Mathematics	395
17 Statistics	407
18 Radiation safety	417
19 Astronomical catalogs	423
Diskettes for PCs	431
<i>Index</i>	433

Preface

I have compiled the tables, graphs, diagrams, and formulae in this book in order to provide a ready reference and working tool for the practicing space astronomer and astrophysicist. Ground-based astronomers and advanced amateur astronomers will find much here of interest, too. The material represents a diversified selection based upon the circumstance that the space astronomer and astrophysicist must draw upon a knowledge of atomic physics, nuclear physics, relativity, plasma physics, electromagnetism, mathematics, statistics, geophysics, experimental physics, *et cetera*, in addition to the classical branches of astronomy. My hope is that this book will replace hunting through many separate works or a trip to the reference library. In that spirit, I welcome suggestions of material for inclusion in a later edition and, of course, corrections or criticism.

There are 19 chapters in the book. The first chapter contains general data, physical constants, unit conversions, etc. Chapters 2-8 cover general astronomy and astrophysics, radio, infrared, ultraviolet, X-ray, and gamma-ray astronomy, and cosmic rays. Chapter 9 contains information on the Earth's atmosphere and environment relevant to space science. Chapter 10 deals with special and general relativity and chapter 11 provides relevant information in atomic physics. Electromagnetic radiation and plasma physics are the subjects of chapters 12 and 13. The remaining chapters essentially deal with the tools of the trade, viz., information on radiation and particle interactions, detectors, launch vehicles, useful mathematical relations and statistical formulae, laboratory radiation safety, and a comprehensive list of astronomical catalogs. Each chapter ends with a brief bibliographical list for further reading on the subject of the chapter or for more extensive reference material. The book has a complete index. References are given and they should be consulted for derivations or comprehensive explanations of the material.

The question of units is always a problem in a book of this type; sticking to one consistent set (SI, for example) is not very useful to the practitioner. I have tried to use the unit systems common to the particular field. Thus I have used SI, c.g.s., and Gaussian (e.s.u. c.g.s. units) or occasionally a mixture, whenever customary. What is being used is usually noted and whenever the units are not noted, any consistent system will do. If in doubt, perform a numerical check. Besides a complete up-to-date set of fundamental constants in SI units, I have also provided a subset in c.g.s. units which are more commonly used in the formulae in this book.

I wish to acknowledge colleagues throughout the scientific community for their useful suggestions; especially Gerald Austin, Paul Gorenstein, Charles Maxson, Robert Rosner, Stephen Murray, Daniel Schwartz, and Daniel Fabricant of the Harvard/Smithsonian Center for Astrophysics. I reserve my deepest appreciation for Judith Peritz who tirelessly and with great skill prepared the typescript and never lost patience with my 'just one more addition'.

Martin V. Zombeck
Cambridge, November 1989

Chapter 1

General data

Fundamental physical constants (SI)	2
Fundamental physical constants (c.g.s.)	10
Sun–Earth system constants	10
Cosmological data	11
Unit conversions	11
Conversion tables	12
Numerical constants	16
Mathematical formulae	17
Elementary particles (short list)	17
Elementary particles	18
Energy conversions	19
Prefixes and symbols	19
Periodic table of the elements	20
Greek alphabet	22
Bibliography	22

Fundamental physical constants (SI)

(1986 recommended values of the fundamental physical constants. The digits in parentheses are the one-standard-deviation uncertainty in the last digits of the given value.)

Quantity	Symbol	Value	Units	Relative uncertainty (ppm)
GENERAL CONSTANTS				
<i>UNIVERSAL CONSTANTS</i>				
speed of light in vacuum	c	299 792 458	m s^{-1}	(exact)
permeability of vacuum	μ_0	$4\pi \times 10^{-7}$ = 12.566 370 614 ...	N A^{-2}	(exact)
permittivity of vacuum	ϵ_0	$1/\mu_0 c^2$ = 8.854 187 817 ...	$10^{-12} \text{ F m}^{-1}$	(exact)
Newtonian constant of gravitation	G	6.672 59(85)	$10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$	128
Planck constant	h	6.626 075 5(40)	10^{-34} J s	0.60
in electron volts, $h/(e)$	h	4.135 669 2(12)	10^{-15} eV s	0.30
$h/2\pi$		1.054 572 66(63)	10^{-34} J s	0.60
in electron volts, $h/(e)$		6.582 122 0(20)	10^{-16} eV s	0.30
Planck mass, $(hc/G)^{1/2}$	m_p	2.176 71(14)	10^{-8} kg	64
Planck length, $h/m_p c = (\hbar G/c^3)^{1/2}$	l_p	1.616 05(10)	10^{-35} m	64
Planck time, $l_p/c = (\hbar G/c^5)^{1/2}$	t_p	5.390 56(34)	10^{-44} s	64
ELECTROMAGNETIC CONSTANTS				
elementary charge	e	1.602 177 33(49)	10^{-19} C	0.30
	e/h	2.417 988 36(72)	10^{14} A J^{-1}	0.30
magnetic flux quantum, $h/2e$	Φ_0	2.067 834 61(61)	10^{-15} Wb	0.30
Josephson frequency-voltage ratio	$2e/h$	4.835 976 7(14)	$10^{14} \text{ Hz V}^{-1}$	0.30
quantized Hall conductance	e^2/h	3.874 046 14(17)	10^{-5} S	0.045
quantized Hall resistance, $h/e^2 = \frac{1}{2}\mu_0 c^2/\alpha$	R_H	25 812.805 6(12)	Ω	0.045

Bohr magneton, $e\hbar/2m_e$				$10^{-24} \text{ J T}^{-1}$	0.34
in electron volts, $\mu_B/(1e)$		9.274 015 4(31)		$10^{-5} \text{ eV T}^{-1}$	0.089
in hertz, μ_B/\hbar		5.788 382 63(52)		$10^{10} \text{ Hz T}^{-1}$	0.30
in wavenumbers, μ_B/hc		1.399 624 18(42)		$\text{m}^{-1} \text{ T}^{-1}$	0.30
in kelvins, μ_B/k		46.686 437(14)		K T^{-1}	8.5
nuclear magneton, $e\hbar/2m_p$		0.671 709 9(57)		$10^{-27} \text{ J T}^{-1}$	0.34
in electron volts, $\mu_N/(1e)$		5.050 786 6(17)		$10^{-8} \text{ eV T}^{-1}$	0.089
in hertz, μ_N/\hbar		3.152 451 66(28)		MHz T^{-1}	0.30
in wavenumbers, μ_N/hc		7.622 591 4(23)		$10^2 \text{ m}^{-1} \text{ T}^{-1}$	0.30
in kelvins, μ_N/k		2.542 622 81(77)		10^{-4} K T^{-1}	8.5

ATOMIC CONSTANTS

ATOM					
fine-structure constant, $\frac{1}{2}\mu_0 c e^2/\hbar$	α	7.297 353 08(33)	10^{-3}		0.045
inverse fine-structure constant	α^{-1}	137.035 989 5(61)	m^{-1}		0.045
Rydberg constant, $\frac{1}{2}m_e c \alpha^2/\hbar$	R_∞	10 973 731.534(13)	10^{15} Hz		0.0012
in hertz, $R_\infty c$		3.289 841 949 9(39)	10^{-18} J		0.60
in joules, $R_\infty \hbar c$		2.179 874 1(13)	eV		0.30
in electron volts, $R_\infty \hbar c/(1e)$	u_0	13.605 698 1(40)	10^{-10} m		0.045
Bohr radius, $\alpha/4\pi R_\infty$	E_h	0.529 177 249(24)	10^{-18} J		0.60
Hartree energy, $e^2/4\pi\epsilon_0 a_0 = 2R_\infty \hbar c$	$\hbar/2m_e$	4.359 748 2(26)	eV		0.30
in electron volts, $E_h/(1e)$	$\hbar/m_e c$	27.211 396 1(81)	$10^{-4} \text{ m}^2 \text{ s}^{-1}$		0.089
quantum of circulation		3.636 948 07(33)	$10^{-4} \text{ m}^2 \text{ s}^{-1}$		0.089

ELECTRON

electron mass	m_e	9.109 389 7(54)	10^{-31} kg		0.59
		5.485 799 03(13)	10^{-4} u		0.023
in electron volts, $m_e c^2/(1e)$	m_e/m_μ	0.510 999 06(15)	MeV		0.30
electron-muon mass ratio	m_e/m_p	4.836 332 18(71)	10^{-2}		0.15
electron-proton mass ratio	m_e/m_d	5.446 170 13(11)	10^{-4}		0.020
electron-deuteron mass ratio	m_e/m_x	2.724 437 07(6)	10^{-4}		0.020
electron α -particle mass ratio		1.370 933 54(3)	10^{-4}		0.021

Fundamental physical constants (SI) (cont.)

Quantity	Symbol	Value	Units	Relative uncertainty (ppm)
electron specific charge	$-e/m_e$	-1.758 819 62(53)	$10^{11} \text{ C kg}^{-1}$	0.30
electron molar mass	$M(e), M_e$	5.485 799 03(13)	$10^{-7} \text{ kg mol}^{-1}$	0.023
Compton wavelength, $h/m_e c$	λ_C	2.426 310 58(22)	10^{-12} m	0.089
$\lambda_C/2\pi = \alpha^2/4\pi R_\infty$	$\tilde{\lambda}_C$	3.861 593 23(35)	10^{-13} m	0.089
classical electron radius, $\alpha^2 a_0$	r_e	2.817 940 92(38)	10^{-15} m	0.13
Thomson cross-section, $(8\pi/3)r_e^2$	σ_e	0.665 246 16(18)	10^{-28} m^2	0.27
electron magnetic moment	μ_e	928.477 01(31)	$10^{-26} \text{ J T}^{-1}$	0.34
in Bohr magnetons	μ_e/μ_B	1.001 159 652 193(10)		1×10^{-5}
in nuclear magnetons	μ_e/μ_N	1838.282 000(37)		0.020
electron magnetic moment anomaly, $\mu_e/\mu_B - 1$	a_e	1.159 652 193(10)	10^{-3}	0.0086
electron g-factor, $2(1 + a_e)$	g_e	2.002 319 304 386(20)		1×10^{-5}
electron-muon magnetic moment ratio	μ_e/μ_μ	206.766 967(30)		0.15
electron-proton magnetic moment ratio	μ_e/μ_p	658.210 688 1(66)		0.010
MUON				
muon mass	m_μ	1.883 532 7(11)	10^{-28} kg	0.61
		0.113 428 913(17)	u	0.15
in electron volts, $m_\mu c^2/(e)$		105.658 389(34)	MeV	0.32
muon-electron mass ratio	m_μ/m_e	206.768 262(30)		0.15
muon molar mass	$M(\mu), M_\mu$	1.134 289 13(17)	$10^{-4} \text{ kg mol}^{-1}$	0.15
muon magnetic moment	μ_μ	4.490 451 4(15)	$10^{-26} \text{ J T}^{-1}$	0.33
in Bohr magnetons	μ_μ/μ_B	4.841 970 97(71)	10^{-3}	0.15
in nuclear magnetons	μ_μ/μ_N	8.890 598 1(13)		0.15
muon magnetic moment anomaly, $[\mu_\mu/(eh/2m_\mu)] - 1$	a_μ	1.165 923 0(84)		7.2
muon g-factor, $2(1 + a_\mu)$	g_μ	2.002 331 846(17)	10^{-3}	0.0084
muon-proton magnetic moment ratio	μ_μ/μ_p	3.183 345 47(47)		0.15



PROTON

proton mass	m_p	1.672 623 1(10)	10^{-27} kg	0.59
in electron volts, $m_p c^2 / \{e\}$		1.007 276 470(12)	u	0.012
proton-electron mass ratio	m_p / m_e	938.272 31(28)	MeV	0.30
proton-muon mass ratio	m_p / m_μ	1836.152 701(37)		0.020
proton specific charge	e / m_p	8.880 244 4(13)	10^7 C kg ⁻¹	0.15
proton molar mass	$M(p), M_p$	9.578 830 9(29)	10^{-3} kg mol ⁻¹	0.30
proton Compton wavelength, $h / m_p c$	$\lambda_{C,p}$	1.007 276 470(12)	10^{-15} m	0.012
$\lambda_{C,p} / 2\pi$	$\lambda_{C,p} / 2\pi$	1.321 410 02(12)	10^{-16} m	0.089
proton magnetic moment	μ_p	2.103 089 37(19)	10^{-26} J T ⁻¹	0.089
in Bohr magnetons	μ_p / μ_B	1.410 607 61(47)	10^{-3}	0.34
in nuclear magnetons	μ_p / μ_N	1.521 032 202(15)		0.010
diamagnetic shielding correction for protons in pure water, spherical sample, 25 °C, 1 - μ_p / μ_p shielded proton moment (H ₂ O, sph., 25 °C)	σ_{H_2O}	2.792 847 386(63)		0.023
in Bohr magnetons	μ_p' / μ_B	25.689(15)	10^{-6}	0.34
in nuclear magnetons	μ_p' / μ_N	1.410 571 38(47)	10^{-26} J T ⁻¹	0.011
proton gyromagnetic ratio	γ_p	1.520 993 129(17)	10^{-3}	0.023
uncorrected (H ₂ O, sph., 25 °C)	$\gamma_p / 2\pi$	2.792 775 642(64)	10^4 s ⁻¹ T ⁻¹	0.30
	$\gamma_p / 2\pi$	26 752.2128(81)	MHz T ⁻¹	0.30
	$\gamma_p / 2\pi$	42.577 469(13)	10^4 s ⁻¹ T ⁻¹	0.30
	$\gamma_p / 2\pi$	26 751.5255(81)	MHz T ⁻¹	0.30
	$\gamma_p / 2\pi$	42.576 375(13)		0.30

NEUTRON

neutron mass	m_n	1.674 928 6(10)	10^{-27} kg	0.59
in electron volts, $m_n c^2 / \{e\}$		1.008 664 904(14)	u	0.014
neutron-electron mass ratio	m_n / m_e	939.565 63(28)	MeV	0.30
neutron proton mass ratio	m_n / m_p	1838.683 662(40)		0.022
neutron molar mass	$M(n), M_n$	1.001 378 404(9)	10^{-3} kg mol ⁻¹	0.009
neutron Compton wavelength, $h / m_n c$	$\lambda_{C,n}$	1.008 664 904(14)	10^{-15} m	0.014
$\lambda_{C,n} / 2\pi$	$\lambda_{C,n} / 2\pi$	1.319 591 10(12)	10^{-16} m	0.089
	$\lambda_{C,n} / 2\pi$	2.100 194 45(19)		0.089

Fundamental physical constants (SI) (cont.)

Quantity	Symbol	Value	Units	Relative uncertainty (ppm)
neutron magnetic moment ^(a)	μ_n	0.966 237 07(40)	$10^{-26} \text{ J T}^{-1}$	0.41
in Bohr magnetons	μ_n/μ_B	1.041 875 63(25)	10^{-3}	0.24
in nuclear magnetons	μ_n/μ_N	1.913 042 75(45)		0.24
neutron-electron magnetic moment ratio	μ_n/μ_e	1.040 668 82(25)	10^{-3}	0.24
neutron-proton magnetic moment ratio	μ_n/μ_p	0.684 979 34(16)		0.24
DEUTERON				
deuteron mass	m_d	3.343 586 0(20)	10^{-27} kg	0.59
in electron volts, $m_d c^2/\{e\}$		2.013 553 214(24)	u	0.012
deuteron-electron mass ratio	m_d/m_e	1875.613 39(57)	MeV	0.30
deuteron-proton mass ratio	m_d/m_p	3670.483 014(75)		0.020
deuteron molar mass	$M(d), M_d$	1.999 007 496(6)		0.003
deuteron magnetic moment ^(a)	μ_d	2.013 553 214(24)	$10^{-3} \text{ kg mol}^{-1}$	0.012
in Bohr magnetons	μ_d/μ_B	0.433 073 75(15)	$10^{-26} \text{ J T}^{-1}$	0.34
in nuclear magnetons	μ_d/μ_N	0.466 975 447 9(91)	10^{-3}	0.019
deuteron-electron magnetic moment ratio	μ_d/μ_e	0.857 438 230(24)		0.028
deuteron-proton magnetic moment ratio	μ_d/μ_p	0.466 434 546 0(91)	10^{-3}	0.019
		0.307 012 203 5(51)		0.017

PHYSICO-CHEMICAL CONSTANTS

Avogadro constant	N_A, L	6.022 136 7(36)	10^{23} mol^{-1}	0.59
atomic mass constant, $m_u = \frac{1}{12}m(^{12}\text{C})$	m_u	1.660 540 2(10)	10^{-27} kg	0.59
in electron volts, $m_u c^2/\{e\}$		931.494 32(28)	MeV	0.30
Faraday constant	F	96 485.309(29)	C mol^{-1}	0.30

molar Planck constant	$N_A h$	3.990 313 23(36)	$10^{-10} \text{ J s mol}^{-1}$	0.089
molar gas constant	$N_A h c$	0.119 626 58(11)	J m mol^{-1}	0.089
Boltzmann constant, R/N_A	R	8.314 510(70)	$\text{J mol}^{-1} \text{ K}^{-1}$	8.4
in electron volts, k/e_1	k	1.380 658(12)	$10^{-23} \text{ J K}^{-1}$	8.5
in hertz, k/h		8.617 385(73)	$10^{-5} \text{ eV K}^{-1}$	8.4
in wavenumbers, k/hc		2.083 674(18)	$10^{10} \text{ Hz K}^{-1}$	8.4
molar volume (ideal gas), RT/p		69.503 87(59)	$\text{m}^3 \text{ K}^{-1}$	8.4
$T = 273.15 \text{ K}$, $p = 101 325 \text{ Pa}$	V_m	22.414 10(19)	L mol^{-1}	8.4
Loschmidt constant, N_A/V_m	n_0	2.686 763(23)	10^{25} m^{-3}	8.5
$T = 273.15 \text{ K}$, $p = 100 \text{ kPa}$	V_m	22.711 08(19)	L mol^{-1}	8.4
Sackur-Tetrode constant (absolute entropy constant), ^(a) $\frac{5}{2} + \ln\{(2\pi m_0 k T_1 / h^2) k T_1 / p_0\}$	S_0/R	-1.151 693(21)		18
$T_1 = 1 \text{ K}$, $p_0 = 100 \text{ kPa}$		-1.164 856(21)		18
$p_0 = 101 325 \text{ Pa}$	σ	5.670 51(19)	$10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$	34
Stefan-Boltzmann constant, $(\pi^2/60)k^4/h^3c^2$	c_1	3.741 774 9(22)	10^{-16} W m^2	0.60
first radiation constant, $2\pi hc^2$	c_2	0.014 387 69(12)	m K	8.4
second radiation constant, hc/k	b	2.897 756(24)	10^{-3} m K	8.4
Wien displacement law constant, $b = \lambda_{\text{max}} T = c_2/4.965 114 23 \dots$				

^(a) The scalar magnitude of the neutron moment is listed here. The neutron magnetic dipole is directed oppositely to that of the proton, and corresponds to the dipole associated with a spinning negative charge distribution. The vector sum, $\mu_d = \mu_p + \mu_n$, is approximately satisfied.

^(b) The entropy of an ideal monatomic gas of relative atomic weight A_r is given by

$$S = S_0 + \frac{5}{2}R \ln A_r - R \ln(p/p_0) + \frac{3}{2}R \ln(T/K) \dots$$

Fundamental physical constants (SI) (cont.)

MAINTAINED UNITS AND STANDARD VALUES

A summary of 'maintained' units and 'standard' values and their relationship to SI units, based on a least-squares adjustment with 17 degrees of freedom. The digits in parentheses are the one-standard-deviation uncertainty in the last digits of the given value.

Quantity	Symbol	Value	Units	Relative uncertainty (ppm)
electron volt, $(e/C) J = \{e\} J$	eV	1.602 177 33(49)	$10^{-19} J$	0.30
(unified) atomic mass unit, $1 u = m_0 = \frac{1}{12} m(^{12}C)$	u	1.660 540 2(10)	$10^{-27} kg$	0.59
standard atmosphere	atm	101 325	Pa	(exact)
standard acceleration of gravity	g_n	9.806 65	$m s^{-2}$	(exact)
'AS-MAINTAINED' ELECTRICAL UNITS				
BIPM ^(a) maintained ohm, Ω_{69-BI}	Ω_{BIS}	$1 - 1.563(50) \times 10^{-6}$ $= 0.999 998 437(50)$	Ω	0.050
drift rate of Ω_{69-BI}	$\frac{d\Omega_{69-BI}}{dt}$	-0.0566(15)	$\mu\Omega/a$	—
BIPM maintained volt.	V_{76-BI}	$1 - 7.59(30) \times 10^{-6}$ $= 0.999 992 41(30)$	V	0.30
BIPM maintained ampere.	A_{BIS}	$1 - 6.03(30) \times 10^{-6}$ $= 0.999 993 97(30)$	A	0.30
X-RAY STANDARDS				
Cu x-unit: $\lambda(CuK\alpha_1) \equiv 1537.400$ xu	xu(CuK α_1)	1.002 077 89(70)	$10^{-13} m$	0.70
Mo x-unit: $\lambda(MoK\alpha_1) \equiv 707.831$ xu	xu(MoK α_1)	1.002 099 38(45)	$10^{-13} m$	0.45
\dot{A}^* : $\lambda(WK\alpha_1) \equiv 0.209 100 \dot{A}^*$	\dot{A}^*	1.000 014 8 1(92)	$10^{-10} m$	0.92

lattice spacing of Si (in vacuum, 22.5 C) ^(a)	a	0.543 101 96(11)	nm	0.21
$d_{220} = a/\sqrt{8}$	d_{220}	0.192 015 540(40)	nm	0.21
molar volume of Si, $M(\text{Si})/\rho(\text{Si}) = N_A a^3/8$	$V_m(\text{Si})$	12.058 817 9(89)	cm ³ mol ⁻¹	0.74

^(a) BIPM: Bureau International des Poids et Mesures.

^(b) The lattice spacing of single-crystal Si can vary by parts in 10⁷ depending on the preparation process. Measurements at Physikalisch-Technische Bundesanstalt (PTB) indicate also the possibility of distortions from exact cubic symmetry of the order of 0.2 ppm.

(Reprinted with permission from CODATA Bulletin, Number 63, Cohen, E. Richard & Taylor, Barry N., *The 1986 Adjustment of the Fundamental Physical Constants*, Copyright 1987, Pergamon Press, Ltd.)

Fundamental physical constants (c.g.s.)

Speed of light in vacuum	$c = 2.997\,924\,58 \times 10^{10} \text{ cm s}^{-1}$
Gravitational constant	$G = 6.672\,59 \times 10^{-8} \text{ dyn cm}^2 \text{ g}^{-2}$
Planck's constant	$h = 6.626\,075\,5 \times 10^{-27} \text{ erg s}$
Electron charge	$e = 4.803\,206\,8 \times 10^{-10} \text{ esu}$
Mass of electron	$m_e = 9.109\,389\,7 \times 10^{-28} \text{ g}$
Mass of proton	$m_p = 1.672\,623\,1 \times 10^{-24} \text{ g}$
Mass of neutron	$m_n = 1.674\,928\,6 \times 10^{-24} \text{ g}$
Atomic mass unit (amu)	$m_u = 1.660\,540\,2 \times 10^{-24} \text{ g}$
Proton-electron mass ratio	$m_p/m_e = 1\,836.152\,701$
Fine structure constant	$hc/2\pi e^2 = 1/\alpha = 137.035\,989\,5$
Classical electron radius	$e^2/m_e c^2 = r_e = 2.817\,940\,92 \times 10^{-13} \text{ cm}$
Bohr radius	$h^2/4\pi^2 m_e e^2 = a_0 = 0.529\,177\,249 \times 10^{-8} \text{ cm}$
Electron Compton wavelength	$h/m_e c = \lambda_c = 2.426\,310\,58 \times 10^{-10} \text{ cm}$
Rydberg constant	$2\pi^2 m_e e^4 / ch^3 = R_\infty = 109\,737.315\,34 \text{ cm}^{-1}$
Boltzmann constant	$k = 1.380\,658 \times 10^{-16} \text{ erg K}^{-1}$
Stefan-Boltzmann constant	$\sigma = 2\pi^5 k^4 / 15 h^3 c^2$ $= 5.670\,51 \times 10^{-5} \text{ erg cm}^{-2} \text{ K}^{-4} \text{ s}^{-1}$
Thomson cross-section	$8\pi r_e^2 / 3 = \sigma_T = 0.665\,246\,16 \times 10^{-24} \text{ cm}^2$
Bohr magneton	$eh/4\pi m_e = \mu_B = 9.274\,015\,4 \times 10^{-21} \text{ gauss cm}^3$

(Based on constants recommended by the 1986 CODATA Committee in previous table.)

Sun-Earth system constants

Equatorial radius for Earth	$a_e = 6378.140 \text{ km}$
Dynamical form-factor for Earth	$J_2 = 0.001\,082\,63$
Gravity at Earth's surface (mean)	$g_e = 980.7 \text{ cm s}^{-2}$
Ratio of mass of Moon to that of Earth	$\mu = 0.012\,300\,02$
Lunar distance (mean)	$3.844\,01 \times 10^{10} \text{ cm}$
Astronomical unit	$\text{AU} = 1.495\,978\,70 \times 10^{13} \text{ cm}$
Mass of the Earth	$M_e = 5.976 \times 10^{27} \text{ g}$
Solar parallax	$\pi_\odot = 8''.794\,148$
Tropical year (1900.0)	365.242 days $3.1557 \times 10^7 \text{ ephemeris seconds}$
Ephemeris day	86 400 ephemeris seconds
Constant of aberration (2000.0)	$K = 20''.495\,52$
Obliquity of the ecliptic (2000.0)	$\varepsilon = 23^\circ 26' 21''.448$
General precession in longitude per Julian century (2000.0)	$p = 5029''.0966$
Constant of nutation (2000.0)	$N = 9''.2055$
Earth's magnetic dipole moment	$M_e = 8.02 \times 10^{22} \text{ Am}^2$
Angular rotation rate of Earth	$\Omega = 7.292\,115 \times 10^{-5} \text{ rad s}^{-1}$
Average depth of ocean	$H = 3800 \text{ m}$
Flattening of Earth	$1/f = 298.257$

Cosmological data

Hubble constant	$H_0 = (50-100) \text{ km s}^{-1} \text{ Mpc}^{-1}$ $= (1.6-3.2) \times 10^{-18} \text{ s}^{-1}$
Hubble time	$1/H_0 = (19.6-9.78) \times 10^9 \text{ y}$
Hubble distance	$R = c/H_0 = (6050-3025) \text{ Mpc}$
Critical density	$\rho_c = 3H_0^2/8\pi G$ $= (5-20) \times 10^{-30} \text{ g cm}^{-3}$
Volume	$\frac{4}{3}\pi R^3 = (9-1) \times 10^{11} \text{ Mpc}^3$
Smoothed density of galactic material throughout universe (Allen 1973)	$2 \times 10^{-31} \text{ g cm}^{-3}$ $1 \times 10^{-7} \text{ atom cm}^{-3}$ $3 \times 10^9 M_\odot \text{ Mpc}^{-3}$
Space density of galaxies	0.02 Mpc^{-3}
Luminous emission from galaxies	$3 \times 10^8 L_\odot \text{ Mpc}^{-3}$
Mean sky brightness from galaxies	$1.4(m_V = 10) \text{ deg}^{-2}$
Cosmic background thermodynamic temperature	$2.74 \pm 0.09 \text{ K}$
Weak coupling constant	$g_{\text{wk}} = 1.435 \times 10^{-49} \text{ erg cm}^3$

Unit conversions

1 keV: $hc/E = 12.39854 \times 10^{-8} \text{ cm}$	1 keV = $1.602177 \times 10^{-9} \text{ erg}$
1 keV: $E/h = 2.417965 \times 10^{17} \text{ Hz}$	1 joule = 10^7 erg
1 keV: $E/k = 11.6048 \times 10^6 \text{ K}$	1 calorie = 4.184 joule
1.0 EHz: $h\nu = 4.13571 \text{ keV}$	
1 parsec = $3.261633 \text{ light years} = 3.085678 \times 10^{18} \text{ cm}$	
1 light year = $9.460530 \times 10^{17} \text{ cm}$	
1 XU = $1.00209 \times 10^{-11} \text{ cm}$	
1 Ångstrom $\equiv 1 \times 10^{-8} \text{ cm}$	
1 amu: $Mc^2 = 1.49241 \times 10^{-3} \text{ erg} = 931.494 \text{ MeV}$	
760 torr = $1.013 \times 10^6 \text{ dyn cm}^{-2} = 1 \text{ atmos.} = 1.013 \text{ bars} = 1.013 \times 10^5 \text{ Pascals}$	
1 Rayleigh $\equiv (1/4\pi) \times 10^6 \text{ photons cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$	
1 Uhuru ct $\text{s}^{-1} = 1.7 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$ (2-6 keV)	
$= 2.4 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$ (2-10 keV)	
1 flux unit $\equiv 10^{-26} \text{ watt m}^{-2} \text{ Hz}^{-1} \equiv 1 \text{ Jansky}$	
1.0 $\mu\text{Jy} = 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ EHz}^{-1}$	
$= 0.242 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1} = 1.509 \times 10^{-3} \text{ keV cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1}$	
1 curie: amount of material undergoing $3.7 \times 10^{10} \text{ disintegrations s}^{-1}$	
1 nautical mile = 1852 m	
1 statute mile = 1609.344 m	
intensity ($\text{erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1}$) = $3.33 \times 10^{-19} \lambda^2(\text{Å}) \text{ intensity} (\text{erg cm}^{-2} \text{ s}^{-1} \text{ Å}^{-1})$	
1 barn = 10^{-24} cm^2	
1 tesla = 10^4 Gauss	
0 °C = 273.15 K	

Conversion tables

(A given amount of a physical quantity, expressed in the units of one system, is expressed as an equivalent number of units in another system.)

Quantity	Amount	Unit	Amount	Unit
LENGTH				
	1	meter (SI) = $1.000\ 00E + 02$		centimeter (cgs)
	1	light year = $9.460\ 53E + 15$		meter (SI)
	1	parsec = $3.085\ 68E + 16$		meter (SI)
	1	Ångstrom = $1.000\ 01E - 10$		meter (SI)
	1	Ångstrom = $1.000\ 01E - 08$		centimeter (cgs)
	1	micron = $1.000\ 00E - 06$		meter (SI)
	1	XU = $1.002\ 09E - 13$		meter (SI)
	1	fermi = $1.000\ 00E - 15$		meter (SI)
	1	nautical mile = $1.852\ 00E + 03$		meter (SI)
	1	statute mile = $1.609\ 34E + 03$		meter (SI)
	1	astron. unit (AU) = $1.495\ 99E + 11$		meter (SI)
	1	solar radius = $6.959\ 90E + 08$		meter (SI)
	1	centimeter (cgs) = $3.240\ 78E - 19$		parsec
	1	centimeter (cgs) = $6.684\ 56E - 14$		astron. unit (AU)
	1	meter (SI) = $3.240\ 78E - 17$		parsec
	1	meter (SI) = $6.684\ 54E - 12$		astron. unit (AU)
	1	inch (Eng) = $2.540\ 00E - 02$		meter (SI)
MASS				
	1	kilogram (SI) = $1.000\ 00E + 03$		gram (cgs)
	1	at. mass unit (amu) = $1.660\ 54E - 24$		gram (cgs)
	1	at. mass unit (amu) = $1.660\ 54E - 27$		kilogram (SI)
	1	solar mass = $1.989\ 10E + 33$		gram (cgs)
	1	solar mass = $1.989\ 10E + 30$		kilogram (SI)

1 gram (cgs) = 6.022 14E + 23
 1 gram (cgs) = 5.027 40E - 34
 1 kilogram (SI) = 6.022 14E + 26
 1 kilogram (SI) = 5.027 40E - 31
 1 kilogram (SI) = 2.204 62E + 00
 1 kilogram (SI) = 3.527 40E + 01
 1 pound (avdp.) = 4.535 92E - 01
 1 pound (avdp.) = 1.600 00E + 01
 1 ounce (avdp.) = 2.834 95E + 01
 1 gram (cgs) = 3.527 40E - 02

 1 joule (SI) = 1.000 00E + 07
 1 joule (SI) = 6.241 51E + 18
 1 erg (cgs) = 1.000 00E - 07
 1 erg (cgs) = 6.241 51E + 11
 1 electron volt = 1.602 18E - 12
 1 amu x c² = 9.314 95E + 08
 1 gm (cgs) x c² = 5.609 59E + 32
 1 calorie = 4.184 00E + 00

ENERGY

1 joule (SI) = 1.000 00E + 07
 1 joule (SI) = 6.241 51E + 18
 1 erg (cgs) = 1.000 00E - 07
 1 erg (cgs) = 6.241 51E + 11
 1 electron volt = 1.602 18E - 12
 1 amu x c² = 9.314 95E + 08
 1 gm (cgs) x c² = 5.609 59E + 32
 1 calorie = 4.184 00E + 00

FORCE

1 newton (SI) = 1.000 00E + 05
 1 dyne (cgs) = 1.000 00E - 05

PRESSURE

1 pascal (SI) = 1.000 00E + 00
 1 bar = 1.000 00E + 06
 1 bar = 9.869 23E - 01
 1 torr = 1.333 22E - 03

POWER

1 watt (SI) = 1.000 00E + 07
 1 horsepower = 7.457 00E + 02
 1 Btu s⁻¹ (Eng) = 1.055 80E + 03

at. mass unit (amu)
 solar mass
 at. mass unit (amu)
 solar mass
 pound (avdp.)
 ounce (avdp.)
 kilogram (SI)
 ounce (avdp.)
 gram (cgs)
 ounce (avdp.)

erg (cgs)
 electron volt (eV)
 joule (SI)
 electron volt
 erg (cgs)
 electron volt
 electron volt
 joule (SI)

dyne (cgs)
 newton (SI)

newton m⁻² (SI)
 dyne cm⁻² (cgs)
 atmosphere
 bar

erg s⁻¹ (cgs)
 watt (SI)
 watt (SI)

Conversion tables (cont.)

Quantity	Amount	Unit	Amount	Unit
TIME				
	1	second (SI) = 1		second (cgs)
	1	minute = 6.000 00E + 01		second
	1	hour = 3.600 00E + 03		second
	1	day = 8.640 00E + 04		second
	1	tropical year = 3.155 69E + 07		second
	1	tropical year = 3.652 42E + 02		day
	1	second = 3.168 88E - 08		tropical year
	1	sidereal second = 9.972 70E - 01		second
	1	sidereal year = 3.652 56E + 02		day
TEMPERATURE				
	<i>T</i>	kelvin = $T - 273.15$		celsius
	<i>T</i>	kelvin = $(9/5) \times (T - 273.15) + 32$		fahrenheit
	<i>T</i>	celsius = $T + 273.15$		kelvin
	<i>T</i>	fahrenheit = $(5/9) \times (T - 32) + 273.15$		kelvin
	<i>T</i>	celsius = $(9/5) \times T + 32$		fahrenheit
	<i>T</i>	fahrenheit = $(5/9) \times (T - 32)$		celsius
Energy equivalence	1	electron volt: 1.160 48E + 04		kelvin
Temperature equivalence	1	kelvin: 8.617 12E - 05		electron volt
ELECTRICITY AND MAGNETISM				
Charge	1	coulomb = 2.997 92E + 09		statcoulomb
Charge density	1	coulomb m ⁻³ = 2.997 92E + 03		statcoul cm ⁻³
Current	1	ampere (coul s ⁻¹) = 2.997 92E + 09		statampere
Current density	1	ampere m ⁻² = 2.997 92E + 05		statamp cm ⁻²
Electric field	1	volt m ⁻¹ = 3.335 65E 05		statvolt cm ⁻¹
Potential	1	volt = 3.335 65E - 03		statvolt

Resistance	1	ohm = 1.112 65E - 12	s cm^{-1}
Resistivity	1	ohm m = 1.112 65E - 10	s
Conductance	1	siemens, mho = 8.987 52E + 11	cm s^{-1}
Conductivity	1	mho m^{-1} = 8.987 52E + 09	s^{-1}
Capacitance	1	farad = 8.987 52E + 11	cm
Magnetic flux	1	weber = 1.000 00E + 08	gauss cm^2 (maxwell)
Magnetic flux density	1	tesla = 1.000 00E + 04	gauss
Magnetic field	1	ampere-turn m^{-1} = 1.256 64E - 02	oersted
Inductance	1	henry = 1.112 65E - 12	$\text{s}^2 \text{cm}^{-1}$
MISCELLANEOUS			
Radio-activity	1	curie (SI) = 3.700 00E + 10	disinteg. s^{-1}
Intensity	1	rayleigh = 7.957 75E + 04	$\text{ph cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$
Flux density	1	fu or jansky = 1.000 00E - 26	$\text{watt m}^{-2} \text{Hz}^{-1}$
Flux density	1	jansky = 1.000 00E - 08	$\text{erg cm}^{-2} \text{s}^{-1} \text{EHZ}^{-1}$
Flux density	1	jansky = 2.420 00E - 09	$\text{erg cm}^{-2} \text{s}^{-1} \text{keV}^{-1}$
Flux density	1	jansky = 1.509 00E + 00	$\text{keV cm}^{-2} \text{s}^{-1} \text{keV}^{-1}$
Energy equivalence	1	eV: 1.239 85E + 04	Angstrom
Energy equivalence	1	eV: 2.417 97E + 14	Hz
Wavelength equivalence	1	Angstrom: 1.239 85E + 04	eV
Angle	1	arcsec = 4.848 14E - 06	radian
Angle	1	arcmin = 2.908 88E - 04	radian
Angle	1	degree = 1.745 33E - 02	radian
Solid angle	1	arcsec ² = 2.350 40E - 11	steradian
Solid angle	1	arcmin ² = 8.461 70E - 08	steradian
Solid angle	1	deg ² = 3.046 20E - 04	steradian

Numerical constants

$\pi = 3.141\ 592\ 7$	$\text{rad} = 57.295\ 78\ \text{deg}$
$e = 2.718\ 281\ 8$	$= 3.437\ 747 \times 10^4\ \text{arcmin}$
$\ln 2 = 0.693\ 147\ 2$	$= 2.062\ 648 \times 10^5\ \text{arcsec}$
$\log_{10} 2 = 0.301\ 030\ 0$	$\text{steradian} = 32\ 400/\pi^2 = 3.2828 \times 10^3\ \text{deg}^2$
$\ln 10 = 2.302\ 585\ 1$	$= 1.1818 \times 10^7\ \text{arcmin}^2$
$\log_{10} e = 0.434\ 294\ 5$	$= 4.2545 \times 10^{10}\ \text{arcsec}^2$
$(2\pi)^{1/2} = 2.506\ 628$	$\text{degree} = 0.017\ 453\ 3\ \text{rad}$
$\pi^2 = 9.869\ 604$	$\text{arcmin} = 2.908\ 88 \times 10^{-4}\ \text{rad}$
$2^{10} = 1024$	$\text{arcsec} = 4.848\ 137 \times 10^{-6}\ \text{rad}$
$e^{-1} = 0.367\ 879\ 4$	$\text{deg}^2 = 3.0462 \times 10^{-4}\ \text{steradian}$
$\Gamma(n+1) = n!$, integer n	$\text{arcmin}^2 = 8.4617 \times 10^{-8}\ \text{steradian}$
$\Gamma(1/2) = \pi^{1/2}$	$\text{arcsec}^2 = 2.3504 \times 10^{-11}\ \text{steradian}$
Feigenbaum's number: $\delta = 4.669\ 201\ 6$	

n	<i>Powers of 2:</i>		<i>Number system conversions:</i>			
	2^n	$\log 2^n$	Decimal	Octal	Binary	Hexadecimal
0	1	0.00	0	0	0000	0
1	2	0.30	1	1	0001	1
2	4	0.60	2	2	0010	2
3	8	0.90	3	3	0011	3
4	16	1.20	4	4	0100	4
5	32	1.51	5	5	0101	5
6	64	1.81	6	6	0110	6
7	128	2.11	7	7	0111	7
8	256	2.41	8	10	1000	8
9	512	2.71	9	11	1001	9
10	1024	3.01	10	12	1010	A
11	2048	3.31	11	13	1011	B
12	4096	3.61	12	14	1100	C
13	8192	3.91	13	15	1101	D
14	16384	4.21	14	16	1110	E
15	32768	4.52	15	17	1111	F
20	1048576	6.02	16	20	10000	10
25	33554432	7.53				
n		$0.301n$				

Mathematical formulae

$$(a+x)^n = a^n + na^{n-1}x + \frac{n(n-1)}{2!}a^{n-2}x^2 + \frac{n(n-1)(n-2)}{3!}a^{n-3}x^3 + \dots + na^{n-1}x^{n-1} + x^n,$$

where n is any positive integer.

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots,$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \text{ for } -1 < x \leq 1.$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots,$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots,$$

$$\int u \, dv = uv - \int v \, du + C,$$

$$\int_0^\pi \sin^2 nx \, dx = \int_0^\pi \cos^2 nx \, dx = \frac{\pi}{2}$$

for an integer, $n \neq 0$.

$$\int_0^x e^{-a^2x^2} \, dx = \pi^{1/2}/2a,$$

$$\int_0^x x^n e^{-ax} \, dx = \Gamma(n+1)/a^{n+1}.$$

$$F(u) = \int_{-x}^x f(x) e^{-2\pi iux} \, dx \leftrightarrow f(x) = \int_{-x}^x F(u) e^{2\pi iux} \, du.$$

$$n! \approx (2\pi)^{1/2} n^{n+1/2} e^{-n} \text{ (for large } n).$$

$$\int_a^b f(x) \, dx = - \int_b^a f(x) \, dx,$$

$$\int_a^b f(x) \, dx = \int_a^c f(x) \, dx + \int_c^b f(x) \, dx.$$

$$\frac{d}{dx} f[u(x)] = \frac{df}{du} \frac{du}{dx},$$

$$\frac{d}{dx} [u(x)v(x)] = v \frac{du}{dx} + u \frac{dv}{dx}.$$

$$\frac{d}{dx} \ln y(x) = y'(x)/y(x),$$

$$2.30 \log_{10} x = \log_e x$$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B.$$

$$\cos(A+B) = \cos A \cos B - \sin A \sin B.$$

$$e^{ix} = \cos x + i \sin x.$$

Elementary particles (short list)

Particle	Charge	Mass (amu)	Spin	Magnetic moment	Mean life ^(a) (s)
Photon	0	0	1	0	stable
π -meson	+1, -1	0.149 84	1	0	2.60×10^{-8}
	0	0.144 90	0	0	0.83×10^{-16}
Neutrino	0	$< 10^{-6}$	1/2	~ 0	stable
Electron, positron	-1, +1	0.000 548 6	1/2	$1.001 160 \mu_B^{(b)}$	stable
μ -meson	-1, +1	0.1134	1/2	$0.004 842 \mu_B^{(b)}$	2.20×10^{-6}
Proton	+1, -1	1.007 276	1/2	$2.792 85 \mu_N^{(c)}$	stable
Neutron	0	1.008 665	1/2	$-1.913 04 \mu_N$	917 ± 14

^(a) half-life = mean life $\times \ln 2$

^(b) $\mu_B = eh/4\pi m_e c = 9.274 015 4(31) \times 10^{-24} \text{ J T}^{-1}$

^(c) $\mu_N = eh/4\pi m_p c = 5.050 786 6(17) \times 10^{-27} \text{ J T}^{-1}$.

Data from 'Reviews of Particle Properties', *Rev. Mod. Phys.* **52**, No. 2, April 1980)

Elementary Particles

(The second column is the isospin t , while the next column is the spin and parity, J^P . Masses and lifetimes have generally been rounded; see the original reference for error bars and a complete listing of particle properties.)

Particle	t	J^P	Mass (MeV)	Mean life (s)
LEPTONS				
e		$\frac{1}{2}$	0.511 003	Stable
μ		$\frac{1}{2}$	105.6594	$2.197\ 14 \times 10^{-6}$
τ		$\frac{1}{2}$	1784	5×10^{-13}
NONSTRANGE BARYONS				
p	$\frac{1}{2}$	$\frac{1}{2}^+$	938.280	Stable
n	$\frac{1}{2}$	$\frac{1}{2}^+$	939.573	925
Δ	$\frac{3}{2}$	$\frac{3}{2}^+$	1232	6×10^{-24}
STRANGENESS = -1 BARYONS				
Λ	0	$\frac{1}{2}^+$	1115.60	2.63×10^{-10}
Σ^+	1	$\frac{1}{2}^+$	1189.36	8.00×10^{-11}
Σ^0	1	$\frac{1}{2}^+$	1192.46	6×10^{-20}
Σ^-	1	$\frac{1}{2}^+$	1197.34	1.48×10^{-10}
STRANGENESS = -2 BARYONS				
Ξ^0	$\frac{1}{2}$	$\frac{1}{2}^+$	1314.9	2.9×10^{-10}
Ξ^-	$\frac{1}{2}$	$\frac{1}{2}^+$	1321.3	1.64×10^{-10}
STRANGENESS = -3 BARYON				
Ω^-	0	$\frac{3}{2}^+$	1672.5	8.2×10^{-11}
NONSTRANGE CHARMED BARYON				
Λ_c^+	0	$\frac{1}{2}^+$	2282	1×10^{-13}
NONSTRANGE MESONS				
π^\pm	1	0^-	139.567	2.603×10^{-8}
π^0	1	0^-	134.963	8.3×10^{-17}
η	0	0^-	548.8	8×10^{-19}
ρ	1	1^-	769	4.3×10^{-24}
ω	0	1^-	782.6	6.6×10^{-23}
η'	0	0^-	957.6	2.4×10^{-21}
ϕ	0	1	1019.6	1.6×10^{-22}
J/Ψ	0	1^-	3096.9	1.0×10^{-20}
Υ		1^-	9456	1.6×10^{-20}
STRANGENESS = -1 MESONS				
K^\pm	$\frac{1}{2}$	0^-	493.67	1.237×10^{-8}
K^0, \bar{K}^0	$\frac{1}{2}$	0^-	497.7	$K_S: 8.92 \times 10^{-11}$ $K_L: 5.18 \times 10^{-8}$
CHARMED NONSTRANGE MESONS				
D^\pm	$\frac{1}{2}$	0^-	1869.4	9×10^{-13}
D^0, \bar{D}^0	$\frac{1}{2}$	0^-	1864.7	5×10^{-13}
CHARMED STRANGE MESON				
F^\pm	0	0^-	2021	2×10^{-13}

(From Shapiro, S. L. & Teukolsky, S. A., *Black Holes, White Dwarfs, and Neutron Stars*, John Wiley and Sons, 1983, with permission.)

Energy conversions

1 erg	= 1 dyne-centimeter = 10^{-7} joule
1 joule	= 1 newton-meter
1 foot-pound	= 1.356 joule
1 calorie	= 4.184 joule
1 Btu	= 1.055×10^3 joule
1 horsepower-hour	= 2.6845×10^6 joule
1 kilowatt-hour	= 3.6×10^6 joule = 3.413×10^3 Btu
1 MeV	= 1.6×10^{-13} joule
Energy of fission of 1 atom of ^{235}U	= 199 MeV = 3.2×10^{-11} joule
Energy equivalent of 1 ton of TNT	= 4.2×10^9 joule
Energy of fission of 1 kilogram of ^{235}U	= 20 kilotons of TNT
Hydrogen fusion:	$D + T \rightarrow {}_2\text{He}^4 + n + 17.6 \text{ MeV}$
Energy equivalent of 1 gram of matter	= 9×10^{13} joule
High heat value of 1 ton of coal	= 26×10^6 Btu
High heat value of 1 cord of red oak	= 30×10^6 Btu
High heat value of 100 gallons of fuel oil	= 15×10^6 Btu
High heat value of 20 000 cu ft natural gas	= 20×10^6 Btu
US energy consumption	= 10^{20} joule yr^{-1} (proj. 1970–2000)
Earth's daily receipt of solar energy	= 1.49×10^{22} joule = 4.2×10^{12} Mwh
Earth's rotational energy	= 2.2×10^{29} joule
Earth's total heat content	= 3×10^{31} joule
1 D-cell flashlight battery	= 10^4 watt-s = 10^4 joule

Prefixes and symbols

(used with SI units to indicate decimal multiples and submultiples)

Multiples			Submultiples		
Factor	Prefix	Symbol	Factor	Prefix	Symbol
10^{18}	exa	E	10^{-1}	deci	d
10^{15}	peta	P	10^{-2}	centi	c
10^{12}	tera	T	10^{-3}	milli	m
10^9	giga	G	10^{-6}	micro	μ
10^6	mega	M	10^{-9}	nano	n
10^3	kilo	k	10^{-12}	pico	p
10^2	hecto	h	10^{-15}	femto	f
10	deca	da	10^{-18}	atto	a

Periodic table of the elements

PERIODIC TABLE OF THE ELEMENTS

GROUP IA		GROUP IIA		GROUP IIIB		GROUP IVB		GROUP VB		GROUP VIB		GROUP VIIB		GROUP VIIIB		GROUP VIII	
1 100797 -252.7 -259.2 0.071 H HYDROGEN		3 6.939 1330 180.5 0.53 Li LITHIUM	4 9.0122 2770 1277 185 Be BERYLLIUM														
11 22.9898 892 87.8 0.97 Na SODIUM		12 24.312 1107 650 1.74 Mg MAGNESIUM															
19 39.102 760 63.7 0.86 K POTASSIUM	20 40.08 1440 838 1.55 Ca CALCIUM	21 44.956 2730 1539 3.0 Sc SCANDIUM	22 47.88 3260 1858 4.5 Ti TITANIUM	23 50.942 3450 1800 6.1 V VANADIUM	24 51.996 2865 1875 7.19 Cr CHROMIUM	25 54.938 2150 1245 7.43 Mn MANGANESE	26 55.847 3000 1536 7.88 Fe IRON	27 58.933 2900 1495 8.9 Co COBALT	28 58.71 2730 1453 8.9 Ni NICKEL								
37 85.47 688 36.9 1.33 Rb RUBIDIUM	38 87.62 1380 768 2.6 Sr STRONTIUM	39 88.905 2927 1509 4.47 Y YTRIUM	40 91.22 3580 1852 6.49 Zr ZIRCONIUM	41 92.906 3500 2488 8.4 Nb NIOBIUM	42 95.94 3350 2610 10.2 Mo MOLYBDENUM	43 (98) 2140 11.5 Tc TECHNETIUM	44 101.07 4900 2500 12.2 Ru RUTHENIUM	45 102.905 4500 1968 12.4 Rh RHODIUM	46 106.4 3860 1552 12.0 Pd PALLADIUM								
55 132.905 690 78.7 1.80 Cs CESIUM	56 137.34 1640 7.4 3.5 Ba BARIUM	57 138.91 3470 920 6.17 La LANTHANUM	72 178.49 3400 2222 13.1 Hf HAFNIUM	73 180.948 5425 2996 16.6 Ta TANTALUM	74 183.85 5930 3410 19.3 W WOLFRAM	75 186.2 5900 3180 21.0 Re RHENIUM	76 190.2 5900 3000 22.6 Os OSMIUM	77 192.2 5300 2454 22.5 Ir IRIDIUM	78 195.09 4530 1769 21.4 Pt PLATINUM								
87 (223) 127 Fr FRANCIUM	88 (226) 700 5.0 Ra RADIUM	89 (227) 1090 Ac ACTINIUM	104														
58 140.12 3446 795 5.57 Ce CERIUM	59 140.907 3127 835 8.77 Pr PRASEODYMIUM	60 144.24 3027 1024 7.00 Nd NEODYMIUM	61 (147) (1027) Pm PROMETHIUM	62 150.35 1900 972 7.54 Sm SAMARIUM	63 151.96 1459 826 5.26 Eu EUROPIUM	64 157.25 3000 1317 7.69 Gd GADOLINIUM											
90 232.038 3850 1730 11.7 Th THORIUM	91 (231) (1830) 15.4 Pa PROTACTINIUM	92 238.03 3616 1132 19.07 U URANIUM	93 (237) 637 19.5 Np NEPTUNIUM	94 (242) 3735 840 Pu PLUTONIUM	95 (243) 117 Am AMERICIUM	96 (247) Cm CURIUM											

VALUES FOR GASEOUS ELEMENTS ARE FOR LIQUIDS AT THE BOILING POINT.
 OUTLINE - SYNTHETICALLY PREPARED.

KEY

ATOMIC NUMBER	30	65.37	ATOMIC WEIGHT
BOILING POINT, °C	906		
MELTING POINT, °C	419.5	Zn	SYMBOL
DENSITY (g/ml)	7.14	ZINC	NAME

Periodic table of the elements (cont.)

		IIIA		IVA		VA		VIA		VIIA		VIII			
		5 10.811		6 12.011 5		7 14.0067		8 15.9994		9 18.9984		10 20.183			
		(2030) 2.34 B		4630 3727.9 2.26 C		195.8 210 9.8 N		115 216.8 1.4 O		-188.7 -79.6 90.5 F		-246 -248.6 120 Ne			
		BORON		CARBON		NITROGEN		OXYGEN		FLUORINE		NEON			
		13 26.9815		14 28.086		15 30.9738		16 32.064		17 35.453		18 39.948			
		2450 860 2.70 Al		7680 1410 2.33 Si		280.4 44.24 1.824 P		444.6 19.0 2.07 S		-34.7 -101.0 -3.5 Cl		-185.8 -189.4 -1.9 Ar			
		ALUMINUM		SILICON		PHOSPHORUS		SULFUR		CHLORINE		ARGON			
IB		IIB													
29 63.54		30 65.37		31 69.72		32 72.59		33 74.922		34 78.96		35 79.909		36 83.80	
2595 1093 8.96 Cu		906 419.5 7.14 Zn		2237 29.8 5.91 Ga		2830 937.4 5.32 Ge		513 8.7 5.72 As		685 217 4.79 Se		58 -0.2 3.12 Br		152 -157.3 2.6 Kr	
COPPER		ZINC		GALLIUM		GERMANIUM		ARSENIC		SELENIUM		BROMINE		KRYPTON	
47 107.870		48 112.40		49 114.82		50 118.69		51 121.75		52 127.60		53 126.904		54 131.30	
2210 1660.8 10.5 Ag		763 320.9 8.85 Cd		7000 156.2 7.31 In		2270 231.9 7.30 Sn		1380 530.5 6.62 Sb		989.8 448.5 6.24 Te		183 113.7 4.94 I		-108.0 -119 3.06 Xe	
SILVER		CADMIUM		INDIUM		TIN		ANTIMONY		TELLURIUM		IODINE		XENON	
79 196.967		80 200.59		81 204.37		82 207.19		83 208.980		84 210		85 (210)		86 (222)	
2970 1663 19.3 Au		357 -38.4 13.6 Hg		1457 303 11.85 Tl		1725 327.4 11.4 Pb		1560 271.3 9.8 Bi		254 (9.2) Po		252 At		211 Rn	
GOLD		MERCURY		THALLIUM		LEAD		BISMUTH		POLONIUM		ASTATINE		RADON	
65 158.924		66 162.50		67 164.930		68 167.26		69 168.934		70 173.04		71 174.97			
2800 1356 8.27 Tb		2600 1407 8.54 Dy		2600 1461 8.80 Ho		2900 1497 9.05 Er		1727 1945 9.33 Tm		1427 623 6.96 Yb		3527 1632 9.84 Lu			
TERBIUM		DYSPROSIUM		HOLMIUM		ERBIUM		THULIUM		YTERBIUM		LUTETIUM			
97 (247)		98 (249)		99 (251)		100 (253)		101 (256)		102 (258)		103 (259)			
Bk		Cf		Es		Fm		Md		No		Lw			
BERKELIUM		CALIFORNIUM		EINSTEINIUM		FERMIUM		MENDELEVIUM		NOBELIUM		LAWRENCIUM			

Greek alphabet

A	α	Alpha	N	ν	Nu
B	β	Beta	Ξ	ξ	Xi
Γ	γ	Gamma	O	o	Omicron
Δ	δ	Delta	Π	π	Pi
E	ϵ	Epsilon	P	ρ	Rho
Z	ζ	Zeta	Σ	σ	Sigma
H	η	Eta	T	τ	Tau
Θ	θ	Theta	Υ	υ	Upsilon
I	ι	Iota	Φ	ϕ	Phi
K	κ	Kappa	X	χ	Chi
Λ	λ	Lambda	Ψ	ψ	Psi
M	μ	Mu	Ω	ω	Omega

Bibliography

Reviews of Particle Properties, *Rev. Mod. Phys.*, **52**, 1980.

Handbook of Chemistry and Physics, Weast, R. C., ed., CRC Press, Inc.

Landolt-Börnstein: Zahlenwerte und Funktionen aus Physik, Chemie,

Astronomie, Geophysik, und Technik, Springer-Verlag.

International Critical Tables of Numerical Data, Physics, Chemistry, and Technology, McGraw-Hill Book Company.

Chapter 2

Astronomy and astrophysics

The Solar System	25
<i>The Sun</i>	25
<i>Solar System elemental abundances</i>	30
<i>The planets (physical elements)</i>	31
<i>The planets (orbital elements)</i>	32
<i>Natural satellites in the Solar System</i>	33
<i>Short period comets</i>	36
<i>Principal meteor streams</i>	37
Stars	38
<i>Star charts</i>	38
<i>Constellations</i>	40
<i>The brightest stars</i>	44
<i>The 100 visually brightest stars</i>	45
<i>Stars within 5 pc</i>	50
<i>Nearest stars</i>	52
<i>Stars of large proper motion</i>	53
<i>The 21 fastest pulsars</i>	54
<i>Pulsars</i>	56
<i>Binary radio pulsars</i>	58
<i>OB associations</i>	59
<i>Orbital elements of binary stars</i>	61
<i>Classification of variable stars</i>	62
<i>Galactic supernova remnants</i>	64
<i>Henry Draper spectral classification</i>	64
<i>Spectral type and luminosity class</i>	65
<i>Hertzsprung–Russell diagram</i>	66
<i>Infrared and visible spectral features</i>	67
<i>The main sequence</i>	68

<i>Giant stars</i>	70
<i>Classification and absolute magnitude of stars</i>	71
<i>Stellar mass, luminosity, radius and density</i>	72
<i>Present-day mass function</i>	74
<i>Star number densities</i>	76
<i>Relative number of stars</i>	78
<i>Integrated star light</i>	78
<i>Mean star density vs visual magnitude</i>	79
<i>Star counts</i>	80
<i>Luminosity functions</i>	80
<i>Parameters of the interstellar gas</i>	81
Galaxies	82
<i>Properties of the galaxy</i>	82
<i>The local group</i>	83
<i>Hubble's classification</i>	84
<i>Nearby extragalactic objects</i>	84
<i>Selected brighter galaxies</i>	85
<i>Named galaxies</i>	86
<i>Representative active galactic nuclei</i>	87
<i>Objects with large redshifts</i>	90
<i>Prominent clusters of galaxies</i>	91
Energy content of constituents of space	93
Mass–radius–density data for astronomical objects	94
The Messier catalog	95
Astronomical photometry	98
<i>Standard photometric systems</i>	100
<i>Interstellar reddening</i>	103
<i>Absolute magnitude</i>	105
<i>Moon, night sky, sun, and planetary brightness</i>	105
Spherical astronomy	107
<i>Time</i>	107
<i>The celestial sphere</i>	109
<i>Celestial coordinates</i>	110
<i>The Zodiac</i>	110
<i>Astronomical coordinate transformations</i>	111
<i>Approximate reduction of astronomical coordinates</i>	113
Major ground-based astronomical facilities	114
The Hubble Space Telescope	120
<i>Description of the Hubble Space Telescope</i>	121
Glossary of astronomical terms	124
Bibliography	140




The Solar System

The Sun


Mass	1.9891×10^{33} g
Radius	6.9599×10^{10} cm
Magnetic field strengths (typical):	
Sunspots	3000 G
Polar field	1 G
Bright, chromospheric network	25 G
Ephemeral (unipolar) active regions	20 G
Chromospheric plages	200 G
Prominences	10–100 G
Mean density	1.410 g cm^{-3}
Gravity at surface	$2.7398 \times 10^4 \text{ cm s}^{-2}$
Mean distance from Earth	1 AU = 1.49598×10^{13} cm
Solar constant (1980)	$0.1368 \text{ watts cm}^{-2}$
M_{ν}	+4.83
m_{ν}	−26.74
M_{bol}	+4.75
m_{bol}	−26.82
Effective temperature	5770 K
Luminosity	$3.826 \times 10^{33} \text{ erg s}^{-1}$
Spectral type	G2 V
Inclination of equator to ecliptic	7° 15'
Sidereal rotation (func. of lat.)	$14.44 - 3.0 \sin^2 \phi$ per day
Synodic rotation (func. of lat.)	$13.45 - 3.0 \sin^2 \phi$ per day
Central density	$140\text{--}180 \text{ g cm}^{-3}$
Central temperature	$(14.9\text{--}15.7) \times 10^6 \text{ K}$
Surface area	$6.087 \times 10^{22} \text{ cm}^2$
Volume	$1.412 \times 10^{33} \text{ cm}^3$
Moment of inertia	$5.7 \times 10^{53} \text{ g cm}^2$
Specific surface emission	$6.329 \times 10^{10} \text{ erg cm}^{-2} \text{ s}^{-1}$
Specific mean energy production	$1.937 \text{ erg g}^{-1} \text{ s}^{-1}$
Escape velocity at surface	$6.177 \times 10^7 \text{ cm s}^{-1}$

The Sun (cont.)

Annual mean sunspot number, AD 1610-1975. (From Eddy, J. in *The Solar Output and its Variation*, O. White, ed., Colorado Association University Press, 1977.)




Yearly mean sunspot numbers 1945-87. (US Department of Commerce.)




The Sun (cont.)

Temperature and density as a function of distance from the solar surface. (Courtesy of G. Withbroe, Harvard/Smithsonian Center for Astrophysics.)




Solar temperature and electron density. (Adapted from Carrigan, A. L. & Skrivanek, *Aerospace Environment*, Air Force Cambridge Research Laboratories, Massachusetts, 1974.)




The Sun (cont.)

Solar spectral irradiance. (Adapted from Carrigan, A. L. & Skrivanek, *Aerospace Environment*, Air Force Cambridge Research Laboratories, Massachusetts, 1974.)




The solar spectral irradiance at 1 AU between 10 and 300 Å. Three states of solar activity are shown for the region 10-30 Å. The vertical extent of the shaded areas is representative of the variability of the spectral irradiance for changing solar conditions. (Adapted from Manson, J. E. in *The Solar Output and Its Variation*, O. R. White, ed., Colorado Associated University Press, Boulder, 1977.)



The Sun (cont.)

Solar EUV flux distribution incident on Earth's atmosphere (moderately active, non-flaring sun). (Adapted from Carrigan, A. L. & Skrivanek, *Aerospace Environment*, Air Force Cambridge Research Laboratories, Massachusetts, 1974.)



Solar System elemental abundances (normalized to Si = 10⁶)

Element	Abundance	Element	Abundance	Element	Abundance
1 H	2.66×10^{10}	29 Cu	540	58 Ce	1.2
2 He	1.8×10^9	30 Zn	1260	59 Pr	0.18
3 Li	60	31 Ga	38	60 Nd	0.79
4 Be	1.2	32 Ge	117	62 Sm	0.24
5 B	9	33 As	6.2	63 Eu	0.094
6 C	1.11×10^7	34 Se	67	64 Gd	0.42
7 N	2.31×10^6	35 Br	9.2	65 Tb	0.076
8 O	1.84×10^7	36 Kr	41.3	66 Dy	0.37
9 F	780	37 Rb	6.1	67 Ho	0.092
10 Ne	2.6×10^6	38 Sr	22.9	68 Er	0.23
11 Na	6.0×10^4	39 Y	4.8	69 Tm	0.035
12 Mg	1.06×10^6	40 Zr	12	70 Yb	0.20
13 Al	8.5×10^4	41 Nb	0.9	71 Lu	0.035
14 Si	1.00×10^6	42 Mo	4.0	72 Hf	0.17
15 P	6500	44 Ru	1.9	73 Ta	0.020
16 S	5.0×10^5	45 Rh	0.40	74 W	0.15
17 Cl	4740	46 Pd	1.3	75 Re	0.051
18 Ar	1.06×10^5	47 Ag	0.46	76 Os	0.69
19 K	3500	48 Cd	1.55	77 Ir	0.72
20 Ca	6.25×10^4	49 In	0.19	78 Pt	1.41
21 Sc	31	50 Sn	3.7	79 Au	0.21
22 Ti	2400	51 Sb	0.31	80 Hg	0.42
23 V	254	52 Te	6.5	81 Tl	0.19
24 Cr	1.27×10^4	53 I	1.27	82 Pb	2.6
25 Mn	9300	54 Xe	5.84	83 Bi	0.14
26 Fe	9.0×10^5	55 Cs	0.39	90 Th	0.045
27 Co	2200	56 Ba	4.8	92 U	0.027
28 Ni	4.78×10^4	57 La	0.37		

The practical mass fractions of hydrogen, helium, and the heavier elements are: $X = 0.77$, $Y = 0.21$, $Z = 0.02$. (Adapted from Cameron, A. G. W. in "Essays in Nuclear Astrophysics", Cambridge University Press, 1981).

The planets (physical elements)

Planet	Mass (Earth = 1)	Equatorial radius (Earth = 1)	Mean density (g cm^{-3})	Equatorial surface gravity (Earth = 1)	Equatorial escape velocity (km s^{-1})	Sidereal rotation period	Inclination of equator to orbit (°)	No. of satellites
Mercury	0.0558	0.382	5.42	0.387	4.3	58.65 d	0	0
Venus	0.8150	0.949	5.25	0.879	10.3	243.01 (retro)d	-2	0
Earth	1.0000	1.000	5.52	1.000	11.2	23.9345 h	23.45	1
Moon	0.01230	0.2725	3.34	0.166	2.38	27.322 d	6.68	2
Mars	0.1074	0.532	3.94	0.380	5.0	24.6229 h	23.98	15
Jupiter	317.893	11.27	1.314	2.339	59.5	9.841 h	3.08	17
Saturn	95.147	9.44	0.69	0.925	35.6	10.233 h	29	5
Uranus	14.54	4.10	(1.19)	0.794	21.22	15.5 h	97.92	2
Neptune	17.23	3.88	1.66	1.125	23.6	15.8 h	28.8	1
Pluto	(0.0017)	(0.17)	(0.6-1.7)	(0.44)	(5.3)	6.3874 d	(≥ 50)	

Values in parentheses are uncertain by more than $10^0\%$.

For the Earth: equatorial radius = 6378 km, mass = 5.976×10^{27} g,
equatorial surface gravity = 978 cm s^{-2} (uncorrected for centrifugal force),

magnetic dipole moment = 8.00×10^{25} Gauss cm^3 ,

Earth-Moon mean distance = 3.84401×10^5 km.

The planets (orbital elements)

Planet	Mean distance from Sun (AU)	Sidereal period (d)	Synodic period (d)	Mean orbital velocity (km s ⁻¹)	Eccentricity	Inclination to the ecliptic (°)
Mercury	0.387 099	87.969	115.88	47.89	0.2056	7.00
Venus	0.723 332	224.701	583.92	35.03	0.0068	3.39
Earth	1.000 000	365.256	—	29.79	0.0167	—
Mars	1.523 691	686.980	779.94	24.13	0.0934	1.85
Jupiter	5.202 803	4 332.589	398.88	13.06	0.0485	1.30
Saturn	9.538 84	10 759.22	378.09	9.64	0.0556	2.49
Uranus	19.1819	30 685.4	369.66	6.81	0.0472	0.77
Neptune	30.0578	60 189.0	367.49	5.43	0.0086	1.77
Pluto	39.44	90 465.0	366.73	4.74	0.250	17.2

1 AU = $1.495\,979 \times 10^{13}$ cm. The sidereal period is the period of revolution measured with respect to the fixed stars; the synodic period is the interval between successive oppositions of a superior planet or successive inferior conjunctions of an inferior planet.

Natural satellites in the Solar System

Number	Name	Year found	Orbital radius (10 ³ km)	Orbital period (d)	Eccentricity ()	Inclination ()	Radius (km)	Magnitude† (m _v)
SATELLITE OF EARTH								
	Moon	—	384.4	27.322	0.055	5.1	1738	-12.7
SATELLITES OF MARS (2)								
MI	Phobos	1877	9.38	0.319	0.015	1.02	14 11 9	11.6
MII	Deimos	1877	23.46	1.262	0.001	1.82	(8) 6 (5)	12.7
SATELLITES OF JUPITER (16)								
(J16)	(Metis)	1979-80	127.96	0.295	(0)	(0)	(20)	17.4
(J15)	(Adrastea)	1979	128.98	0.298	(0)	(0)	12 10 8	18.9
JV	Amalthea	1892	181.3	0.498	0.003	0.45	135 85 75	14.1
JXIV	Thebe	1979-80	221.9	0.675	0.013	(0.9)	55 ? 45	(15.5)
J1	Io	1610	421.6	1.769	0.004	0.04	1815	5.0
JII	Europa	1610	670.9	3.551	0.009	0.47	1569	5.3
JIII	Ganymede	1610	1070	7.155	0.002	0.21	2631	4.6
JIV	Callisto	1610	1880	16.689	0.007	0.51	2400	5.6
JXIII	Leda	1974	11 094	238.7	0.148	26.1	(5)	20.2
JVI	Himalia	1904 05	11 480	250.6	0.158	27.6	(90)	14.8
JX	Lysithea	1938	11 720	259.2	0.107	29.0	(10)	18.4
JVII	Elara	1904-05	11 737	259.7	0.207	24.8	(40)	16.7
JXII	Ananke	1951	21 200	631	0.17	147	(10)	18.9
JXI	Carme	1938	22 600	692	0.21	164	(15)	18.0
JVIII	Pasiphae	1908	23 500	735	0.38	145	(20)	17.7
JIX	Sinope	1914	23 700	758	0.28	153	(15)	18.3

† At mean opposition.

Number	Name	Year found	Orbital radius (10 ³ km)	Orbital period (d)	Eccentricity	Inclination (°)	Radius (km)	Magnitude (m _v)
SATELLITES OF SATURN (17)								
(SXV)	(Atlas)	1980	137.67	0.602	0.002	0.3	(19) ? (13)	18
1980 S 27		1980	139.35	0.613	0.004	0.0	70 (50) (37)	16.5
1980 S 26		1980	141.70	0.629	0.004	0.1	(55) (42) (33)	16
SX	Janus	1966	151.47	0.695	0.007	0.1	110 95 80	14.5
SXI	Epimetheus	1966	151.42	0.694	0.009	0.3	(70) (57) (50)	15.5
SI	Mimas	1789	185.54	0.942	0.020	1.52	196	12.9
SII	Enceladus	1789	238.04	1.370	0.004	0.02	250	11.8
SIII	Tethys	1684	294.67	1.888	0.000	1.86	530	10.3
SXIII	Telesto	1980	294.67	1.888	?	?	? (12) (11)	19
SXIV	Calypso	1980	294.67	1.888	?	?	(15) (12) (8)	18.5
SIV	Dione	1684	377.42	2.737	0.002	0.02	560	10.4
1980 S 6		1980	377.42	2.737	0.005	0.2	(18) ? (<15)	18.5
SV	Rhea	1672	527.04	4.518	0.001	0.35	765	9.7
SVI	Titan	1655	1221.86	15.945	0.029	0.33	2575	8.4
SVII	Hyperion	1848	1481.1	21.277	0.104	0.43	175 117 (100)	14.2
SVIII	Iapetus	1671	3561.3	79.331	0.028	(7.52)	730	10.2-11.9
SIX	Phoebe	1898	12 954	550.4	0.163	175	110	16.5
SATELLITES OF URANUS (5)								
UV	Miranda	1948	129.4	1.414	0.027	4.22	(200)	16.5
UI	Ariel	1851	191.0	2.520	0.003	0.31	665	14.4
UII	Umbriel	1851	266.3	4.144	0.005	0.36	555	15.3
UIII	Titania	1787	435.9	8.706	0.002	0.14	800	14
UIV	Oberon	1787	583.5	13.463	0.001	0.10	815	14.2

SATELLITES OF NEPTUNE (2)								
NI	Triton	1846	355.3	5.877	<0.001	159	(1750)	13.6
NII	Nereid	1949	5510	360.21	0.75	27.6	(200)	18.7
SATELLITE OF PLUTO								
PI	Charon	1978	19.7	6.387	(0)	(0)	(580)	17

† At mean opposition.

Parenthetic names and satellite numbers await approval by the International Astronomical Union. Parenthetic values are uncertain by at least 10%. *Inclinations* and *eccentricities* are the total values; 'forced' components are important for the eccentricities of the satellites Io, Europa, Enceladus, and Hyperion. Inclinations are with respect to planetary equators for inner satellites and to orbital planes for outer ones. *Compound radii* are the values for the 'best-fit' triaxial ellipsoid. *Magnitudes* are the visual mean opposition values. (Reproduced by permission from *Sky and Telescope* astronomy magazine, November 1983 issue, page 405.)

Short period comets

Comet name	Perihelion epoch	Perihelion longitude, $\bar{\omega}$ ($^{\circ}$)	Longitude of ascending node, Ω ($^{\circ}$)	Period, T_p (yr)	Semi-major axis of orbit, a (AU)	Eccentricity of orbit, e	Inclination of orbit, i ($^{\circ}$)	Perihelion distance (AU)
Encke	1974.32	160.1	334.2	3.30	2.209	0.847	12.0	0.338
Temple 2	1972.87	310.2	119.3	5.26	3.024	0.549	12.5	1.364
Haneda-Campos	1978.77	12.016	131.7000	5.37	3.066	0.64152	5.805	1.099
Schwassmann-Wachmann 2	1974.70	123.3	126.0	6.51	3.489	0.386	3.7	2.142
Borrelly	1974.36	67.8	75.1	6.76	3.576	0.632	30.2	1.316
Whipple	1970.77	18.2	188.4	7.47	3.821	0.351	10.2	2.480
Oterma	1958.44	150.0	155.1	7.88	3.958	0.144	4.0	3.388
Schaumasse	1960.29	138.1	86.2	8.18	4.054	0.705	12.0	1.196
Comas Sola	1969.83	102.9	62.8	8.55	4.182	0.577	13.4	1.769
Schwassmann-Wachmann 1	1974.12	334.1	319.6	15.03	6.087	0.105	9.7	5.448
Neujmin 1	1966.94	334.0	347.2	17.93	6.858	0.775	15.0	1.543
Crommelin	1956.82	86.4	250.4	27.89	9.173	0.919	28.9	0.743
Olbers	1956.46	150.0	85.4	69.47	16.843	0.930	44.6	1.179
Pons-Brooks	1954.39	94.2	255.2	70.98	17.200	0.955	74.2	0.774
Halley	1986.112	170.0110	58.1540	76.0081	17.9435	0.9673	162.2384	0.587

(Adapted from Duffet-Smith, P., *Practical Astronomy with your Calculator*, Cambridge University Press.)



Principal meteor streams

Stream	Maximum	Normal period of visibility	Radiant		Geocentric velocity km s ⁻¹
			α	δ	
Quadrantids	Jan. 3	Jan. 2-4	231	+49	43
Lyrids	Apr. 23	Apr. 20-22	271	+33	47
η Aquarids	May 4	May 2-7	336	0	64
δ Aquarids	July 30	July 20-Aug. 14	339	-10	41
Perseids	Aug. 12	July 29-Aug. 18	46	+58	60
Draconids	Oct. 10	Oct. 10	265	+54	24
Orionids	Oct. 21	Oct. 17-24	95	+15	66
Taurids	Nov. 4	Oct. 20-Nov. 25	55	+17	30
Leonids	Nov. 16	Nov. 14-19	153	+22	72
Andromedids	Nov. 20	Nov. 15-Dec. 6	13	+55	20
Geminids	Dec. 13	Dec. 8-15	112	+32	36
Ursids	Dec. 22	Dec. 19-23	213	+76	36
Permanent daytime streams					
Arietids	June 8	May 29 June 17	44	+23	39
ξ Perseids	June 9	June 1-15	61	+23	29
β Taurids	June 30	June 23 July 7	86	+19	31



(Adapted from Allen, C. W., *Astrophysical Quantities*, The Athlone Press, 1973.)

Stars

Star charts



Star charts (cont.)



(Adapted from *The Nautical Almanac*)

Constellations

Constellation	Gen. ending	Contr.	Description	α	δ	Area ^(a) (sq. deg.)	Order of size
Andromeda	-dae	And	the chained maiden ^(b)	1 ^h	40° N	722.278	19
Antlia	-liae	Ant	the air pump	10	35° S	238.901	62
Apus	-podis	Aps	the bird of paradise	16	75° S	206.327	67
Aquarius	-rii	Aqr	the water pourer	23	15° S	979.854	10
Aquila	-lae	Aql	the eagle	20	5° N	652.473	22
Ara	-rae	Ara	the altar	17	55° S	237.057	63
Aries	-ietis	Ari	the ram	3	20° N	441.395	39
Auriga	-gae	Aur	the wagoner	6	40° N	657.438	21
Boötes	-tis	Boo	the ploughman	15	30° N	906.831	13
Caelum	-aeli	Cae	the burin	5	40° S	124.865	81
Camelopardalis	-di	Cam	the giraffe	6	70° N	756.828	18
Cancer	-cri	Cnc	the crab	9	20° N	505.872	31
Canes Venatici	-num -corum	CVn	the hunting dogs	13	40° N	465.194	38
Canis Major	-is -ris	CMa	the large dog	7	20° S	380.118	43
Canis Minor	-is -ris	CMi	the small dog	8	5° N	183.367	71
Capricornus	-ni	Cap	the goat	21	20° S	413.947	40
Carina	-nae	Car	the keel	9	60° S	494.184	34
Cassiopeia	-peiae	Cas	Cassiopeia	1	60° N	598.407	25
Centaurus	-ri	Cen	the centaur	13	50° S	1060.422	9
Cepheus	-phei	Cep	King Cepheus	22	70° N	587.787	27
Cetus	-ti	Cet	the sea monster	2	10° S	1231.411	4
Chamaeleon	-ntis	Cha	the chamaeleon	11	80° S	131.592	79
Circinus	-ni	Cir	the pair of compasses	15	60° S	93.353	85
Columba	-bae	Col	the dove	6	35° S	270.184	54
Coma Berenices	-mae -cis	Com	the hair of Berenice	13	20° N	386.475	42

Corona Australis	-nae -lis	CrA	the southern crown	19	40 S	127.696	80
Corona Borealis	-nae -lis	CrB	the northern crown	16	30 N	178.710	73
Corvus	-vi	CrV	the crow	12	20 S	183.801	70
Crater	-eris	CrI	the cup	11	15 S	282.398	53
Crux	-ucis	CrU	the southern cross	12	60 S	68.447	88
Cygnus	-gni	CyG	the swan	21	40 N	803.983	16
Delphinus	-ni	Del	the dolphin	21	10 N	188.549	69
Dorado	-dus	Dor	the swordfish	5	65 S	179.173	72
Draco	-onis	Dra	the dragon	17	65 N	1082.952	8
Equuleus	-lei	Equ	the foal	21	10 N	71.641	87
Eridanus	-ni	Eri	the river	3	20 S	1137.919	6
Fornax	-acis	For	the furnace	3	30 S	397.502	41
Gemini	-norum	Gem	the twins	7	20 N	513.761	30
Grus	-ruis	Gru	the crane	22	45 S	365.513	45
Hercules	-lis	Her	Hercules	17	30 N	1225.148	5
Horologium	-gii	Hor	the clock	3	60 S	248.885	58
Hydra	-drae	Hya	the water snake	10	20 S	1302.844	1
Hydrus	-dri	Hyi	the sea serpent	2	75 S	243.035	61
Indus	-di	Ind	the Indian	21	55 S	294.006	49
Lacerta	-tae	Lac	the lizard	22	45 N	200.688	68
Leo	-onis	Leo	the lion	11	15 N	946.964	12
Leo Minor	-onis -ris	LMi	the small lion	10	35 N	231.956	64
Lepus	-poris	Lep	the hare	6	20 S	290.291	51
Libra	-rae	Lib	the balance	15	15 S	538.052	29
Lupus	-pi	Lup	the wolf	15	45 S	333.683	46
Lynx	-ncis	Lyn	the lynx	8	45 N	545.386	28
Lyra	-rae	Lyr	the lyre	19	40 N	286.476	52
Mensa	-sae	Men	the table ⁽¹⁾	5	80 S	153.484	75
Microscopium	-pii	Mic	the microscope	21	35 S	209.513	66
Monoceros	-rotis	Mon	the unicorn	7	5 S	481.569	35

Constellations (cont.)

Constellation	Gen. ending	Contr.	Description	α	δ	Area ^(a) (sq. deg.)	Order of size
Musca	-cae	Mus	the fly	12	70°S	138.355	77
Norma	-mae	Nor	the square	16	50 S	165.290	74
Octans	-ntis	Oct	the octant	22	85 S	291.045	50
Ophiuchus	-chi	Oph	the serpent bearer	17	0	948.340	11
Orion	-nis	Ori	the hunter	5	5 N	594.120	26
Pavo	-vonis	Pav	the peacock	20	65 S	377.666	44
Pegasus	-si	Peg	the winged horse	22	20 N	1120.794	7
Perseus	-sei	Per	Perseus	3	45 N	614.997	24
Phoenix	-nisis	Phe	the Phoenix	1	50 S	469.319	37
Pictor	-ris	Pic	the easel	6	55 S	246.739	59
Pisces	-cium	Psc	the fishes	1	15 N	889.417	14
Piscis Austrinus	-is -ni	PsA	the southern fish	22	30 S	245.375	60
Puppis	-ppis	Pup	the poop	8	40 S	673.434	20
Pyxis (= Malus)	-xidis	Pyx	the compass	9	30 S	220.833	65
Reticulum	-li	Ret	the net	4	60 S	113.936	82
Sagitta	-tae	Sge	the arrow	20	10 N	79.923	86
Sagittarius	-rii	Sgr	the archer	19	25 S	867.432	15
Scorpius	-pii	Sco	the scorpion	17	40 S	496.783	33
Sculptor	-ris	ScI	the sculptor	0	30 S	474.764	36
Scutum	-ti	ScI	the shield ^(d)	19	10 S	109.114	84
Serpens (Caput and Cauda)	-ntis	Ser	the serpent	18	5 S	636.928	23
Sextans	-ntis	Sex	the sextant	10	0	313.515	47
Taurus	-ri	Tau	the bull	4	15 N	797.249	17
Telescopium	-pii	Tel	the telescope	19	50 S	251.512	57
Triangulum	-li	Tri	the triangle	2	30 N	131.847	78

Triangulum Australe	-li -lis	TriA	the southern triangle	16	65 S	109,978	83
Tucana	-nae	Tuc	the toucan	0	65 S	294,557	48
Ursa Major	-sae -ris	UMa	the great bear	11	50 N	1279,660	3
Ursa Minor	-sae -ris	UMi	the little bear	15	70 N	255,864	56
Vela	-lorum	Vel	the sails	9	50 S	499,649	32
Virgo	-ginis	Vir	the virgin	13	0	1294,428	2
Volans	-ntis	Vol	the flying fish	8	70 S	141,354	76
Vulpecula	-lae	Vul	the little fox	20	25 N	268,165	55

^(a) From *Sky and Telescope*, June 1983.

^(b) Daughter of Cepheus and Cassiopeia.

^(c) After Table Mountain in South Africa.

^(d) Of John Sobieski, the Polish hero.

The brightest stars in the sky (limiting magnitude, $V = 1.35$)

Rank	Name	Designation	V magnitude	α 2000	δ 2000
1	Sirius	α Canis Majoris	-1.46	6 ^h 45 ^m 8.9 ^s	-16° 42' 58"
2	Canopus	α Carinae	-0.72	6 23 57.1	-52 41 44
3	Rigel Kentaurus	α Centauri	-0.27	14 39 36.7	-60 50 02
4	Arcturus	α Boötis	-0.04	14 15 39.6	+19 10 57
5	Vega	α Lyrae	+0.03	18 36 56.2	+38 47 01
6	Capella	α Aurigae	+0.08	5 16 41.3	+45 59 53
7	Rigel	β Orionis	+0.12	5 14 32.2	-8 12 06
8	Procyon	α Canis Minoris	+0.38	7 39 18.1	+5 13 30
9	Achernar	α Eridani	+0.46	1 37 42.9	-57 14 12
10	Betelgeuse	α Orionis	+0.50 (var)	5 55 10.2	+7 24 26
11	Hadar	β Centauri	+0.61	14 03 49.4	-60 22 22
12	Altair	α Aquilae	+0.77	19 50 46.8	+8 52 06
13	Aldebaran	α Tauri	+0.85 (var)	4 35 55.2	+16 30 33
14	Acrux	α Crucis	+0.87	12 26 35.9	-63 05 56
15	Antares	α Scorpii	+0.96 (var)	16 29 24.3	-26 25 55
16	Spica	α Virginis	+0.98	13 25 11.5	-11 09 41
17	Pollux	β Geminorum	+1.14	7 45 18.9	+28 01 34
18	Fomalhaut	α Piscis Austrini	+1.16	22 57 38.9	-29 37 20
19	Deneb	α Cygni	+1.25	20 41 25.8	+45 16 49
20	Mimosa	β Crucis	+1.25	12 47 43.2	-59 41 19
21	Regulus	α Leonis	+1.35	10 08 22.2	+11 58 02

The 100 visually brightest stars (limiting magnitude, $V = 2.59$)

Star name*	α 2000	δ 2000	$\mu(\alpha)$	$\mu(\delta)$	V	$B - V$	M_v	Spec	RV (km s^{-1})	d (pc)	Notes
21 α And	0 ^h 08 ^m 23 ^s .2	+29°05'26"	+0.010	-0.16	2.06	-0.11	0.3	A0 p	-12	22 mn	Alpheratz, m
11 β Cas	0 09 10.6	+59 08 59	+0.068	-0.18	2.27	0.34	1.7	F2 IV	+12	13 ts	Caph, m
α Phe	0 26 17.0	-42 18 22	+0.019	-0.39	2.39	1.09	0.5	K0 III	+75	24 s	(Ankaal, m)
18 α Cas	0 40 30.4	+56 32 15	+0.006	-0.03	2.23	1.17	-0.6	K0 II III	-4	37 s	Schedar, m
16 β Cet	0 43 35.3	-17 59 12	+0.016	+0.04	2.04	1.02	0.4	K0 III	+13	21 ts	Deneb Kait, Diphda
27 γ Cas	0 56 42.4	+60 43 00	+0.003	0.00	2.47	-0.15	-4.4	B0 IV e	-7	240 s	Cih, m, v
43 β And	1 09 43.8	+35 37 14	+0.015	-0.11	2.06	1.58	-0.1	M0 III	0	27 ts	Mirach, m
α Eri	1 37 42.9	-57 14 12	+0.013	-0.03	0.46	-0.16	-1.6	B5 IV	+19	26 s	Achernar
57 γ^1 And	2 03 53.9	+42 19 47	+0.004	-0.05	2.18	1.20	-0.7	K2 III	-12	37 mn	Almach
57 γ^2 And	2 03 54.7	+42 19 51	+0.003	-0.05	5.03			A0 p	-14	37 mn	m
13 α Ari	2 07 10.3	+23 27 45	+0.014	-0.14	2.00	1.15	-0.1	K2 III	-14	26 ts	Hamal
68 ρ Cet	2 19 20.6	-2 58 39	-0.001	-0.23	2.0	1.7	-0.3	Md	+64	29 mn	Mira, m, v
1 α UMi	2 31 50.4	+89 15 51	+0.232	-0.01	2.02	0.60	-4.6	F8 Ib	-17		Polaris, m
92 α Cet	3 02 16.7	+4 05 23	-0.001	-0.07	2.53	1.64	-0.5	M2 III	-26	40 mn	Menkar
26 β Per	3 08 10.1	+40 57 21	0.000	0.00	2.12	-0.05	-0.2	B8 V	+4	29 ts	Algol, m, v
33 α Per	3 24 19.3	+49 51 40	+0.003	-0.02	1.80	0.48	-4.6	F5 Ib	-2	190 s	Mirfak, m
87 α Tau	4 35 55.2	+16 30 33	+0.005	-0.19	0.85	1.54	-0.8	K5 III	+54	21 ts	Aldebaran, m
13 α Aur	5 16 41.3	+45 59 53	+0.008	-0.42	0.08	0.80	-0.5	G8 III	+30	13 t	Capella, m
19 β Ori	5 14 32.2	-8 12 06	0.000	0.00	0.12	-0.03	-7.1	B8 Ia	+21	280 s	Rigel, m
24 γ Ori	5 25 07.8	+6 20 59	-0.001	-0.01	1.64	-0.22	-3.6	B2 III	+18	110 s	Bellatrix, m
112 β Tau	5 26 17.5	+28 36 27	+0.002	-0.17	1.65	-0.13	-1.4	B7 III	+8	40 mx	El Nath, m
34 δ Ori	5 32 00.3	-0 17 57	0.000	0.00	2.23	-0.22		O9.5 II	+16		Mintaka, m
11 α Lep	5 32 43.7	-17 49 20	0.000	0.00	2.58	0.21	-4.7	F0 Ib	+25	290 s	Arneb, m

* See page 49 for explanation of column headings.

The 100 visually brightest stars (cont.)

Star name*	α 2000	δ 2000	$\mu(\alpha)$	$\mu(\delta)$	V	$B - V$	M_v	Spec	RV (km s^{-1})	d (pc)	Notes
46 ϵ Ori	5 ^h 36 ^m 12 ^s .7	-1° 12' 07"	0.000	0.00	1.70	-0.19	-6.1	B0 Ia	+26	370 s	Alnilam, m
50 ζ Ori	5 40 45.5	-1 56 34	0.000	0.00	1.77	-0.21	-5.9	O9.5 Ib	+18	340 s	Alnitak, m
53 κ Ori	5 47 45.3	-9 40 11	0.000	-0.01	2.06	-0.17	-6.7	B0.5 Ia	+21	560 s	Saiph
58 α Ori	5 55 10.2	+7 24 26	+0.002	+0.01	0.50	1.85	-6.0	M2 Iab	+21	200 s	Betelgeuse, m
34 β Aur	5 59 31.7	+44 56 51	-0.005	0.00	1.90	0.03	0.2	A2 IV	-18	22 ts	Menkalinan, m
2 β CMa	6 22 41.9	-17 57 22	-0.001	0.00	1.98	-0.23	-4.7	B1 II-III	+34	220 s	Mirzam, m
α Car	6 23 57.1	-52 41 44	+0.003	+0.02	-0.72	0.15	-8.5	F0 Ia	+21	360 s	Canopus
24 γ Gem	6 37 42.7	+16 23 57	+0.003	-0.04	1.93	0.00	-0.1	A0 IV	-13	26 ts	Alhena, m
9 α CMa	6 45 08.9	-16 42 58	-0.038	-1.21	-1.46	0.01	1.4	A1 V	-8	2.7 t	Sirius, m
21 ϵ CMa	6 58 37.5	-28 58 20	0.000	0.00	1.50	-0.21	-4.4	B2 II	+27	150 s	Adhara, m
25 δ CMa	7 08 23.4	-26 23 36	-0.001	0.00	1.86	0.65	-8.0	F8 Ia	+34	940 s	Wezen
31 η CMa	7 24 05.6	-29 18 11	-0.001	0.00	2.44	-0.07	-7.0	B5 Ia	+41	760 s	Aludra, m
66 α Gem	7 34 35.9	+31 53 18	-0.013	-0.10	1.58	0.04	0.8	A1 V	+4	14 ts	Castor, m
10 α CMi	7 39 18.1	+5 13 30	-0.047	-1.03	0.38	0.42	2.7	F5 IV	-3	3.5 t	Procyon, m
78 β Gem	7 45 18.9	+28 01 34	-0.047	-0.05	1.14	1.00	0.9	K0 III	+3	11 ts	Pollux, m
ζ Pup	8 03 35.0	-40 00 12	-0.003	+0.01	2.25	-0.26		O5.8	-24		Naos
γ Vel	8 09 31.9	-47 20 12	-0.001	0.00	1.78	-0.22		WC7	+35		m
ϵ Car	8 22 30.8	-59 30 34	-0.003	+0.01	1.86	1.27	-2.1	K0 II	+12	63 s	(Avior)
δ Vel	8 44 42.2	-54 42 30	+0.003	-0.08	1.96	0.04	0.3	A0 V	+2	21 ts	m
λ Vel	9 07 59.7	-43 25 57	-0.002	+0.01	2.21	1.66	-3.7	K5 Ib	+18	150 ts	Suhail
β Car	9 13 12.2	-69 43 02	-0.029	+0.10	1.68	0.00	-0.4	A0 III	-5	26 s	Miaplacidus
ι Car	9 17 05.4	-59 16 31	-0.002	0.00	2.25	0.18	-4.7	F0 Ib	+13	250 s	Aspidiske, Scutulum
κ Vel	9 22 06.8	-55 00 38	-0.001	+0.01	2.50	-0.18	-2.9	B2 IV	+22	120 s	m
30 α Hya	9 27 35.2	-8 39 31	-0.001	+0.03	1.98	1.44	-0.1	K3 III	-4	26 mn	Alphard

32 α Leo	10 08 22.2	+11 58 02	-0.017	0.00	1.35	-0.11	-0.7	B7 V	+4	26 ts	Regulus, m
41 γ^1 Leo	10 19 58.3	+19 50 30	+0.022	-0.15	2.28	1.08	0.2	K0 III	-37		Algeba, m
41 γ^2 Leo	10 19 58.6	+19 50 25	+0.022	-0.17	3.58			G7 III	-36		m
48 β UMa	11 01 50.4	+56 22 56	+0.010	+0.03	2.37	-0.02	1.0	A1 V	-12	19 ts	Merak
50 α UMa	11 03 43.6	+61 45 03	-0.017	-0.07	1.79	1.07	0.0	K0 III	-9	23 ts	Dubhe, m
68 δ Leo	11 14 06.4	+20 31 25	+0.010	-0.14	2.56	0.12	1.5	A4 V	-21	16 ts	Zosma, m
94 β Leo	11 49 03.5	+14 34 19	-0.034	-0.12	2.14	0.09	1.7	A3 V	0	12 mx	Denebolan, m
64 γ UMa	11 53 49.7	+53 41 41	+0.011	+0.01	2.44	0.00	0.6	A0 V	-13	23 s	Phecda
4 γ Crv	12 15 48.3	-17 32 31	-0.011	+0.02	2.59	-0.11	-1.2	B8 III	-4	57 s	Gienah
α^1 Cru	12 26 35.9	-63 05 56	-0.004	-0.02	1.41	0.1	-3.8	B1 IV	-11	110 s	Acrux, m
α^2 Cru	12 26 36.5	-62 05 58	-0.005	-0.02	1.88		-3.3	B3 n	-1	110 s	m
γ Cru	12 31 09.9	-57 06 47	+0.003	-0.27	1.63	1.59	-0.5	M3 III	+21	27 s	(Gaerux), m
γ Cen	12 41 30.9	-48 57 34	-0.019	-0.01	2.17	-0.01	-0.5	A0 III	-8	34 s	Muhlifain, m
β Cru	12 47 43.2	-59 41 19	-0.005	-0.02	1.25	-0.23	-4.3	B0 III	+20	130 mx	Mimesa
77 ϵ UMa	12 54 01.7	+55 57 35	+0.013	-0.01	1.77	-0.02	0.4	A0 p	-9	19 mn	Alioth
79 ζ UMa	13 23 55.5	+54 55 31	+0.014	-0.02	2.27	0.02	1.0	A2 V	-9	18 ts	Mizar, m
67 α Vir	13 25 11.5	-11 09 41	-0.003	-0.03	0.98	-0.23	-3.5	B1 V	+1	79 s	Spica, m, v
ϵ Cen	13 39 53.2	-53 27 58	-0.002	-0.02	2.30	-0.22	-3.6	B1 V	+6	150 s	m
85 η UMa	13 47 32.3	+49 18 48	-0.013	-0.01	1.86	-0.19	-0.7	B3 V	-11	33 mx	Alkaid
ζ Cen	13 55 32.3	-47 17 17	-0.006	-0.04	2.55	-0.22	-2.7	B2 IV	+7	110 mx	m
β Cen	14 03 49.4	-60 22 22	-0.003	-0.02	0.61	-0.24	-5.1	B1 II		140 s	Hadar, m
5 θ Cen	14 06 40.9	-36 22 12	-0.043	-0.52	2.06	1.01	1.3	K0 III-IV	+1	14 ts	(Menkent)
16 α Boo	14 15 39.6	+19 10 57	-0.077	-2.00	-0.04	1.23	-0.2	K2 III p	-5	11 t	Arcturus
η Cen	14 35 30.3	-42 09 28	-0.003	-0.04	2.31	-0.19	-2.9	B3 III	0	110 s	
α^2 Cen	14 39 35.4	-60 50 13	-0.494	+0.69	1.39		5.8	K1 V	-21	1.3 t	m
α^1 Cen	14 39 36.7	-60 50 02	-0.494	+0.69	0.00	0.68	4.4	G2 V	-25	1.3 t	Rigel Kent, m
α Lup	14 41 55.7	-47 23 17	-0.002	-0.02	2.30	-0.20	-4.3	B1 III	+7	210 s	m
36 ϵ Boo	14 44 59.1	+27 04 27	-0.004	+0.02	2.37	0.97	-0.9	K0 II-III	-17	46 s	Izar
7 β UMi	14 50 42.2	+74 09 19	-0.009	+0.01	2.08	1.47	-0.2	K4 III	+17	29 ts	Kochab

Star name	α 2000	δ 2000	$\mu(\alpha)$	$\mu(\delta)$	V	$B - V$	M_V	Spec	RV (km s^{-1})	d (pc)	Notes
5 α CrB	15 ^h 34 ^m 41 ^s .2	+26° 42' 53"	+0.009	-0.09	2.23	-0.02	0.3	A0 V	+2	24 ts	Alphecca, m
7 δ Sco	16 00 19.9	-22 37 18	-0.001	-0.03	2.32	-0.12	-3.8	B0 V	-14	170 s	Dzuba
8 β^1 Sco	16 05 26.1	-19 48 19	0.000	-0.02	2.64	-0.07	-4.3	B0.5 V	-7	250 s	Acraab, m
8 β^2 Sco	16 05 26.4	-19 48 07	-0.001	-0.02	4.92	-0.02	-2.5	B2 V	-5	250 s	
21 α Sco	16 29 24.3	-26 25 55	0.000	-0.02	0.96	1.83	-4.0	M1 Ib	-3	100 s	Antares, m, v
13 ζ Oph	16 37 09.4	-10 34 02	+0.001	+0.02	2.56	0.02	-3.6	O9.5 V	-19	170 s	
α TrA	16 48 39.8	-69 01 39	+0.005	-0.03	1.92	1.44	0.8	K2 III	-4	17 s	(Atria)
26 ϵ Sco	16 50 09.7	-34 17 36	-0.049	-0.25	2.29	1.15	0.8	K2 III	-3	20 mx	
35 η Oph	17 10 22.5	-15 43 30	+0.003	+0.09	2.43	0.06	1.2	A2 V	-1	18 ts	Sabik, m
35 λ Sco	17 33 36.4	-37 06 14	0.000	-0.03	1.63	-0.22	-3.0	B2 IV	0	84 s	Shaula, m
55 α Oph	17 34 55.9	+12 33 36	+0.008	-0.23	2.08	0.15	0.7	A5 III	+13	19 ts	Ras-Alhague
θ Sco	17 37 19.0	-42 59 52	+0.001	0.00	1.87	0.40	-5.4	F0 I-II	+1	280 s	
κ Sco	17 42 29.0	-39 01 48	-0.001	-0.03	2.41	-0.22	-3.0	B2 IV	-10	120 s	
33 γ Dra	17 56 36.2	+51 29 20	-0.001	-0.02	2.23	1.52	-0.2	K5 III	-28	31 s	Eltanin, m
20 ϵ Sgr	18 24 10.2	-34 23 05	-0.003	-0.13	1.85	-0.03	-0.2	B9 IV	-11	26 s	Kaus Australis, m
3 α Lyr	18 36 56.2	+38 47 01	+0.017	+0.28	0.03	0.00	0.5	A0 V	-14	8.1 t	Vega, m
34 σ Sgr	18 55 15.7	-26 17 48	+0.001	-0.05	2.02	-0.22	-2.0	B3 IV-V	-11	64 s	Nunki, m
53 α Aql	19 50 46.8	+8 52 06	+0.036	+0.39	0.77	0.22	2.2	A7 IV-V	-26	5.1 t	Altair, m
37 γ Cyg	20 22 13.5	+40 15 24	0.000	0.00	2.20	0.68	-4.6	F8 Ib	-8	230 s	Sadir, m
α Pav	20 25 38.7	-56 44 06	+0.002	-0.08	1.94	-0.20	-2.3	B3 IV	+2	71 s	(Peacock), m
50 α Cyg	20 41 25.8	+45 16 49	0.000	+0.01	1.25	0.09	-7.5	A2 Ia	-5	560 s	Deneb, m
53 ϵ Cyg	20 46 12.5	+33 58 13	+0.028	+0.33	2.46	1.03	0.5	K0 III	-10	25 ts	Gienar, m
5 α Cep	21 18 34.6	+62 35 08	+0.022	+0.05	2.44	0.22	1.7	A7 IV-V	-10	14 ts	Alderamin, m
8 ϵ Peg	21 44 11.0	+9 52 30	+0.002	0.00	2.38	1.52	-3.6	K2 Ib	+5	160 s	Enif, m

α Gru	22 08	13.8	-46 57	40	+0.013	-0.15	1.74	-0.13	0.1	B5 V	+12	21 mx	(Al Na'ir), m
β Gru	22 42	39.9	-46 53	05	+0.014	-0.01	2.11	1.62	-1.5	M3 II	+2	53 mx	
24 α PsA	22 57	38.9	-29 37	20	+0.026	-0.16	1.16	0.09	2.0	A3 V	+7	6.7 t	Fomalhaut
53 β Peg	23 03	46.3	+28 04	58	+0.014	+0.14	2.42	1.67	-1.2	M2 II-III	+9	54 s	Scheat, m, v
54 α Peg	23 04	45.5	+15 12	19	+0.004	-0.04	2.49	-0.04	0.0	B9 V	-4	31 ts	Markab

Column headings

Star name: the Flamsteed number, Bayer letter, and constellation abbreviation.

α 2000: right ascension for equator, equinox, and epoch 2000.0.

δ 2000: declination for equator, equinox, and epoch 2000.0.

$\mu(\alpha)$: proper motion in right ascension in seconds of time per year.

$\mu(\delta)$: proper motion in declination in seconds of arc per year.

V : visual magnitude in the standard U, B, V photometric system.

$B - V$: the color index in the U, B, V system.

M_v : the absolute visual magnitude ($M_v = V + 5 - 5 \log d$ (pc)).

Spec: the spectral type and luminosity class in the MKK system.

RV : radial velocity. Positive means receding from the Solar System.

d (pc): distance to the star in parsecs. 't' denotes the distance is based on a trigonometric parallax; 's' denotes spectroscopic parallax; 'mn' denotes a minimum likely value; 'mx' denotes a maximum likely value.

Notes: the classical names are given here; 'm' means the star is part of a multiple system; 'v' means the star is variable. Only the dominant star is listed. If more than one member of a multiple system is visually bright, all the visually bright members are listed ($V < 8.0$). The combined magnitude of a pair of close stars is given by:

$$m = m_2 - 2.5 \log(R + 1) \quad \text{where } R = 10^{0.4(m_2 - m_1)}, m_1 \text{ is the magnitude of the brightest component, and } m_2 \text{ is that of the fainter star}$$

(Data are from *Sky Catalogue 2000.00*, Hirshfeld, A. & Sinnott, R., eds., Sky Publishing Corp., 1982.)

Stars within 5 pc

No. Name	z 1950	δ 1950	Trig. parallax, π	Proper motion, μ , yr $^{-1}$	Radial velocity, v_r , km s $^{-1}$	Spec	l	b	l'	b'	$R-l$	M	Luminosity (L_{\odot})
1 Sun													
2 Proxima Cen	14 ^h 26 ^m 53	-62 28'	0.772 \pm 0.007	5.85	-16	G2 V	-26.72	0.65	0.10		1.65	4.85	1.0
z Cen A	14 36.2	-60 38'	0.750 \pm 0.010	3.68	22	G2 V	11.05	1.97			0.22	15.49	0.000106
z Cen B						K0 V	0.01	0.68			0.24	4.37	1.6
3 Barnard's star	17 55.4	+4 33'	0.545 \pm 0.003	10.31	-108	M5 V	1.33	0.88	1.29	1.74	1.25	5.71	0.45
4 Wolf 359	10 54.1	+7 19'	0.421 \pm 0.006	4.70	+13	dM8 e	9.54	2.01	1.54	1.85	1.85	13.22	0.00045
5 BD + 36 2147	11 00.6	+36 18'	0.397 \pm 0.004	4.78	-84	M2 V	15.53	1.51	1.12	0.91	0.91	16.65	0.00002
6 L 726 - 8 = A	1 36.4	-18 13'	0.387 \pm 0.012	3.36	129	dM6 e	7.80	1.85	1.09	1.6	1.6	10.50	0.0035
V V Cen = B					+32	dM6 e	12.52	0.00				15.46	0.000106
7 Sirius A	6 42.9	-16 39'	0.377 \pm 0.006	1.33	-8	A1 V	13.07	0.00	-0.04	0.12	0.12	15.96	0.000104
Sirius B						D8	1.46	-0.12				1.42	23.5
8 Ross 154	18 46.7	-23 53'	0.345 \pm 0.012	0.72	-4	GMS e	8.3	1.70	1.17	1.30	1.30	13.14	0.00048
9 Ross 248	23 39.4	-43 55'	0.314 \pm 0.004	1.60	-81	dM6 e	12.29	1.91	1.48	1.56	1.56	14.78	0.00011
10 z Eri	3 30.6	9 38'	0.305 \pm 0.004	0.98	-16	K2 V	3.53	0.88	0.58	0.30	0.30	6.14	0.70
11 Ross 128	11 45.1	+1 06'	0.298 \pm 0.006	1.38	13	GMS	11.10	1.76	1.30	1.30	1.30	13.47	0.00036
12 61 Cyg A	21 04.7	+38 30'	0.294 \pm 0.006	5.22	64	K5 V	5.22	1.17	1.11	0.47	0.47	7.56	0.082
61 Cyg B						K7 V	6.03	1.37	1.23	0.60	0.60	8.37	0.039
13 z Ind	21 59.6	-57 00'	0.291 \pm 0.010	4.70	40	K5 V	4.68	1.05	1.00	0.40	0.40	7.00	0.14
14 BD + 43 44A	0 15.5	+43 44'	0.290 \pm 0.006	2.90	+13	M1 V	8.08	1.56	1.24	0.88	0.88	10.39	0.0061
+ 43 44B					-20	M6 V e	11.06	1.80	1.40	1.22	1.22	13.37	0.00039
15 L 789 - 6	22 35.7	-15 36'	0.290 \pm 0.007	3.26	-60	dM7 e	12.18	1.96	1.54	1.66	1.66	14.49	0.00014
16 Procyon A	7 36.7	+5 21'	0.285 \pm 0.006	1.25	-3	F5 IV V	0.37	0.42	0.03	0.14	0.14	2.64	7.65
Procyon B						D8	10.7					13.0	0.00055
17 BD + 59 1915A	18 42.2	+59 33'	0.282 \pm 0.004	2.29	0	dM4	8.90	1.54	1.11	1.07	1.07	11.15	0.0030
+ 59 1915B				2.27	+10	dM5	9.69	1.59	1.14	1.14	1.14	11.94	0.0015
18 CD - 36 15693	23 02.6	-36 09'	0.279 \pm 0.024	6.90	-10	M2 V	7.55	1.48	1.18	0.85	0.85	9.58	0.013
19 G 51 - 15	8 26.9	+26 57'	0.278 \pm 0.004	1.27			14.81	2.06	1.79	1.79	1.79	17.03	0.00001
20 z Cen	1 41.7	-16 12'	0.277 \pm 0.007	1.92	-16	G8 V	3.80	0.72	0.22	0.26	0.26	5.72	0.45
21 BD + 5 1668	7 24.7	+5 23'	0.266 \pm 0.006	3.77	-26	dM5	9.82	1.56	1.12	1.19	1.19	11.94	0.0015
22 L 725 - 32	1 09.9	-17 16'	0.261 \pm 0.012	1.32	+28	dM5 e	12.04	1.83	1.46	1.44	1.44	14.12	0.00020


23	CD - 39 14192	21 14.3	-39 04	0.260 ± 0.012	3.46	-21	M0 V	6.66	1.40	1.20	0.69	8.74	0.028
24	Kapleyn's star	5 09.7	-45 00	0.256 ± 0.010	8.72	+245	sdM0 pec	8.84	1.56	1.05	0.77	10.88	0.0039
25	Kruger 60A	22 26.2	+52 27	0.253 ± 0.004	0.86	26	dM3	9.85	1.62	1.25	1.14	11.87	0.0016
	Kruger 60B						dM5 e	11.3	1.8	1.5		13.3	0.0004
26	BD - 12 4523	16 27.5	-12 32	0.247 ± 0.007	1.18	-13	dM5	10.11	1.60	1.16	1.20	12.07	0.0013
27	Ross 614 A	6 26.8	-2 46	0.246 ± 0.004	1.00	+34	dM7 e	11.10	1.71	1.15	1.40	13.12	0.00049
	Ross 614 B							14 }				16	0.00014
28	Van Maanen's star	0 46.5	+5 09	0.232 ± 0.004	2.99	+54	DG	12.37	0.56	0.02	0.16	14.20	0.00018
29	Wolf 424 A	12 30.9	+9 18	0.230 ± 0.006	1.76	-5	dM6 e	13.16 }	1.80	1.18	1.62	14.97	0.00009
	Wolf 424 B						dM6 e	13.4 }				15.2	0.000107
30	CD - 37 15492	0 02.5	-17 36	0.225 ± 0.012	6.11	+23	M4 V	8.56	1.46	1.03	0.92	10.32	0.00065
31	L 1159 - 16	1 57.5	+12 50	0.224 ± 0.004	2.09		dM8 e	12.26	1.82	1.35	1.35	14.01	0.00022
32	BD + 50 1725	10 08.3	+49 42	0.222 ± 0.010	1.45	26	K7 V	6.59	1.36	1.28	0.60	8.32	0.041
33	CD - 46 11540	17 24.9	-46 51	0.216 ± 0.012	1.06		dM4	9.37	1.53	1.21	1.03	11.04	0.0033
34	G 158 - 27	0 04.2	-7 48	0.214 ± 0.007	2.04		dM	13.74	1.95	1.21	1.52	15.39	0.00006
35	CD - 49 13515	21 30.2	-49 13	0.214 ± 0.010	0.81	+8	M1 V	8.67	1.46	1.05	0.93	10.32	0.00065
36	CD - 44 11909	17 33.5	-44 17	0.213 ± 0.007	1.16		M5	10.96	1.65	1.20	1.26	12.60	0.00079
37	BD + 68 946	17 36.7	+68 23	0.212 ± 0.006	1.31	-22	M3.5 V	9.15	1.50	1.08	1.10	10.79	0.0042
38	G 208 - 44 - A	19 52.3	+44 18	0.211 ± 0.004	0.74			13.41	1.91			15.03	0.00008
	G 208 - 44 - B							13.99	1.98			15.61	0.000105
39	BD - 15 6290	22 50.6	-14 31	0.209 ± 0.007	1.14	+9	dM5	10.17	1.60	1.15	1.22	11.77	0.0017
40	o(40)En A	4 13.0	-7 44	0.207 ± 0.003	4.08	-42	K1 V	4.43	0.82	0.44	0.31	6.01	0.34
	40 En B	4 13.1	-7 44		4.07	-21	DA	9.52	0.03	-0.68	0.30	11.10	0.0032
	40 En C					-45	dM4 e	11.17	1.66	0.83	1.21	12.75	0.00069
41	BD + 20 2465	10 16.9	+20 07	0.206 ± 0.006	0.49	+11	M4.5 V e	9.43	1.54	1.06	1.12	11.00	0.0035
42	L 145 - 141	1 43.0	64 33	0.206 ± 0.012	2.68		DC	11.50	0.19	-0.60	0.04	13.07	0.00052
43	70 Oph A	18 02.9	-2 31	0.203 ± 0.006	1.12	-7	K0 V	4.22 }	0.86	0.51	0.30	5.76	0.43
	70 Oph B						K5 V	6.00 }				7.54	0.084
44	BD + 43 4305	22 44.7	+44 05	0.200 ± 0.004	0.83	-2	dM5 e	10.2	1.6	1.1	1.15	11.7	0.0018
45	Altair	19 48.3	+8 44	0.198 ± 0.006	0.66	-26	A7 IV V	0.76	0.22	0.08	0.02	2.24	11.1
46	AC + 79 3888	1 44.6	+78 58	0.193 ± 0.007	0.89	-119	sdM4	10.8	1.60	1.18	1.18	12.25	0.0011
47	G 9 - 38 = A	8 55.4	+19 57	0.192 ± 0.004	0.89		m	14.06	1.84			15.48	0.00016
	I.P. 426 - 40 = B						m	14.92	1.93			16.34	0.000125
48	BD - 15 2620	17 43.2	+15 10	0.192 ± 0.007	1.30	+15	M4 V	8.49	1.44	1.10	0.86	9.91	0.0095

: denotes approximate value.

(Adapted from Landolt-Börnstein, VI/2C, Springer-Verlag, 1982.)

Nearest stars

Known stars within 5 parsecs of the Sun (in a sphere, projected on a plane). The numbers correspond to the order of radial distance from the Sun. The angular position on the plot is the star's right ascension. Stars that appear close together on the plot are not necessarily close in space, since the third dimension - for example, the star's declination - cannot be indicated. (Adapted from Roach, F. E. & Gordon, J. L., *The Light of the Night Sky*, D. Reidel Publishing Co., 1973.)



Stars of large proper motion

Name	Constellation	V mag.	Annual motion	α 2000	δ 2000
Groombridge 1830	Ursa Major	6.45	7.05	11 ^h 52 ^m 58.7	+ 37 43 07"
Lacaille 9352	Piscis Aus.	7.34	6.90	23 05 52.0	- 35 51 12
61 Cygni	Cygnus	4.84	5.22	21 06 53.9	+ 38 44 58
Lalande 21185	Ursa Major	7.49	4.77	11 03 20.2	+ 35 58 13
ϵ Indi	Indus	4.69	4.70	22 03 21.5	- 56 47 10
ρ^2 Eridani	Eridanus	4.43	4.08	04 15 16.2	- 07 39 10
μ Cassiopeiae	Cassiopeia	5.17	3.75	01 08 16.2	+ 54 55 15
α^2 Centauri	Centaurus	1.39	3.68	14 39 35.4	- 60 50 13
α^1 Centauri	Centaurus	0.00	3.68	14 39 36.7	- 60 50 02
Lacaille 8760	Microscopium	6.68	3.46	21 17 15.1	- 38 52 05
82 G Eridani	Eridanus	4.27	3.14	03 19 55.7	- 43 04 10
268 G Ceti	Cetus	5.82	2.32	02 36 04.8	+ 06 53 13
Arcturus	Boötes	-0.04	2.28	14 15 39.6	+ 19 10 57

(List from *Sky Catalogue 2000.0*, Hirshfeld, A. & Sinnott, R., eds., Sky Publishing Corp., 1982.)

The 21 fastest pulsars (as of August 1986)

Pulsar PSR	α 1950	δ 1950	P (s)	\dot{P} (10^{-9} s d $^{-1}$)	JD - 2440000	DM (pc cm $^{-3}$)
1937 + 214 ^(c)	19 ^h 37 ^m 28 ^s .72	21° 28' 17.3	0.001 557 806 449 023	1.07×10^{-5}	5303	71.20
1855 + 09 ^(c)	18 55 13.68	09 39 13.3	0.005 362 100 452 39	1.8×10^{-6}	6433	13.2943
1953 + 29 ^(c)	19 53 26.7	29 00 42.0	0.006 133 17	≤ 0.050	5427	104.5
0531 + 21 ^(a)	5 31 31.43	21 58 0.7	0.033 200 385 8	36.46	1994	56.791
1913 + 16 ^(c)	19 13 12.48	16 01 8.4	0.059 029 995 272	0.000 76	2322	167.000
0833 - 45 ^(b)	8 33 39.35	-45 00 8.6	0.089 234 713 88	10.8026	2701	69.080
1356 - 60	13 56 26.18	-60 23 36.2	0.127 500 776 85	0.547 65	3556	295.000
1930 + 22	19 30 12.49	22 15 19.0	0.144 427 889 857 53	4.992 17	2676	219.00
0355 + 54	3 55 0.45	54 04 42.6	0.156 380 055 92	0.379	1536	57.030
1804 - 08	18 04 53.91	-8 48 10.5	0.163 727 360 007	0.002 48	3557	112.800
0740 - 28	7 40 47.87	-28 15 32.6	0.166 753 902 088	1.454 79	3556	73.770
1541 - 52	15 41 12.59	-52 59 22.9	0.178 553 799 688	0.005 24	3556	35.200
1449 - 64	14 49 25.30	-64 01 0.5	0.179 483 983 915	0.237 42	3556	70.500
1821 - 19	18 21 2.78	-19 47 29.0	0.189 332 134 77	0.452 55	3557	225.700
1557 - 50	15 57 8.83	-50 35 55.7	0.192 598 318 85	0.4375	2554	270.00
1556 - 57	15 56 14.66	-57 42 47.4	0.194 453 886 93	0.183 58	3557	176.900
1915 + 13	19 15 21.57	13 48 28.7	0.194 626 341 491	0.622 33	2302	94.00

0656 + 64 ^(c)	6 56 30.00	64 20 0.0	0.1955	3513	5
1055 - 52	10 55 48.59	-52 10 51.9	0.197 107 608 187	3556	30.100
0743 - 53	7 43 50.50	-53 44 1.0	0.214 836 351 4	3780	122.300
1221 - 63	12 21 34.71	-63 51 16.3	0.216 474 842 97	3556	96.900

α, δ = position,

P = pulse period,

\dot{P} = the rate of change of the period,

JD = Julian date of observation,

DM = dispersion measure = $\int_{f_0}^{f_d} N_e df$.

where d = distance of the pulsar from the Sun.

N_e = electron number density in interstellar space.

The pulse arrival time for two different observing frequencies f_1 and f_2 differs by:

$$t_2 - t_1 = \frac{e^2}{2\pi m_e c} \left(\frac{1}{f_2^2} - \frac{1}{f_1^2} \right) DM.$$

(a) Crab pulsar.


(b) Vela pulsar.

(c) Binary pulsar.


(Data from *Landolt-Börnstein*, Band 2c, Springer-Verlag, 1982, except PSR 1937 + 214: Baker, D. C. & Kulkarni, S. R., *Nature*, **301**, 314, 1983, PSR 1953 + 29: Boriakoff, V. *et al.*, *Nature*, **304**, 417, 1983 and PSR 1855 + 09: Segelstein, D. J. *et al.*, *Nature*, **322**, 714, 1986.)

Pulsars

Galactic distribution of pulsars. In the adopted coordinates, 0° latitude corresponds to the Galactic plane, while 0° longitude, 0° latitude corresponds to the direction of the Galactic center. (Courtesy of Y. Terzian, Cornell University.)




Distribution of pulsars in Galactic latitude. Latitude 0° corresponds to the Galactic plane. (Courtesy of Y. Terzian, Cornell University.)



Pulsars (cont.)

Distribution of periods and period derivatives for 353 pulsars. The seven known binary pulsars, indicated by circles around the dots, have unusually small period derivatives and hence relatively weak magnetic fields. (Dewey, R. J. *et al.*, *Nature*, **322**, 712, 1986, with permission.)



Parameters of binary radio pulsars

PSR	P (ms)	$\log \dot{P}$	$\log B^{(a)}$ (Gauss)	$ z ^{(b)}$ (pc)	$a_1 \sin i^{(c)}$ (R_{\odot})	P_b (d)	e	$f(m_1, m_2)^{(d)}$ (M_{\odot})	Likely $m_2^{(e)}$ (M_{\odot})
1855+09	5.4	-19.7	8.5	20	4.0	12.33	0.00002	0.0052	0.2-0.4
1953+29	6.1	-19.5	8.6	20	13.6	117.35	0.0003	0.0027	0.2-0.4
0655+64	195.6	-18.2	10.0	120	1.8	1.03	<0.00005	0.0712	0.7-1.3
1913+16	59.0	-17.1	10.3	190	1.0	0.32	0.6171	0.1322	1.4
1831-00	520.9	<-17.0	<10.9	190	0.3	1.81	<0.005	0.00012	0.06-0.13
0820+02	864.9	-16.0	11.5	280	70.0	1232.40	0.0119	0.0030	0.2-0.4
2303+46	1066.4	-15.4	11.8	480	14.1	12.34	0.6584	0.2463	1.2-2.5

^(a) $B = 3.2 \times 10^{19} (P\dot{P})^{1/2}$.

^(b) The distance above the galactic plane.

^(c) R_{\odot} represents the Sun's radius as a unit of measurement.

^(d) $f(m_1, m_2) = \frac{(m_2 \sin i)^3}{(m_1 + m_2)^2} = \frac{G}{P_b^2} \frac{P_b^2}{G}$, where m_1 and m_2 are the masses of the pulsar and companion, respectively, a_1 is the orbital

semi-major axis; i is the angle between the plane of the orbit and the plane of the sky.

^(e) M_{\odot} represents the Sun's mass as a unit of measurement.

(List taken from Dewey, R. J. *et al.*, *Nature*, **322**, 712, 1986.)

Prominent OB associations

Name	α 2000	δ 2000	Diameter ($'$)	Distance (pc)	O stars	B stars	RV (km s $^{-1}$)	Clusters	Stars
Cas OB4	0 ^h 28 ^m 4	+62 42'		2880	5	12	-43	103	
Cas OB14	0 28.8	+63 22		1110	0	3	-8		κ Cas
Cas OB1	1 00.8	+61 30	120	2510	0	5	-38	381?	
Cas OB8	1 46.2	+61 19		2880	1	10	-30	581, 663; 654?	
Per OB1	2 14.5	+57 19	360	2290	9	56	-41	h, γ Per	
Cas OB6	2 43.2	+61 23	480	2190	17	8	-47	IC 1805	
Cam OB1	3 31.6	+58 38		1000	3	9	-6	1444? 1502?	
Per OB3	3 27.8	+49 54		170					α, δ Per
Per OB2	3 42.2	+33 26	480 \times 300	400	1	3			ξ, η, ζ Per
Aur OB2	5 28.3	+34 54		3160	5	3	-13	1893, IC 410	
Aur OB1	5 21.7	+33 52	360 \times 300	1320	5	5	-3	1912, 60; 1931?	
Gem OB1	6 09.8	+21 35	300	1510	4	13	+13	2175?	ζ^2 Ori
Ori OB1	5 31.4	-2 41	960	460	9	6		Trapezium	$\theta, \beta, \gamma, \delta, \epsilon$ Ori
Mon OB1	6 33.1	+8 50	840 \times 300	550	1	0	+22	2264	S Mon
Mon OB2	6 37.2	+4 50	360 \times 250	1510	10	7	+28	2244	Plaskett's star
CMa OB1	7 07.0	-10 28	240	1320	4	3	+27	2335, 53; 2343?	
Pup OB1	7 54.8	-27 05	240 \times 180	2510	7	0	+43	2467?	
Vel OB2	8 11.8	-47 50		460					Vela pulsar?
Vel OB1	8 49.9	-45 00	360 \times 240	1400	5	11		2659?	
Car OB1	10 46.7	-59 05	120 \times 66	2510	6	15		3293; IC 2581?	
Car OB2	11 06.0	-59 51	330 \times 150	2000	8	6		3572, Tr 18	
Car OB4	11 08.3	-60 31	114 \times 54					3590	
Cen OB2	11 35.3	-62 36	84 \times 48	2500				IC 2944	ζ Cen
Cen OB1	13 04.8	-62 04	360	2510	2	19		4755	ζ Cru
Nor OB1	15 58.7	-54 30		2500	0	6		6031?	

Name	α 2000	δ 2000	Diameter ($''$)	Distance (pc)	O stars	B stars	RV (km s^{-1})	Clusters	Stars
Sco-Cen	16^{h} $^{\text{m}}$	-25:		160				IC 2602?	α CMa, α Car, α Eri
Ara OB1	16 39.5	-46 46	270×180	1380				6169, 93	μ Nor
Sco OB1	16 53.5	-41 57	96×66	1910	18	10	-18	6231	γ Sco
Sco OB2	16 14.9	-25 55		160	0	3			α, β^1, δ Sco
Sgr OB1	18 07.9	-21 28	570×240	1580	8	9	-4	6514, 30 1	μ Sgr
Sgr OB4	18 14.4	-19 03		2400	1	6	+3	6603	
Ser OB1	18 20.8	-14 35	300×180	2190	9	9	-23	6611	
Ser OB2	18 18.6	-11 58	500:	2000:	9	6	+6	6604?	
Vul OB1	19 44.0	+24 13		2000	5	7	+8	6823	
Cyg OB3	20 04.7	+35 50		2290	9	15	0	6871?	Cyg X-1
Cyg OB1	20 17.8	+37 38	420×240	1820	12	28	-7	6913, IC 4996	
Cyg OB9	20 23.3	+39 56		1200	7	7	-20	6910	
Cyg OB2	20 32.4	+41 17	30	1820	13	2			
Cyg OB4	21 13.1	+37 52		1000					
Cyg OB7	21 02.7	+49 43		830	3	6	-10		σ Cyg
Cep OB2	21 47.9	+61 04	460	830	8	9	-20	7160, IC 1396	α Cyg
Cep OB1	22 24.6	+55 14	210	3470	7	26	-51	7380?	μ, ν, λ Cep
Lac OB1	22 41.2	+39 05	900×540	600	1	0			β Cep
Cep OB3	23 00.4	+64 03		870	3	3	-21		10 Lac
Cas OB5	23 58.7	+60 22	150	2510	5	10	-46	7788; 7790?	ρ Cas
Cep OB4	23 59.5	+67 35		840	2	0			

: denotes approximate value.

(Adapted from Sky Catalogue 2000.0, Vol. 2, Sky Publishing Corp., 1985.)

The orbital elements of some binary stars

Name	α 2000	δ 2000	Period, P (mean solar years)	Epoch of periastron, t	Longitude of periastron, ω ($^\circ$)	Eccen- tricity, e	Semi- major axis of orbit, a (arcsec)	Inclin. orbit, i ($^\circ$)	Position angle of ascending node, Ω ($^\circ$)	Distance, d (pc)
η CrB	15 ^h 23 ^m 21 ^s	30° 17'	41.623	1934.008	219.907	0.2763	0.907	59.025	23.717	14
γ Vir	12 41 40	-01 27	171.37	1836.433	252.88	0.8808	3.746	146.05	31.78	11
η Cas	00 49 06	57 49	480	1889.6	268.59	0.497	11.9939	34.76	278.42	5.9
ζ Ori	05 40 46	-01 56	1508.6	2070.6	47.3	0.07	2.728	72.0	155.5	340
α CMa (Sirius)	06 45 09	-16 43	50.09	1894.13	147.27	0.5923	7.500	136.53	44.57	2.7
δ Gem	07 20 07	21 59	1200	1437	57.19	0.1100	6.9753	63.28	18.38	18
α Gem (Castor)	07 34 36	31 53	420.07	1965.3	261.43	0.33	6.295	115.94	40.47	14
α CMi (Procyon)	07 39 18	5 14	40.65	1927.6	269.8	0.40	4.548	35.7	284.3	3.5
α Cen	14 39 36	-60 50	79.920	1955.56	231.560	0.516	17.583	79.240	204.868	1.3
α Sco (Antares)	16 29 24	-26 26	900	1889.0	0.0	0.0	3.21	86.3	273.0	100

a semi-major axis,
 P period of revolution,
 M_1, M_2 stellar masses,

$$\frac{a^3}{P^2} = \frac{G}{4\pi^2} (M_1 + M_2), \text{ where } G \text{ is the gravitational constant (Kepler's third law).}$$

If we express masses in solar masses, periods in years, and distances in astronomical units, we have

$$(M_1 + M_2)P^2 = a^3 \quad (a \text{ (AU)} = a \text{ (arcsec)} \times d \text{ (pc)}).$$

e eccentricity,
 t epoch of the periastron passage (the closest approach of the stars),
 Ω position of the ascending node. The nodes are points of intersection of the relative orbit and a plane tangential to the celestial sphere at the position of the bright component,
 ω longitude of the periastron, the angle between the radius vector to the ascending node and that in the direction of the periastron, measured from the node to the periastron in the direction of the orbital motion,
 i inclination, the angle between the orbital plane and the plane tangential to the celestial sphere.
 (Adapted from Duffett-Smith, P., *Practical Astronomy With Your Calculator*, Cambridge University Press.)

Main class	Subclass	Brightness variation (mag)	Period (d)	Typical representative	Brightness ^(a) (mag)		Period (d)
					max	min	
Cepheids	C	0.1-2.0	1-50 or 70	TW CMa	9.5	11.0 p	6.99
	C δ	0.1-2.0	1-50 or 70	δ Cep	4.1	5.2 p	5.37
	CW	0.1-2.0	1-50 or 70	W Vir	9.9	11.3 p	17.29
	I	RX Cep	7.5	7.8 v	
	Ia	V395 Cyg	7.8	8.4 v	
Long-period variables	Ib	CO Cyg	9.6	10.6 v	
	Ic	TZ Cas	9.2	10.5 v	
	M	2.5-5.0	80-1000	α Cet	2.0	10.1 v	331.62
		and more					
Red giant variables	SR	1-2.0	30-1000	VW UMa	8.4	9.1 p	125
	SRa	< 2.5		Z Aqr	9.5	12.0 p	136.9
	SRb			AF Cyg	7.4	9.4 p	94.1
	SRc			μ Cep	3.6	5.1 v	
RR Lyrae variables	SRd			UU Her	8.5	10.6 p	
	RR	< 1-2.0	0.05-1.2	V756 Oph	12.3	13.7 p	
	RRa	< 1.5	0.5 and 0.7	RR Lyr	6.94	8.03 p	0.567
	RRc	3	0.3	SX UMa	10.6	11.2 p	0.307
	RV	3	30-150	EP Lyr	10.2	11.6 p	83.43
RV Tauri variables	RVa	3	30-150	AC Her	7.4	9.2 p	75.46
	RVb	3	30-150	R Sge	9.0	11.2 p	70.594
	β C	0.1	0.1-0.3	β Cep	3.3	3.35 p	0.190
				β Canis Major V.			
	δ Sc	< 0.25	1.0	δ Sct	4.9	5.19 p	0.194
χ^2 CV	< 0.1	1-25	χ^2 CVn	3.0	3.1 p	5.47	

Galactic supernova remnants

Name	Galactic coordinates $l^{\text{II}}, b^{\text{II}}$	α 1950	δ 1950	Radio size	Optical size
CTA 1	119.53, +9.77	00 ^h 04 ^m 18 ^s	+72 04.5	90'	50' × 90'
Tycho	120.09, +1.41	00 22 33	+63 51.8	8'	8'
HB 3	132.70, +1.30	02 14	+62 18	140'	...
HB 9	160.39, +2.75	04 57	+46 36	130' × 155'	90' × 125'
OA 184	166.07, +4.40	05 15 38	+41 46	70' × 90'	70' × 80'
VRO 42.05.01	166.27, +2.53	05 23 21	+43 00	70' × 75'	35' × 40'
S 147	180.33, -1.68	05 36 45	+27 44.5	175'	195' × 200'
Crab	184.55, -5.78	05 31 31	+21 58.9	290" × 420"	290" × 420"
IC 443	189.01, +3.02	06 14 06	+22 37.2	47' × 54'	48'
Monoceros	205.62, -0.10	06 35	+06 30	210'	180' × 200'
Puppis	260.40, -3.42	08 20 30	-42 50	45' × 65'	50' × 80'
Vela	263.37, -3.01	08 32	-45 00	300'	270'
MSH 10 - 53	284.17, -1.78	10 15 40	-58 40.5	33' × 50'	1' × 5'
RCW 86	315.44, -2.33	14 39 08	-62 15	55'	8' × 31'
RCW 89	320.36, -0.97	15 09 30	-58 46	8'	450" × 580"
RCW 103	332.43, -0.39	16 13 54	-50 55.8	7'	5.7 × 9.5
Kes 45	342.05, +0.13	16 50 11	-43 30.3	30'	... × 20'
Kepler	004.52, +6.82	17 27 41	-21 26.6	3'	21" × 64"
W28	006.46, -0.09	17 57 36	-23 25	30'	30'
3C 400.2	053.62, -2.23	19 36 30	+17 08	20'	4' × 6'
DR 4	078.13, +1.81	20 20 38	+40 03.4	≤ 3'	2' × 3'
Cygnus	074.27, -8.49	20 49 30	+30 45	160' × 240'	160' × 210'
Cas A	111.73, -2.13	23 21 10	+58 32.4	130'	130'
CTB 1	116.94, +0.18	23 56 45	+62 10	35' × 45'	32'

: denotes approximate value.

(Adapted from van den Bergh *et al.*, *Ap. J. Suppl.*, **26**, 19, 1973.)


Henry Draper (HD) spectral classification

Class	Class characteristics
O	Hot stars with He II absorption
B	He I absorption; H developing later
A	Very strong H, decreasing later; Ca II increasing
F	Ca II stronger; H weaker; metals developing
G	Ca II strong; Fe and other metals strong; H weaker
K	Strong metallic lines; CH and CN bands developing
M	Very red; TiO bands developing strongly

Spectral type and luminosity class (MK, or Yerke's classification)


Luminosity class	Examples:	Spectral type
Ia	Supergiants	α Boo (Arcturus) K2 III
Ib	Supergiants	α CMi (Procyon) F5 IV
II	Bright giants	β Gem (Pollux) K0 III
III	Giants	α Lyr (Vega) A0 V
IV	Subgiants	α UMi (Polaris) F8 Ib
V	Main sequence (dwarfs)	α CMa (Sirius) A1 V
VI	Subdwarfs	α Cyg (Deneb) A2 Ia
VII	White dwarfs	α Leo (Regulus) B7 V
		β Ori (Rigel) B8 Ia
		Sun G2 V

Spectral type and luminosity class of the MK classification:
dependence on color index $B - V$ and visual absolute magnitude M_v .
(Adapted from Unsöld, A., *The New Cosmos*, Springer-Verlag, 1969.)



Hertzsprung-Russell diagram

Hertzsprung-Russell or temperature luminosity diagram. (Adapted from Goldberg, L. & Dyer, E. R. in *Science in Space*, L. V. Berkner & H. Odishaw, eds., McGraw-Hill Book Company, 1961.)



An incomplete list of astrophysically important infrared and visible spectral features

Identification	Wavelength (Å) [†]	Identification	Wavelength (Å)
He I	10 830	O II	4649
C I	10 691	N III	4641
Si I	10 689	N III	4634
Si I	10 627	Mg I	4571
Si I	10 603	He II	4541
Si I	10 371	Mg II	4481
He II	10 124	He I	4472
He II	10 120	He I	4471
Na I	9961	He I	4388
C III	9710	Fe I	4384
O I	8446	[O III]	4363
He II	8237	H I γ	4340
O I	7774	O II	4318
He I	7065	G band ^(b)	4300
H I α	6563	Ca I	4227
[O I]	6363 ^(a)	He II	4200
[O I]	6300	H I δ	4102
Na I (D)	5896	He II	4100
Na I (D)	5890	N III	4097
He I	5876	Si IV	4089
He II	5412	O II	4073
Fe XIV	5303	He I	4026
Mg I	5175 ^(b)	H I ϵ	3970
Mg I	5173	Ca II (H)	3968
[O III]	5007	[Ne III]	3968
[O III]	4959	He I	3965
He I	4922	Ca II (K)	3934
H I β	4861	[Ne III]	3869
He II	4686	He I	3820
C IV	4658	O III	3760
O II	4650	[O II]	3727
C III	4650	[Ne V]	3426

[†] $h\nu$ (eV) = 123 98.54/ λ (Å).

^(a) Forbidden transitions are noted by brackets.

^(b) Superposition of CH band and metallic lines.

The main sequence^(a)

Sp (V)	U-V	B-V	V-R	V-I	V-J	V-K	V-L	V-M	V-N	BC	T _e (K)
O5-7	-1.46	-0.32	-0.15	-0.47	-0.73	-0.94	-1.01	-	-	-	38 000
O8-9	-1.44	-0.31	-0.15	-0.47	-0.73	-0.94	-1.01	-	-	-	35 000 ^(b)
O9.5	-1.40	-0.30	-0.14	-0.46	-0.73	-0.94	-1.00	-	-	-3.34	31 900
B0	-1.38	-0.30	-0.13	-0.42	-0.70	-0.93	-0.99	-	-	-3.17	30 000
B0.5	-1.29	-0.28	-0.12	-0.39	-0.66	-0.88	-0.93	-	-	-2.80	27 000 ^(b)
B1	-1.19	-0.26	-0.11	-0.36	-0.61	-0.81	-0.86	-	-	-2.50	24 200
B2	-1.10	-0.24	-0.10	-0.32	-0.55	-0.74	-0.77	-	-	-2.23	22 100
B3	-0.91	-0.20	-0.08	-0.27	-0.45	-0.61	-0.63	-	-	-1.77	18 800
B5	-0.72	-0.16	-0.06	-0.22	-0.35	-0.47	-0.48	-	-	-1.39	16 400
B6	-0.63	-0.14	-0.06	-0.19	-0.30	-0.41	-0.41	-	-	-1.21	15 400
B7	-0.54	-0.12	-0.04	-0.17	-0.25	-0.35	-0.34	-	-	-1.04	14 500
B8	-0.39	-0.09	-0.02	-0.12	-0.17	-0.24	-0.22	-	-	-0.85	13 400
B9	-0.25	-0.06	0.00	-0.06	-0.09	-0.14	-0.11	-	-	-0.66	12 400
A0	0.00	0.00	+0.02	0.00	-0.01	-0.03	0.00	-0.03	-0.03	-0.40	10 800
A2	+0.12	+0.06	+0.08	+0.09	+0.11	+0.13	+0.16	+0.13	+0.13	-0.25	9730
A5	+0.25	+0.14	+0.16	+0.22	+0.27	+0.36	+0.40	+0.36	+0.36	-0.15	8620
A7	+0.30	+0.19	+0.19	+0.28	+0.35	+0.46	+0.52	+0.46	+0.46	-0.12	8190
F0	+0.37	+0.31	+0.30	+0.47	+0.58	+0.79	+0.86	+0.79	+0.79	-0.08	7240
F2	+0.39	+0.36	+0.35	+0.55	+0.68	+0.93	+1.07	+0.93	+0.93	-0.06	6930
F5	+0.43	+0.43	+0.40	+0.64	+0.79	+1.07	+1.25	+1.07	+1.07	-0.04	6540

F8	+0.60	+0.54	+0.47	+0.76	+0.96	+1.27	+1.45	+1.27	+1.27	-0.05	6200
G0	+0.70	+0.59	+0.50	+0.81	+1.03	+1.35	+1.53	+1.35	+1.35	-0.06	5920
G2	+0.79	+0.63	+0.53	+0.86	+1.10	+1.44	+1.61	+1.44	+1.44	-0.07	5780
G5	+0.86	+0.66	+0.54	+0.89	+1.14	+1.49	+1.67	+1.49		-0.10	5610
G8	+1.06	+0.74	+0.58	+0.96	+1.24	+1.63	+1.85	+1.63		-0.15	5490
K0	+1.29	+0.82	+0.64	+1.06	+1.38	+1.83	+2.00	+1.83		-0.19	5240
K2	+1.60	+0.92	+0.74	+1.22	+1.57	+2.15	+2.24	+2.15		-0.25	4780
K5	+2.18	+1.15	+0.99	+1.62	+2.04	+2.75	+2.84	+2.75			
K7	+2.52	+1.30	+1.15	+1.93	+2.36	+3.21	+3.40	+3.21		-0.65	4410
M0	+2.67	+1.41	+1.28	+2.19	+2.71	+3.60	+3.78	+3.60		-0.90	4160
M1	+2.70	+1.48	+1.40	+2.45	+3.06	+3.95	+4.15	+3.95		-1.20	3920
M2	+2.69	+1.52	+1.50	+2.69	+3.37	+4.27	+4.47	+4.27		-1.48	3680
M3	+2.70	+1.55	+1.60	+2.94	+3.66	+4.57	+4.79	+4.57		-1.76	3500
M4	+2.70	+1.56	+1.70	+3.19	+3.97	+4.87	+5.20	+4.87		-2.03	3360
M5	+2.80	+1.61	+1.80	+3.47	+4.28	+5.17	(+5.54)	+5.17		-2.31	3230
M6	+2.99	+1.72	+1.93	+3.76	+4.63	+5.58	(+6.03)	+5.58		-2.62	3120
M7	+3.24	+1.84	+2.20	+4.20	+5.20	+6.18		+6.18			
M8	(+3.50)	(+2.00)	(+2.50)	(+4.70)	(+5.80)	(+6.75)		(+6.75)		4.20	2660

Bolometric corrections, BC , effective temperatures, T_e , and colors for stars of various spectral types, $m_b = BC + V$, where m_b and V are the bolometric and visual magnitudes, respectively.

^(a) Morgan & Keenan classification of spectral types.

^(b) Interpolated values.

(Adapted from Johnson, H., *Ann. Rev. Astron. and Astrophys.*, **4**, 193, 1966 and for BC and T_e , Lang, K., *Astrophysical Formulae*, Springer-Verlag, 1974.)

Giant stars

Sp (III)	$U - V$	$B - V$	$V - R$	$V - I$	$V - J$	$V - K$	$V - L$	$V - M$	$V - N$	BC	T_e (K)
G5	+1.55	+0.92	+0.69	+1.17	+1.52	+2.08	+2.18	+2.02	+2.05	-0.20	5010
G8	+1.64	+0.95	+0.70	+1.18	+1.56	+2.16	+2.27	+2.09	+2.12	-0.21	4870
K0	+1.93	+1.04	+0.77	+1.30	+1.71	+2.35	+2.47	+2.25	+2.28	-0.30	4720
K1	+2.13	+1.10	+0.81	+1.37	+1.80	+2.48	+2.61	+2.36	+2.39	-0.36	4580
K2	+2.32	+1.16	+0.84	+1.42	+1.87	+2.59	+2.73	+2.45	+2.48	-0.42	4460
K3	+2.74	+1.30	+0.96	+1.61	+2.12	+2.92	+3.07	+2.75	+2.80	-0.59	4210
K4	+3.07	+1.41	+1.06	+1.81	+2.36	+3.24	+3.39	+3.05	+3.11	-0.79	4010
K5	+3.34	+1.54	+1.20	+2.10	+2.71	+3.67	+3.83	+3.47	+3.54	-1.08	3780
M0	+3.43	+1.55	+1.23	+2.17	+2.82	+3.79	+3.96	+3.59	+3.65	-1.17	3660
M1	+3.48	+1.56	+1.28	+2.27	+2.90	+3.92	+4.09	+3.72	+3.78	-1.25	3600
M2	+3.51	+1.59	+1.34	+2.44	+3.08	+4.11	+4.29	+3.91	+3.97	-1.41	3500
M3	+3.51	+1.60	+1.48	+2.79	+3.51	+4.58	+4.77	+4.39	+4.45	-1.80	3300
M4	+3.32	+1.59	+1.74	+3.39	+4.26	+5.24	+5.44	+5.10	+5.14	-2.44	3100
M5	+3.00	+1.55	+2.18	+4.14	+5.04	+6.06	+6.31	+6.00	+6.00	-3.23	2950
M6	+2.43	+1.54	+2.80	+5.06	+5.86	+7.01	+7.39			-4.15	2800

Bolometric corrections, BC , effective temperatures, T_e , and colors for stars of various spectral types, $m_h = BC + V$, where m_h and V are the bolometric and visual magnitudes respectively.

Classification and absolute magnitude of stars (M_v)

Sp	Supergiants		Bright giants II	Giants III	Sub-giants IV	Main sequence dwarfs V		Population II				
	Ia	Ib				ZAMS ^(a) V	White dwarfs VII	Sub-dwarfs VI	Red branch	Horiz. branch		
O5	-6.4			-5.4		-5.7						
B0	-6.7	-6.1	-5.4	-5.0	-4.7	-4.1	-3.3	+10.2				
B5	-6.9	-5.7	-4.3	-2.4	-1.8	-1.1	-0.2	+10.7				+2.3
A0	-7.1	-5.3	-3.1	-0.2	+0.1	+0.7	+1.5	+11.3				+0.8
A5	-7.7	-4.9	-2.6	+0.5	+1.4	+2.0	+2.4	+12.2				+0.5
F0	-8.2	-4.7	-2.3	+1.2	+2.0	+2.6	+3.1	+12.9				+0.4
F5	-7.7	-4.7	-2.2	+1.4	+2.3	+3.4	+3.9	+13.6		+4.8	+4.8	+0.4
G0	-7.5	-4.7	-2.1	+1.1	+2.9	+4.4	+4.6	+14.3		+5.7	+4.1	+0.3
G5	-7.5	-4.7	-2.1	+0.7	+3.1	+5.1	+5.2	+14.9		+6.4	+2.0	-0.1
K0	-7.5	-4.6	-2.1	+0.5	+3.2	+5.9	+6.0	+15.3		+7.3	-0.2	-0.6
K5	-7.5	-4.6	-2.2	-0.2		+7.3	+7.3	+15		+8.4	+2.2	-2.2
M0	-7.5	-4.6	-2.3	-0.4		+9.0	+9.0	+15		+10	-3	
M2			-2.4	-0.6		+10.0	+10.0			+12		
M5				-0.8		+11.8	+11.8			+14		
M8						+16				+16		

Relation between absolute magnitude M_v and emission line width W (FWHM in km s^{-1}):

Emission line	Relation
Ca II K	$M_v = 27.59 - 14.94 \log W$ (Wilson-Bappu)
Mg II K	$M_v = 34.93 - 15.15 \log W$
H δ	$M_v = (40.2 \pm 4.5) - (14.7 \pm 1.6) \log W$

^(a) Zero age main sequence.

(After Allen, C. W., *Astrophysical Quantities*, The Athlone Press, 1973.)

Stellar mass, luminosity, radius and density (luminosity and radius with mass; white dwarfs omitted)

$\log(M/M_{\odot})$	$\log(L/L_{\odot})$	M_{bol}	M_V	M_B	$\log(R/R_{\odot})$ main seq.
-1.0	-2.9	+12.1	15.5	+17.1	-0.9
-0.8	-2.5	+10.9	13.9	+15.5	-0.7
-0.6	-2.0	+9.7	12.2	+13.9	-0.5
-0.4	-1.5	+8.4	10.2	+11.8	-0.3
-0.2	-0.8	+6.6	7.5	+8.7	-0.14
0.0	0.0	+4.7	4.8	+5.5	0.00
+0.2	+0.8	+2.7	2.7	+3.0	+0.10
+0.4	+1.6	+0.7	1.1	+1.1	+0.32
+0.6	+2.3	-1.1	-0.2	-0.1	+0.49
+0.8	+3.0	-2.9	-1.1	-1.2	+0.58
+1.0	+3.7	-4.6	-2.2	-2.4	+0.72
+1.2	+4.4	-6.3	-3.4	-3.6	+0.86
+1.4	+4.9	-7.6	-4.6	-4.9	+1.00
+1.6	+5.4	-8.9	-5.6	-6.0	+1.15
+1.8	+6.0	-10.2	-6.3	-6.9	+1.3

(After Allen, C. W., *Astrophysical Quantities*, Athlone Press, 1973.)

Stellar mass, luminosity, radius and density (mass, radius, luminosity, and mean density with spectral class)^(a)

Sp	log(M/M_{\odot})					log(R/R_{\odot})					log ρ (g cm^{-3})					log(L/L_{\odot})				
	I	III	V	I	III	I	III	V	I	III	I	III	V	I	III	I	III	V		
O5	+2.2		+1.6					+1.25					-2.0					+5.7		
B0	+1.7		+1.25	+1.3	+1.2			+0.87					-1.2			+5.4		+4.3		
B5	+1.4		+0.81	+1.5	+1.0			+0.58					-0.78			+4.8		+2.9		
A0	+1.2		+0.51	+1.6	+0.8			+0.40					-0.55			+4.3		+1.9		
A5	+1.1		+0.32	+1.7				+0.24					-0.26			+4.0		+1.3		
F0	+1.1		+0.23	+1.8				+0.13					-0.01			+3.9		+0.8		
F5	+1.0		+0.11	+1.9	+0.6			+0.08				+0.03				+3.8		+0.4		
G0	+1.0	+0.4	+0.04	+2.0	+0.8			+0.02				-1.8				+3.8	+1.5	+0.1		
G5	+1.1	+0.5	-0.03	+2.1	+1.0			-0.03				-2.4				+3.8	+1.7	-0.1		
K0	+1.1	+0.6	-0.11	+2.3	+1.2			-0.07				-2.9				+3.9	+1.9	-0.4		
K5	+1.2	+0.7	-0.16	+2.6	+1.4			-0.13				-3.4				+4.2	+2.3	-0.8		
M0	+1.2	+0.8	-0.33	+2.7				-0.20				-4				+4.5	+2.6	-1.2		
M2	+1.3		-0.41	+2.9				-0.3				-7.2				+4.7	+2.8	-1.5		
M5			-0.67					-0.5								+3.0		-2.1		
M8			-1.0					-0.9									+3.1		-3.1	

^(a) I = supergiant, III = giant, V = dwarf.

A single column between III and V represents main sequence.

(After Allen, C. W., *Astrophysical Quantities*, Athlone Press, 1973.)

Present-day mass function (PDMF)

M_v	$\phi(M_v)$ (stars pc^{-3} mag^{-1})	$\log M_v M_\odot$	$-\frac{dM_v}{d \log M}$	$2H$ (pc)	$\log T_{\text{MS}}$ (y)	f_{MS}	$\phi_{\text{MS}}(\log M)$ (stars pc^{-2} $\log M^{-1}$)
-6	1.49(-8)*	2.07	3.7	180	6.42	0.40	3.97(-6)*
-5	7.67(-8)	1.80	3.7	180	6.50	0.40	2.04(-5)
-4	3.82(-7)	1.53	3.7	180	6.58	0.41	1.04(-4)
-3	1.80(-6)	1.26	3.7	180	6.84	0.42	5.03(-4)
-2	7.86(-6)	0.99	3.7	180	7.19	0.43	2.25(-3)
-1	3.07(-5)	0.72	3.7	180	7.68	0.46	9.41(-3)
0	1.04(-4)	0.45	10.8	180	8.36	0.50	1.01(-1)
1	2.95(-4)	0.36	10.8	180	8.62	0.56	3.21(-1)
2	6.94(-4)	0.26	10.8	180	8.93	0.64	8.63(-1)
3	1.36(-3)	0.17	10.8	300	9.24	0.78	3.44(+0)
4	2.26(-3)	0.08	10.8	465	9.60	0.98	1.11(+1)
5	3.31(-3)	-0.02	10.8	630	9.83	1.00	2.25(+1)
6	4.41(-3)	-0.11	10.8	650	10.28	1.00	3.10(+1)
7	5.48(-3)	-0.20	10.8	650	—	1.00	3.85(+1)
8	6.52(-3)	-0.29	10.8	650	—	1.00	4.58(+1)
9	7.53(-3)	-0.39	10.8	650	—	1.00	5.29(+1)
10	8.52(-3)	-0.48	10.8	650	—	1.00	5.98(+1)

11	9.54(-3)	-0.57	10.8	650	-	1.00	6.70(+1)
12	1.06(-2)	-0.67	10.8	650	-	1.00	7.44(+1)
13	1.17(-2)	-0.76	10.8	650	-	1.00	8.21(+1)
14	1.29(-2)	-0.85	10.8	650	-	1.00	9.06(+1)
15	1.41(-2)	-0.94	10.8	650	-	1.00	9.90(+1)
16	1.41(-2)	-1.04	10.8	650	-	1.00	9.90(+1)

*Number in parenthesis is power of 10.

$\phi(M_v)$ \equiv luminosity function of field stars.

$\phi_{MS}(\log M) \equiv$ present-day mass function (PDMF) of main-sequence field stars in the solar neighborhood is given by

$$\phi_{MS}(\log M) = \phi(M_v) \left| \frac{dM_v}{d \log M} \right| \frac{2H(M_v)/f_{MS}(M_v)}{2H(M_v)/f_{MS}(M_v)}, \text{ where}$$

$H(M_v)$ is the scale height assuming that stars are distributed as $\exp(-|z|/H)$, where z is the distance measured perpendicular to the Galactic plane.

The factor $f_{MS}(M_v)$ gives the fraction of stars at a given magnitude that are on the main sequence.

T_{MS} = the main-sequence lifetime is given by

$$T_{MS} = \frac{\Delta X_{MS} M E}{L} \approx 13 \times 10^9 (M/M_\odot)^{-2.5} \text{ yr. } M \lesssim 10 M_\odot.$$

where

ΔX_{MS} = the mass fraction of hydrogen burned during the main-sequence phase, ~ 0.13 .

E = energy released per gram in the nuclear fusion reaction, $H \rightarrow He$, $\approx 6.4 \times 10^{18}$ erg g^{-1} .

L = total luminosity of star.

(Adapted from Shapiro, S. L. & Teukolsky, S. A., *Black Holes, White Dwarfs, and Neutron Stars*, John Wiley and Sons, 1983.)

Star number densities ($\log N_m(pg)$) with galactic latitude ^(a)

m_{pg}	Galactic latitude										Mean 0 -90
	0	± 5	± 10	± 20	± 30	± 40	± 50	± 60	± 90		
0.0		-4.0			-4.3				-4.4		-4.25
1.0		-3.4			-3.75				-3.9		-3.70
2.0		-2.83			-3.20				-3.3		-3.18
3.0		-2.32			-2.69				-2.8		-2.60
4.0	-1.75	-1.83	-1.88	-2.01	-2.16	-2.25	-2.30	-2.40			-2.11
5.0	-1.28	-1.36	-1.43	-1.56	-1.69	-1.76	-1.80	-1.89			-1.63
6.0	-0.82	-0.90	-0.97	-1.10	-1.22	-1.29	-1.34	-1.42			-1.14
7.0	-0.39	-0.46	-0.53	-0.66	-0.77	-0.84	-0.89	-0.97			-0.69
8.0	0.05	-0.01	-0.09	-0.22	-0.32	-0.40	-0.45	-0.54			-0.25
9.0	0.52	0.43	0.35	0.22	0.12	0.04	-0.01	-0.12			0.19
10.0	0.97	0.88	0.80	0.66	0.54	0.46	0.40	0.27			0.62
11.0	1.43	1.33	1.23	1.08	0.96	0.87	0.80	0.66			1.05
12.0	1.88	1.77	1.65	1.50	1.37	1.26	1.19	1.03			1.46
13.0	2.30	2.19	2.07	1.90	1.76	1.64	1.54	1.39			1.87
14.0	2.72	2.61	2.48	2.28	2.12	1.98	1.88	1.71			2.26
15.0	3.12	3.00	2.88	2.65	2.46	2.31	2.20	1.97			2.62
16.0	3.48	3.41	3.24	3.00	2.77	2.61	2.48	2.24			2.98
17.0	3.83	3.78	3.60	3.33	3.07	2.84	2.75	2.48			3.33
18.0	4.20	4.10	3.93	3.63	3.35	3.14	2.99	2.72			3.64
19.0	4.5	4.4	4.3	3.9	3.6	3.4	3.2	2.9			3.90
20.0	4.7	4.7	4.6	4.2	3.8	3.6	3.4	3.1			4.17
21.0	5.0	4.9	4.8	4.5	4.0	3.7	3.6	3.2			4.4

^(a) $N_m(pg \approx B)$ = number of stars per square degree brighter than photographic magnitude m_{pg} .

Star number densities (log $N_m(\text{vis})$ with galactic latitude^(b)) (cont.)

m_{vis}	Galactic latitude										Mean 0 -90	
	0°	±5°	±10°	±20°	±30°	±40°	±50°	±60°	±90°			
0.0		-3.9			-4.2					-4.3		-4.1
1.0		-3.3			-3.6					-3.7		-3.56
2.0		-2.7			-3.0					-3.1		-3.00
3.0		-2.14			-2.5					-2.6		-2.43
4.0	-1.55	-1.63	-1.68	-1.81	-1.96	-2.05	-2.10	-2.12	-2.20			-1.90
5.0	-1.08	-1.16	-1.23	-1.36	-1.49	-1.56	-1.60	-1.63	-1.69			-1.41
6.0	-0.60	-0.68	-0.75	-0.88	-1.00	-1.07	-1.12	-1.15	-1.20			-0.93
7.0	-0.16	-0.23	-0.30	-0.43	-0.54	-0.61	-0.66	-0.69	-0.74			0.46
8.0	0.29	0.23	0.15	0.02	-0.08	-0.16	-0.21	-0.24	-0.30			0.00
9.0	0.78	0.69	0.61	0.48	0.38	0.30	0.25	0.20	0.14			0.45
10.0	1.25	1.16	1.08	0.94	0.82	0.74	0.68	0.63	0.55			0.91
11.0	1.73	1.63	1.53	1.38	1.26	1.17	1.10	1.05	0.96			1.34
12.0	2.18	2.07	1.93	1.80	1.67	1.57	1.49	1.42	1.33			1.76
13.0	2.60	2.49	2.37	2.20	2.08	1.94	1.84	1.77	1.69			2.17
14.0	3.02	2.91	2.78	2.60	2.44	2.28	2.18	2.09	2.01			2.56
15.0	3.42	3.30	3.18	2.95	2.78	2.61	2.50	2.40	2.27			2.94
16.0	3.78	3.71	3.54	3.30	3.09	2.91	2.78	2.68	2.54			3.29
17.0	4.13	4.08	3.90	3.60	3.37	3.19	3.05	2.94	2.78			3.64
18.0	4.50	4.40	4.23	3.93	3.65	3.44	3.29	3.17	3.02			3.95
19.0	4.8	4.7	4.6	4.2	3.9	3.7	3.5	3.4	3.2			4.20
20.0	5.0	5.0	4.9	4.5	4.1	3.9	3.7	3.6	3.4			4.5
21.0	5.3	5.2	5.1	4.8	4.3	4.1	3.9	3.7	3.5			4.7

^(b) $N_m(\text{vis} \approx V)$ = number of stars per square degree brighter than visual magnitude m_{vis}
(After Allen, C. W., *Astrophysical Quantities*, Athlone Press, 1973.)

Relative numbers of stars in each class (up to $V = 8.5$ in *HD Catalog*)

Sp	O	B	A	F	G	K	M
% stars	1	10	22	19	14	31	3

(After Allen, C. W., *Astrophysical Quantities*, Athlone Press, 1973.)

Integrated star light as a function of galactic latitude


Star light (10th mag deg ⁻²)			Star light (10th mag deg ⁻²)			Star light (10th mag deg ⁻²)		
Latitude	pg	V	Latitude	pg	V	Latitude	pg	V
0	180	372	20	54	105	60	21	38
5	123	247	30	37	71	70	19	35
10	88	176	40	29	54	80	18	34
15	69	138	50	24	43	90	18	34

Integrated star light from whole sky: 230 zero pg mag stars; 460 zero V mag stars.

Night sky total brightness (zenith, mean sky) $\approx 1(m_v = 22.5)$ star arcsec⁻².

(After Allen, C. W., *Astrophysical Quantities*, Athlone Press, 1973.)

Mean star density vs. visual magnitude



Star counts

A formula for estimating ($\sim 15\%$ accuracy) differential A and integral N star counts for a given galactic longitude l , latitude b , and apparent magnitudes V and B over the ranges $b \geq 20^\circ$, $5 \leq m \leq 30$ for zero obscuration, $\Delta m = 0$, has been derived by Bahcall & Soneira (*Ap. J. Suppl.*, **44**, 73, 1980). (For non-zero obscuration, replace m by $m - \Delta m$, where $\Delta m_V = 0.15 \text{ csc } b$ and $\Delta m_B = 0.20 \text{ csc } b$.) The units of A are stars $\text{mag}^{-1} \text{ deg}^{-2}$ and of N are stars deg^{-2} .

$$D(l, b, m) = \frac{C_1 10^{\beta(m-m^*)}}{[1 + 10^{\alpha(m-m^*)}]^\lambda} \frac{1}{[\sin b(1 - \mu \cot b \cos l)]^{3-5\gamma}}$$

$$+ \frac{C_2 10^{\eta(m-m^*)}}{[1 + 10^{\kappa(m-m^*)}]^\lambda} \frac{1}{(1 - \cos b \cos l)^\sigma}$$


where the constants are $\sigma = 1.45 - 0.20 \cos b \cos l$,

Constant	Range of m		
	$m \leq 12$	$12 < m < 20$	$m \geq 20$
μ	0.03	$0.0075(m - 12) + 0.03$	0.09
γ	0.36	$0.04(12 - m) + 0.36$	0.04


Star count	C_1	C_2	α	β	δ	m^*	κ	η	λ	m^\dagger
$A_V = D$	200	400	-0.2	0.01	2	15	-0.26	0.065	1.5	17.5
$N_V = D$	925	1050	-0.132	0.035	3.0	15.75	-0.180	0.087	2.5	17.5
$A_B = D$	235	370	-0.227	0.0	1.5	17	-0.175	0.06	2.0	18
$N_B = D$	950	910	-0.124	0.027	3.1	16.60	-0.167	0.083	2.5	18

Luminosity functions

Local stellar luminosity function for the disk in the Visual band. The solid line is an analytic approximation. (Adapted from Bahcall, J. H. & Soneira, R. M., *Ap. J. Suppl.* **44**, 73, 1980.)



Local stellar luminosity function for the disk in the Blue band. The solid line is an analytic approximation. (Adapted from Bahcall, J. N. & Soneira, R. M., *Ap. J. Suppl.*, **44**, 73, 1980.)



Parameters of the interstellar gas

Mean density ρ	$3 \times 10^{-24} \text{ g cm}^{-3}$
Typical particle density	
n (HI) in diffuse clouds	20 cm^{-3}
n (HI) between clouds	0.1 cm^{-3}
n_{H} in molecular clouds	$10^3 - 10^6 \text{ cm}^{-3}$
Typical temperature T	
Diffuse HI clouds	80 K
HI between clouds	6000 K
HII photon ionized regions	8000 K
Coronal gas between clouds	$6 \times 10^5 \text{ K}$
Root-mean-square random cloud velocity	10 km s^{-1}
Isothermal sound speed C	
HI cloud at 80 K	0.7 km s^{-1}
HII gas at 8000 K	10 km s^{-1}
Magnetic field B	$2.5 \times 10^{-6} \text{ G}$
Effective thickness $2H$ of HI cloud layer	250 pc

(Adapted from Spitzer, L., *Physical Processes in the Interstellar Medium*, John Wiley and Sons, 1977.)

Galaxies

Properties of the Galaxy

Type of galaxy:

Hubble–van den Bergh system	Sb(–Sb ⁺)I–II
de Vaucouleur's system	SAB(rs)bc II
Morgan's system	gkS 7

 M_V (mag): –20.5

Diameter: 23 kpc

(isophote: 25.0 mag (B) arcsec^{–2})Period of rotation: 2.5×10^8 yr

Mass:

Total mass: $1.8 \times 10^{11} M_\odot$ Gas: $8 \times 10^9 M_\odot$ Age: 1.2×10^{10} yr

Density in solar neighborhood:

Stars: $0.05 M_\odot \text{pc}^{-3}$ Total known: $0.08 M_\odot \text{pc}^{-3}$

Galactic nucleus:

 $R < 0.4 \text{ pc} \approx 5 \times 10^6 M_\odot$ $R < 150 \text{ pc} \approx 1 \times 10^9 M_\odot$ Central bulge ($R < 2.5 \text{ kpc}$): $\approx 4 \times 10^{10} M_\odot$ *Luminosity of the galaxy:*Radio $3 \times 10^{38} \text{ erg s}^{-1}$ Infrared 3×10^{41} Optical 3×10^{43} X-ray $10^{39} - 10^{40}$ γ -ray (> 100 MeV) 5×10^{38} *Energy density in the galaxy:*Starlight $0.7 \times 10^{-12} \text{ erg cm}^{-3}$ Turbulent gas 0.5×10^{-12} Cosmic rays 2×10^{-12} Magnetic field 2×10^{-12} 2.7 K radiation 0.4×10^{-12} Mass–luminosity ratio: M/L_{bol} (solar units) ≈ 10

Stellar radiation emission (solar neighborhood):

 $1.5 \times 10^{-3} (M_{\text{bol}} = 0) \text{ stars pc}^{-3}$ $1.5 \times 10^{-23} \text{ erg cm}^{-3} \text{ s}^{-1}$

Stellar luminous radiation emission (solar neighborhood):

 $6.7 \times 10^{-4} (M_V = 0) \text{ stars pc}^{-3}$

Distance of the Sun from the galactic center:

 $8.7 \pm 0.6 \text{ kpc}$ (IAU, 1985) $7.1 \pm 1.2 \text{ kpc}$ (courtesy of M. Reid, Harvard/Smithsonian)

Height of Sun above galactic disk: 8 pc

Galactic coordinates of the nucleus: $l = -3.34$, $b = -2.75$

Equatorial coordinates of the nucleus:

 α 1950 = $17^{\text{h}} 42^{\text{m}} 29^{\text{s}} 3 \pm 0^{\text{s}} 15$ δ 1950 = $-28^\circ 59' 18'' \pm 3''$

The local group


Name	α 2000	δ 2000	V (mag)	Dimension	Type	M_v	Distance (kpc)	Diameter (kpc)
Andromeda galaxy	0 ^h 42 ^m 7	+41°16'	3.4	178' × 63'	Sb	-21.1	730	38
Milky Way	(17 45.6	-28 56)			Sb+:	-20.5	(8.5)	30:
Triangulum galaxy	1 33.9	+30 39	5.7	62 × 39	Sc	-18.9	900	16
Large Magellanic Cloud	5 23.6	-69 45	0.1	650 × 550	SBm	-18.5	50	9.5
IC 10	0 20.4	+59 18	10.3	5 × 4	Ir+:	-17.6	1300	1.9
Small Magellanic Cloud	0 52.7	-72 50	2.3	280 × 160	SBmp	-16.8	60	4.9
NGC 205	0 40.4	+41 41	8.0	17 × 10	E6:	-16.4	730	3.6
NGC 221	0 42.7	+40 52	8.2	8 × 6	E2	-16.4	730	1.7
NGC 6822	19 44.9	-14 48	9:	10 × 10	Ir+	-15.7	520	1.5
NGC 185	0 39.0	+48 20	9.2	12 × 10	dE0	-15.2	730	2.5
NGC 147	0 33.2	+48 30	9.3	13 × 8	dE4	-14.9	730	2.8
IC 1613	1 04.8	+2 07	9.3	12 × 11	Ir+	-14.8	740	2.6
WLM system	0 02.0	-15 28	10.9	12 × 4	Ir+	-14.7	1600	5.6
Leo A	9 59.4	+30 45	12.6	5 × 3	Ir+	-14.1	2300	3.3
Fornax dwarf galaxy	2 39.9	-34 32	8:	20: × 14:	dE3	-13.6	130	2.2:
IC 5152	22 02.9	-51 17	11:	5 × 3	Ir+	-13.5	1500	2.2
Pegasus dwarf galaxy	23 28.6	+14 45	12.0	5 × 3	Ir+	-13.4	1300	1.9
Sculptor dwarf galaxy	0 59.9	-33 42	10:		dE3	-11.7	85	1.5:
Leo I	10 08.4	+12 18	9.8	11 × 8	dE3	-11.0	230	0.7
Andromeda I	0 45.7	+38 00	13.2		dE0	-11:	730	
Andromeda II	1 16.4	+33 27	13:		dE0	-11:	730	
Andromeda III	0 35.4	+36 31	13:		dE2	-11:	730	
Aquarius dwarf galaxy	20 46.9	-12 51			Ir	-11:	1500	
Sagittarius dwarf galaxy	19 30.0	-17 41	15:		Ir--	-10:	1100	
Leo II	11 13.5	+22 10	11.5	15 × 13	dE0	-9.4	230	1.0
Ursa Minor dwarf galaxy	15 08.8	+67 12	12:	27 × 16	dE6	-8.8	75	0.6
Draco dwarf galaxy	17 20.2	+57 55	11:	34 × 19	dE3	-8.6:	80	0.8
LGS 3	1 03.8	+21 53	15:	2	Ir	-8.5:	900	0.5
Carina dwarf galaxy	6 41.6	-50 58			dE		170	

: denotes approximate value

(Adapted from *Sky Catalogue 2000.0*, Vol. 2, Sky Publishing Corp., 1985.)


Hubble's classification of galaxies

The number n behind the symbol E characterizes the ellipticity: $n = 10(a - b)/a$, where a and b are the major and minor diameters of the ellipse. The letters a, b, c following S and SB characterize the increasing degree of opening of the spiral arms. (Adapted from Landolt-Börnstein, *Astronomy and Astrophysics*, (1982))



Nearby extragalactic objects

Nearby extragalactic objects with our galaxy in the central position. (Adapted from Roach, F. E. & Gordon, J. L., *The Light of the Night Sky*, D. Reidel Publishing Co., 1973.)



Selected brighter galaxies ($V < 9$) (local group is excluded)

Galaxy	NGC IC	Type	l^{II} ($^{\circ}$)	b^{II} ($^{\circ}$)	Ang. ($^{\circ}$)	Diameter		$e^{(a)}$	Distance (Mpc)	V	$B - V$	M_V	Mass ($10^{11} M_{\odot}$)
						Lin. (kpc)	Ang. ($^{\circ}$)						
	55	Sc	333	-76	25	12		0.9	2.3	7.2		-19.9	0.32
	253	Sc	75	-89	22	13		0.8	2.4	7		-20	1.0
	2403	Sc	151	+28	18	11		0.4	3.2	8.4	0.5	-19.2	0.13
M81	3031	Sb	142	+41	20	16		0.5	3.2	6.9	1.0	-20.9	1.6
M82	3034	Ir II	141	+41	8	7		0.7	3	8.2	0.9	-19.6	0.32
	3115	E7	247	+37	4	5		0.7	4	9.1	1.0	-19.3	0.79
M106	4258	Sb	138	+69	15	17		0.6	4.0	8.2	0.8	-20.1	1.0
M87	4486	E1	283	+75	4	13		0.2	13	8.7	1.0	-21.7	40.0
M104 Sombrero	4594	Sa	298	+51	6	8		0.3	12	8.1	1.0	-22	5.0
M94	4736	Sb	123	+76	7	10		0.2	4.5	8.2	0.8	-20.4	1.0
M64	4826	Sb	316	+84	8	12		0.5	3.9	8.4	0.9	-19.7	0.79
	4945	Sb	305	+13	12	14		0.8	4.0	7		-21	
M63	5055	Sb	105	+74	10	15		0.5	4.6	8.4	0.9	-20.0	0.01
Cen A	5128	E0p	310	+19	14	15		0.2	4.4	7		-20	2.0
M51 Whirlpool	5194	Sc	105	+69	9	9		0.4	3.8	8.2	0.6	-19.7	0.79
M83	5236	SBc	315	+32	10	12		0.2	3.2	7.2	0.7	-20.6	
M101 Pinwheel	5457	Sc	102	+60	20	23		0.0	3.8	7.5	0.6	-20.3	1.6
	7793	Sd	4	-77	6	4		0.4	2.6	8.8		-18.4	

^(a) $e = \text{ellipticity} = (a - b)/a$, where a and b are the major and minor diameters, respectively. (After Allen, C. W., *Astrophysical Quantities*, The Athlone Press, 1973.)

Named galaxies

Name	α 2000	δ 2000
Andromeda galaxy = M31 = NGC 224	00 ^h 42 ^m 7	+41° 16'
Andromeda I	00 45.7	+38 00
Andromeda II	01 16.3	+33 25
Andromeda III	00 35.3	+36 30
Andromeda IV	00 42.5	+40 34
BL Lac	22 04.0	+42 26
Capricorn dwarf = Pal 13†	21 46.8	-21 15
Caraffe galaxy	04 28.0	-47 54
Carina dwarf	06 46.3	-51 03
Cartwheel galaxy	00 37.4	-33 45
Centaurus A = NGC 5128 = Arp 135	13 25.4	-43 02
Circinus galaxy	14 13.2	-65 20
Copeland Septet = NGC 3745/54 = Arp 320	11 37.7	+22 01
Cygnus A	19 59.4	+40 44
Draco dwarf = DDO 208	17 20.0	+57 55
Fath 703	15 13.8	-15 28
Fornax A = NGC 1316	03 22.7	-37 12
Fornax dwarf	02 39.9	-34 31
Fourcade Figueroa object	13 35.2	-33 53
GR8 = DDO 155	12 58.7	+14 13
Hardcastle Nebula	13 13.0	-32 42
Hercules A	16 51.2	+05 01
Holmberg I = DDO 63	09 40.5	+71 12
Holmberg II = DDO 50 = Arp 268	08 19.0	+70 43
Holmberg III	09 14.9	+74 14
Holmberg IV = DDO 185	13 54.6	+53 54
Holmberg V	13 40.7	+54 20
Holmberg VI = NGC 1325 A	03 24.8	-21 20
Holmberg VII = DDO 137	12 34.7	+06 18
Holmberg VIII = DDO 166	13 13.3	+36 13
Holmberg IX = DDO 66	09 57.6	+69 03
Hydra A	09 18.1	-12 06
Large Magellanic Cloud	05 23.5	-69 45
Leo I = Harrington-Wilson No. 1 = Regulus Dwarf = DDO 74	10 08.5	+12 18
Leo II = Harrington-Wilson No. 2 = Leo B = DDO 93	11 13.5	+22 10
Leo A = Leo III = DDO 69	09 59.4	+30 45
Lindsay Shapley ring	06 43.1	-74 14
Maffei I	02 36.3	+59 39
Maffei II	02 41.9	+59 36
Mayall's Object = Arp 148 = VV 32	11 03.9	+40 51
Mice = NGC 4676 = Arp 242	12 47.1	+30 38
Pegasus dwarf = DDO 216	23 28.5	+14 44
Perseus A = NGC 1275	03 19.8	+41 31
Reticulum dwarf	04 36.2	-58 50
Reinmuth 80 = NGC 4517 A	01 00.0	-33 42

Named galaxies (cont.)

Name	α 2000	δ 2000
Seashell galaxy	13 ^h 47 ^m 3	-30°25'
Serpens dwarf	15 16.0	-00 08
Seyfert's Sextet = NGC 6027 A-D	15 59.2	+20 45
Sextans A = DDO 75	10 11.1	-04 43
Sextans B = DDO 70	10 00.0	+05 20
Sextans C	10 05.5	+00 04
Small Magellanic Cloud	00 52.8	-72 50
Sombrero galaxy = M104 = NGC 4594	12 40.2	-11 37
Stephan's Quintet = NGC 7317-20 = Arp 319	22 36.0	+33 58
Triangulum galaxy = M33 = NGC 598	01 34.5	+30 39
Ursa Minor dwarf = DDO 199	15 08.8	+67 12
Virgo A = M87 = NGC 4486 = Arp 152	12 30.8	+12 23
Whirlpool galaxy = M51 = NGC 5194	13 29.9	+47 12
Wild's Triplet = Arp 248	11 46.8	-03 50
Wolf-Lundmark-Melotte object = DDO 221	00 02.0	-15 27
Zwicky No. 2 = DDO 105	11 58.5	+38 04
Zwicky's Triplet = Arp 103	16 49.5	+45 28

† Probably a distant globular cluster.

(Adapted from Landolt-Börnstein, *Astronomy and Astrophysics*, VI/2C, Springer-Verlag, 1982.)

Representative active galactic nuclei (AGNs)

Object	α 1950	δ 1950	z	m_v
QUASARS				
Q 0002-422	00 ^h 02 ^m 16 ^s	-42°14'	2.758	17.4
PHL 938	00 58 20	01 55	1.95	17.2
4C 25.05	01 23 57	25 44	2.34	17.5
PHL 1093	01 37 23	01 17	0.262	17.1
PHL 1194	01 48 52	09 03	0.298	17.5
RN 8	02 10 49	86 05	0.184	19.0
Q 0242-410	02 42 02	-41 04	2.214	18.1
Q 0324-407	03 24 29	-40 47	3.056	17.6
PKS 0424-13	04 24 48	-13 10	2.16	17.5
Q 0453-423	04 53 48	-42 21	2.661	17.3
Q 0551-366	05 51 02	-36 38	2.307	17.0
OH 471	06 42 53	44 55	3.39	18.5
PKS 0736+01	07 36 43	01 44	0.192	16.5
4C 05.34	08 05 19	04 41	2.86	18.2
0938+119	09 38 32	11 59	3.19	19.0
3C 232	09 55 25	32 38	0.533	15.8
Ton 490	10 11 06	25 04	1.63	15.4
PKS 1217+02	12 17 39	02 20	0.240	16.5
3C 273	12 26 33	02 20	0.158	12.8
Q 1246-057	12 46 29	-05 43	2.212	17.0
B 340	13 04 48	34 40	0.184	17.0
1331+170	13 31 10	17 04	2.08	16.0

Representative active galactic nuclei (cont.)

Object	α 1950	δ 1950	z	m_v
3C 323.1	15 ^h 45 ^m 31 ^s	21°01'	0.264	16.7
4C 29.50	17 02 11	29 51	1.92	19.1
3C 351	17 04 03	60 49	0.371	15.3
Q 2116-358	21 16 22	-35 49	2.341	17.0
PKS 2135-14	21 35 01	-14 46	0.200	15.5
2256+017	22 56 25	01 48	2.66	18.5
SEYFERT GALAXIES				
<i>Seyfert 1 galaxies</i>				
Mrk 335	00 03 45	19 55	0.025	14.2
I Zw 1	00 51 00	12 25	0.061	14.3
Mrk 376	07 10 36	45 47	0.056	16.0
Mrk 79	07 38 47	49 56	0.020	13.4
Mrk 10	07 43 07	61 03	0.029	15.0
Mrk 110	09 21 44	52 30	0.036	16.1
NGC 3227	10 20 47	20 07	0.0033	13.5
NGC 3516	11 03 24	72 50	0.0093	13.1
NGC 4151	12 08 01	39 41	0.0033	12.0
Mrk 236	12 58 18	61 55	0.052	17.0
Mrk 279	13 51 52	69 33	0.0307	15.4
Mrk 290	15 34 45	58 04	0.0308	15.6
Mrk 486	15 35 21	54 43	0.039	15.0
Mrk 509	20 41 26	-10 54	0.0355	13.0
NGC 7469	23 00 44	08 36	0.0167	13.6
Mrk 541	23 53 30	07 15	0.041	15.5
<i>Seyfert 2 galaxies</i>				
Mrk 1	01 13 19	32 50	0.016	16.6
NGC 1068	02 40 07	-00 14	0.003 63	10.5
Mrk 612	03 21 10	-03 19	0.020 22	16.5
III Zw 55	03 38 38	-01 28	0.0246	14.0
Mrk 3	06 09 48	71 03	0.0137	13.8
Mrk 78	07 37 56	65 18	0.0375	15.6
Mrk 622	08 04 21	39 09	0.022 83	15.6
Mrk 34	10 30 52	60 17	0.051	14.8
Mrk 176	11 29 54	53 14	0.0269	15.5
Mrk 270	13 39 41	56 55	0.009	15.0
Mrk 463E	13 53 40	18 37	0.0505	16.0
Mrk 533	23 25 24	08 30	0.028 73	16.0
BL LAC OBJECTS				
PKS 0215+015	02 15 13	01 31		18.3
AO 0235+164	02 35 53	16 24		15.5
PKS 0521-365	05 21 14	-36 30	0.55	15.0
PKS 0548-323	05 48 50	-32 17	0.069	15.5
OJ 287	08 51 57	20 18		14.0
4C 22.25	09 57 34	22 48		18.0
Mkn 421	11 01 41	38 29	0.308	13.5
Mkn 180	11 33 30	70 25	0.0458	15.0
AP Lib	15 14 45	-24 11	0.049	15.0
Mkn 501	16 52 12	39 50	0.034	13.8
BL Lac	22 00 40	42 02	0.0688	14.5

Representative active galactic nuclei (cont.)

Object	α 1950	δ 1950	z	m_v
RADIO GALAXIES				
<i>BLRGs</i>				
3C 109	04 ^h 10 ^m 55 ^s	11 15'	0.306	18.0
3C 120	04 30 32	05 15	0.033	14.6
3C 227	09 45 07	07 39	0.0855	16.3
3C 234	09 58 57	29 02	0.1846	17.1
3C 287.1	13 29 04	25 24	0.2156	18.5
PKS 1417-19	14 17 02	-19 15	0.1195	17.5
4C 35.37	15 31 45	35 52	0.1565	17.5
3C 332	16 14 44	30 09	0.1515	16.0
3C 381	18 32 28	47 24	0.1614	17.5
3C 382	18 33 12	32 39	0.0586	15.4
3C 390.3	18 45 38.8	79 43	0.0569	15.4
3C 445	22 21 15	-02 21	0.0568	15.8
<i>NLRGs</i>				
3C 33	01 06 14	13 04	0.0595	16.3
3C 98	03 56 10	10 18	0.0306	14.8
3C 178	07 22 33	-09 30	0.0079	16.1
3C 184.1	07 32 20	70 20	0.1182	17
3C 192	08 02 38	24 16	0.0598	16.2
3C 327	15 59 56	02 06	0.1039	16.3
3C 433	21 21 30	24 52	0.1025	15.7
3C 452	22 43 33	39 25	0.082	16.6
PKS 2322-12	23 22 43	-12 24	0.0821	15.8
LINERS				
Mrk 1158	01 32 07	34 47	0.0151	16.2
NGC 1052	02 38 37	-08 28	0.0048	13.2
Ark 160	08 17 52	19 31	0.019	17.6
NGC 2841	09 18 35	51 11	0.0022	13.5
NGC 2911	09 31 05	10 23	0.0106	15.3
NGC 3031	09 51 30	69 18	-0.0001	12.4
NGC 3758	11 33 48	21 52	0.0296	16.6
NGC 3998	11 55 20	55 44	0.0038	13.3
NGC 4036	11 58 54	62 10	0.0046	14.0
NGC 4278	12 17 36	29 34	0.0022	13.6
NGC 5005	13 08 37	37 19	0.0033	14.1
NGC 5077	13 16 53	-12 24	0.0094	14.4
NGC 5371	13 53 33	40 42	0.0086	15.0
Mrk 298	16 03 18	17 56	0.0345	16.2
Mrk 700	17 01 21	31 31	0.034	15
NGC 6764	19 07 01	50 51	0.008	15.5

Redshift $z = \Delta\lambda/\lambda$.

m_v = approximate nuclear visual magnitude.

Luminosity distance $D_L(q_0=0) = \frac{cz}{H_0} (1+0.5z)$. For other values of q_0 see chapter on Relativity.

(Adapted from Landolt-Börnstein, *Astronomy and Astrophysics*, V1/2C, Springer-Verlag, 1982.)

Objects with large redshifts

Object	α 1950	δ 1950	z (em)	V
1208 + 1011	12 ^h 08 ^m 23.73	+10 11' 07".9	3.80	
PKS 2000 - 330	20 00 12.94	-33 00 14.6	3.78	19.0
0055 - 2659	00 55 32.46	-25 29 26.0	3.67	17.1
DHM 0054 - 284	00 53 59.8	-28 24 45	3.61	19.55
OQ 172	14 42 50.48	10 11 12.2	3.53	17.78
1159 + 1223	11 59 14.23	+12 23 11.9	3.51	17.5
222738 - 3928	22 27 38	-39 28 32	3.45	16.8
OH 471	06 42 53.02	-44 54 31.1	3.40	18.49
0042 - 2627	00 42 06.22	-26 27 42.9	3.30	18.5
OX - 146			3.27	
2016 + 112†	20 16 55	11 17 46	3.27	22.5
OS 023			3.21	
024929 - 1826	02 49 29.8	-18 26 21	3.21	18.6
OQ 004			3.20	
024925 - 2212	02 49 25.5	-22 12 37	3.20	18.4
Q 0938 + 119	09 38 31.75	11 59 12.6	3.19	19

† Gravitational lens.

Note added in proof: Over 20 quasars are now known with redshifts greater than 3.5 (see *Sky & Telescope*, 75, 12, 1988).


Prominent clusters of galaxies

Name	Abell No.	α 2000	δ 2000	Diameter (")	RV (km s ⁻¹)	RS type	NGC	Radio source	Notes
Haufen A	151	1 ^h 08 ^m 9	-15° 25'		15 800	cD			
	194	1 25.6	-1 30	0.3	5320	L	541, 5, 7	3C 40	In Perseus supercluster
	400	2 57.6	+6 02		7200	I		3C 75	
Perseus	426	3 18.6	+41 32	4	5460	L	1275	3C 84	In Perseus supercluster: XRS
Fornax II		3 28	-20 45	7	1560		1232		
Fornax I		3 32	-35 20	7	1500		1316	For A	
Gemini	568	7 07.6	+35 03	0.5	23 400	C			
Cancer		8 21	+20 56	3	4800		2563		
Hydra II		8 58	+3 09		60 900				
Leo	1020	10 27.8	+10 25	0.6	19 500				Also Abell 1016?
Hydra I	1060	10 36.9	-27 32		3000	C	3309, 11		XRS
Ursa Major II		10 58	+56 46	0.2	41 000				
Leo A	1185	11 10.9	+28 41		10 500	C	3550		
	1367	11 44.5	+19 50		6150	F	3842, 62	3C 264	In Coma supercluster; XRS
Ursa Major I	1377	11 47.1	+55 44	0.7	15 300	B			
Virgo		12 30	+12 23	12	1200			Vir A	In Local supercluster; XRS
Centaurus		12 50	-41 18	2	3200		4472, 86	PKS	XRS
Coma	1656	12 59.8	+27 59	4	6650	B	4696		
Boötes	1930	14 33	+31 33	0.3	39 300		4889		In Coma supercluster; XRS
Corona Borealis	2065	15 22.7	+27 43	0.5	21 600	C			
Hercules	2151	16 05.2	+17 45	1.7	11 200	F	6040, 47	4C +17.66	In Hercules supercluster; XRS
	2152	16 05.4	+16 27		11 500	~I			In Hercules supercluster
	2197	16 28.2	+40 54		9100	L	6173		In Hercules supercluster
	2199	16 28.6	+39 31	0.2	9200	cD	6166	3C 338	In Hercules supercluster; XRS
Pegasus II		23 10	+7 36	2	12 700		7720	4C +07.61	
Pegasus I		23 22	+9 02	1	4000		7619	PKS	

$z = RV/3.00 \times 10^5$; distance (Mpc) = $RV/50$ ($H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$).
 (Adapted from *Sky Catalogue 2000.0*, Vol. 2, Sky Publishing Corp., 1985.)

Prominent clusters of galaxies (cont.)

Extragalactic systems within about 150 megaparsecs.




(Adapted from Roach, F. E. & Gordon, J. L. *The Light of the Night Sky*.
D. Reidel Publishing Co., 1973.)

Energy content of constituents of space

Constituent	Energy content (average all space including intergalactic space)	Primary sources
A. Photons:		
1. Infraradio	?	?
2. Radio and microwave (exclude 3 K radiation)	$1 \mu\text{eV cm}^{-3}$	Galaxy, radio galaxies, quasars
3. 3 K radiation	0.4 eV cm^{-3}	Big bang
4. Infrared	? may be $> 3 \text{ K}$ radiation	Seyfert galaxies, quasars, N galaxies
5. Visible light	0.01 eV cm^{-3}	Stars in normal galaxies
6. Soft X-rays ($< 1 \text{ keV}$)	$(0.3 \text{ eV cm}^{-3}$ in galaxy)	ISM and discrete sources
7. Hard X-rays (1–500 keV)	$50\text{--}100 \mu\text{eV cm}^{-3}$	Hot plasma: inverse Compton scattering of fast electrons with background radiation: discrete sources (quasars)
8. Gamma rays ($E > 35 \text{ MeV}$)	$2 \mu\text{eV cm}^{-3}$	Uncertain, probably active galaxies
B. Magnetic fields	0.001 eV cm^{-3}	?
C. Cosmic rays and fast electrons	$100 \mu\text{eV cm}^{-3}$	Radio galaxies, quasars, supernovae in normal galaxies
D. Neutrinos	0.1 eV cm^{-3} (?)	Big bang
E. Gravitons	0.1 eV cm^{-3} (?)	Big bang
F. Matter (ρc^2) ($\rho = 2 \times 10^{-31} \text{ g cm}^{-3}$)	100 eV cm^{-3}	Galaxies

Energy density of cosmic photons

R = radio
 M = microwave
 IR = infrared
 O = optical
 UV = ultraviolet
 SXR = soft X-ray
 XR = X-ray
 γ = gamma ray



(Adapted from Rose, W. K., *Astrophysics*, Holt, Rinehart, and Winston, Inc., 1973.)

Mass–radius–density data for astronomical objects

Class of objects	Examples	$\log M$ (g)	$\log R$ (cm)	$\log \rho$ (g cm ⁻³)	$\log \phi$ †
Neutron stars		33.16	5.93	14.75	-0.6?
		32.54	7.44	9.60	-2.5
White dwarfs	L930-80	33.45	8.3:	7.93	-2.7
	α CMaB	33.30	8.77	6.37	-3.2
	vM2	32.90	9.05	4.13	-5.0
Main sequence stars	dM8	32.2	9.95	1.76	-5.6
	Sun	33.30	10.84	0.15	-5.5
	A0	33.85	11.25	-0.55	-4.7
	O5	34.9	12.1:	-2.0	-5.0:
Supergiant stars	F0	34.4	12.65	-4.2	-6.1
	K0	34.4	13.15	-5.7	-6.6
	M2	34.7	13.75	-7.2	-6.9
Protostars	IR	35.3?	16.2?	-13.9?	-8.7?
Compact dwarf elliptical galaxies	M32, core	41.0	19.5?	-18.1	-6.3
	M32, effective	42.5	20.65	-20.0	-5.9
	N4486-B	43.4	20.5	-18.75	-5.0
Spiral galaxies	LMC	43.2	21.75	-22.65	-6.3
	M33	43.5	21.8	-22.5	-6.1
	M31	44.6	22.3	-22.9	-5.5
Giant elliptical galaxies	N3379	44.3	22.0	-22.35	-5.6
	N4486	45.5	22.4	-22.3	-4.7
Compact groups of galaxies	Stephan	45.5	22.6:	-23.1:	-4.7
Small groups of spirals	Sculptor	46.2	24.1	-26.7	-5.7
Dense groups of ellipticals	Virgo E, core Fornax I	46.5	23.7	-25.2	-5.0
Small clouds of galaxies	Virgo S	47.0	24.3	-26.5	-5.1
	Ursa Major				
Small clusters of galaxies	Virgo F	47.2	24.3	-26.3	-4.9
Large clusters of ellipticals	Coma	48.3	24.6	-26.1	-4.9
Superclusters	Local	48.7:	25.5:	-28.4:	-4.7
	HMS sample to $m \simeq 12.5$		26.0:	-29.6	-4.6
	Lick Observatory counts to $m \simeq 19.0$		26.8	-30.5	-4.1

: denotes approximate value; ? denotes large uncertainty

† The filling factor $\phi = \rho/\rho_m$, where $\rho_m = 3c^2/8\pi GR_m^2$; $R_m = 2GM/c^2$.

(Adapted from de Vaucouleurs, G., *Science*, **167**, 1203, 1970.)

The Messier catalog

M	NGC	α 2000	δ 2000	Const.	Dim. (')	V (mag)	Type	Common name
1	1952	5 ^h 34 ^m 05	+22 01'	Tau	6 × 4	8.4:	Di	Crab Nebula
2	7089	21 33.5	-0 49	Aqr	13	6.5	Gb	
3	5272	13 42.2	+28 23	CVn	16	6.4	Gb	
4	6121	16 23.6	-26 32	Sco	26	5.9	Gb	
5	5904	15 18.6	+2 05	Ser	17	5.8	Gb	
6	6405	17 40.1	-32 13	Sco	15	4.2	OC	
7	6475	17 53.9	-34 49	Sco	80	3.3	OC	
8	6523	18 03.8	-24 23	Sgr	90 × 40	5.8:	Di	Lagoon Nebula
9	6333	17 19.2	-18 31	Oph	9	7.9:	Gb	
10	6254	16 57.1	-4 06	Oph	15	6.6	Gb	
11	6705	18 51.1	-6 16	Sct	14	5.8	OC	
12	6218	16 47.2	-1 57	Oph	14	6.6	Gb	
13	6205	16 41.7	+36 28	Her	17	5.9	Gb	Hercules Cluster
14	6402	17 37.6	-3 15	Oph	12	7.6	Gb	
15	7078	21 30.0	+12 10	Peg	12	6.4	Gb	
16	6611	18 18.8	-13 47	Ser	7	6.0	OC	
17	6618	18 20.8	-16 11	Sgr	46 × 37	7:	Di	Omega Nebula
18	6613	18 19.9	-17 08	Sgr	9	6.9	OC	
19	6273	17 02.6	-26 16	Oph	14	7.2	Gb	
20	6514	18 02.6	-23 02	Sgr	29 × 27	8.5:	Di	Trifid Nebula
21	6531	18 04.6	-22 30	Sgr	13	5.9	OC	
22	6656	18 36.4	-23 54	Sgr	24	5.1	Gb	
23	6494	17 56.8	-19 01	Sgr	27	5.5	OC	
24		18 16.9	-18 29	Sgr	90	4.5:		
25	IC 4725	18 31.6	-19 15	Sgr	32	4.6	OC	
26	6694	18 45.2	-9 24	Sct	15	8.0	OC	
27	6853	19 59.6	+22 43	Vul	8 × 4	8.1:	Pl	Dumbbell Nebula
28	6626	18 24.5	-24 52	Sgr	11	6.9:	Gb	
29	6913	20 23.9	+38 32	Cyg	7	6.6	OC	
30	7099	21 40.4	-23 11	Cap	11	7.5	Gb	
31	224	0 42.7	+41 16	And	178 × 63	3.4	S	Andromeda Galaxy
32	221	0 42.7	+40 52	And	8 × 6	8.2	E	
33	598	1 33.9	+30 39	Tri	62 × 39	5.7	S	
34	1039	2 42.0	+42 47	Per	35	5.2	OC	
35	2168	6 08.9	+24 20	Gem	28	5.1	OC	
36	1960	5 36.1	+34 08	Aur	12	6.0	OC	
37	2099	5 52.4	+32 33	Aur	24	5.6	OC	
38	1912	5 28.7	+35 50	Aur	21	6.4	OC	

The Messier catalog (*cont.*)

M	NGC	α 2000	δ 2000	Const.	Dim. (')	V (mag)	Type	Common name
39	7092	21 ^h 32 ^m 2	+48° 26'	Cyg	32	4.6	OC	
40		12 22.4	+58 05	UMa		8:		
41	2287	6 47.0	-20 44	CMa	38	4.5	OC	
42	1976	5 35.4	-5 27	Ori	66 × 60	4:	Di	Orion Nebula
43	1982	5 35.6	-5 16	Ori	20 × 15	9:	Di	
44	2632	8 40.1	+19 59	Cnc	95	3.1	OC	Praesepe
45		3 47.0	+24 07	Tau	110	1.2	OC	Pleiades
46	2437	7 41.8	-14 49	Pup	27	6.1	OC	
47	2422	7 36.6	-14 30	Pup	30	4.4	OC	
48	2548	8 13.8	-5 48	Hya	54	5.8	OC	
49	4472	12 29.8	+8 00	Vir	9 × 7	8.4	E	
50	2323	7 03.2	-8 20	Mon	16	5.9	OC	
51	5194-5	13 29.9	+47 12	CVn	11 × 8	8.1	S	Whirlpool Galaxy
52	7654	23 24.2	+61 35	Cas	13	6.9	OC	
53	5024	13 12.9	+18 10	Com	13	7.7	Gb	
54	6715	18 55.1	-30 29	Sgr	9	7.7	Gb	
55	6809	19 40.0	-30 58	Sgr	19	7.0	Gb	
56	6779	19 16.6	+30 11	Lyr	7	8.2	Gb	
57	6720	18 53.6	+33 02	Lyr	1	9.0:	Pl	Ring Nebula
58	4579	12 37.7	+11 49	Vir	5 × 4	9.8	S	
59	4621	12 42.0	+11 39	Vir	5 × 3	9.8	E	
60	4649	12 43.7	+11 33	Vir	7 × 6	8.8	E	
61	4303	12 21.9	+4 28	Vir	6 × 5	9.7	S	
62	6266	17 01.2	-30 07	Oph	14	6.6	Gb	
63	5055	13 15.8	+42 02	CVn	12 × 8	8.6	S	
64	4826	12 56.7	+21 41	Com	9 × 5	8.5	S	
65	3623	11 18.9	+13 05	Leo	10 × 3	9.3	S	
66	3627	11 20.2	+12 59	Leo	9 × 4	9.0	S	
67	2682	8 50.4	+11 49	Cnc	30	6.9	OC	
68	4590	12 39.5	-26 45	Hya	12	8.2	Gb	
69	6637	18 31.4	-32 21	Sgr	7	7.7	Gb	
70	6681	18 43.2	-32 18	Sgr	8	8.1	Gb	
71	6838	19 53.8	+18 47	Sge	7	8.3	Gb	
72	6981	20 53.5	-12 32	Aqr	6	9.4	Gb	
73	6994	20 58.9	-12 38	Aqr				
74	628	1 36.7	+15 47	Psc	10 × 9	9.2	S	
75	6864	20 06.1	-21 55	Sgr	6	8.6	Gb	
76	650-1	1 42.4	+51 34	Per	2 × 1	11.5:	Pl	
77	1068	2 42.7	-0 01	Cet	7 × 6	8.8	S	
78	2068	5 46.7	+0 03	Ori	8 × 6	8:	Di	
79	1904	5 24.5	-24 33	Lep	9	8.0	Gb	
80	6093	16 17.0	-22 59	Sco	9	7.2	Gb	

The Messier catalog (*cont.*)

M	NGC	α 2000	δ 2000	Const.	Dim. (')	V (mag)	Type	Common name
81	3031	9 ^h 55 ^m 6	+69°04'	UMa	26 × 14	6.8	S	
82	3034	9 55.8	+69 41	UMa	11 × 5	8.4	Ir	
83	5236	13 37.0	-29 52	Hya	11 × 10	7.6	S	
84	4374	12 25.1	+12 53	Vir	5 × 4	9.3	E	
85	4382	12 25.4	+18 11	Com	7 × 5	9.2	E	
86	4406	12 26.2	+12 57	Vir	7 × 6	9.2	E	
87	4486	12 30.8	+12 24	Vir	7	8.6	E	Virgo A
88	4501	12 32.0	+14 25	Com	7 × 4	9.5	S	
89	4552	12 35.7	+12 33	Vir	4	9.8	E	
90	4569	12 36.8	+13 10	Vir	10 × 5	9.5	S	
91	4548	12 35.4	+14 30	Com	5 × 4	10.2	S	
92	6341	17 17.1	+43 08	Her	11	6.5	Gb	
93	2447	7 44.6	-23 52	Pup	22	6.2	OC	
94	4736	12 50.9	+41 07	CVn	11 × 9	8.1	S	
95	3351	10 44.0	+11 42	Leo	7 × 5	9.7	S	
96	3368	10 46.8	+11 49	Leo	7 × 5	9.2	S	
97	3587	11 14.8	+55 01	UMa	3	11.2	PI	Owl Nebula
98	4192	12 13.8	+14 54	Com	10 × 3	10.1	S	
99	4254	12 18.8	+14 25	Com	5	9.8	S	
100	4321	12 22.9	+15 49	Com	7 × 6	9.4	S	
101	5457	14 03.2	+54 21	UMa	27 × 26	7.7	S	
102								M101 reobser- vation
103	581	1 33.2	+60 42	Cas	6	7.4	OC	
104	4594	12 40.0	-11 37	Vir	9 × 4	8.3	S	Sombrero Galaxy
105	3379	10 47.8	+12 35	Leo	4 × 4	9.3	E	
106	4258	12 19.0	+47 18	CVn	18 × 8	8.3	S	
107	6171	16 32.5	-13 03	Oph	10	8.1	Gb	
108	3556	11 11.5	+55 40	UMa	8 × 2	10.0	S	
109	3992	11 57.6	+53 23	UMa	8 × 5	9.8	S	
110	205	0 40.4	+41 41	And	17 × 10	8.0	E?	

: denotes approximate value.

Types: diffuse nebula (Di), globular cluster (Gb), open cluster (OC), planetary nebula (PI), or galaxy (E for elliptical, Ir for irregular, S for spiral).

Magnitudes with a colon are approximate visual magnitudes.

(Adapted from *Sky Catalogue 2000.0*, Vol. 2, Sky Publishing Corp., 1985.)

Astronomical photometry

Following M. Golay (*Introduction to Astronomical Photometry*, D. Reidel Publishing Company, 1974), we can write the following expression for the apparent magnitude difference on the Earth of two stars:

$$m_1 - m_2 = -2.5 \log \frac{\int_{\lambda_a}^{\lambda_b} \alpha_1^2 I_1(\lambda) T_i(\lambda, d_1) T_a(\lambda, d_1) T_t(\lambda) T_f(\lambda) r(\lambda) d\lambda}{\int_{\lambda_a}^{\lambda_b} \alpha_2^2 I_2(\lambda) T_i(\lambda, d_2) T_a(\lambda, d_2) T_t(\lambda) T_f(\lambda) r(\lambda) d\lambda},$$

where

$I_1(\lambda)$ the spectral radiance of star 1.

$I_2(\lambda)$ the same for star 2.

α_1 and α_2 the apparent diameters of stars 1 and 2, which are assumed to be spherical and emit isotropic radiation.

$T_i(\lambda, d_1)$ the fraction of the radiation of star 1 transmitted by interstellar space in the direction d_1 of star 1.

$T_i(\lambda, d_2)$ the same for star 2.

$T_a(\lambda, d_1)$ the fraction of stellar radiation transmitted by the Earth's atmosphere when star 1 is in direction d_1 .

$T_a(\lambda, d_2)$ the same for star 2 when it is in direction d_2 .

$T_t(\lambda)$ the fraction of stellar radiation transmitted by the optical system of the telescope t , whose entry pupil is perpendicular to the star's direction.

$T_f(\lambda)$ the fraction of stellar radiation transmitted by a filter f placed in front of the receiver.

$r(\lambda)$ the response of the receiver r which, for simplicity, is assumed to depend only upon λ .

The limits of integration, λ_a and λ_b , where $\lambda_b > \lambda_a$ are defined by

$$\lambda \geq \lambda_b \quad T_a \cdot T_t \cdot T_f \cdot r \equiv 0,$$

$$\lambda \leq \lambda_a \quad T_a \cdot T_t \cdot T_f \cdot r \equiv 0.$$

Let

$$S(\lambda) = T_t(\lambda) T_f(\lambda) r(\lambda),$$

the response of the photometric system and

$$E(\lambda) = \frac{\alpha^2}{4} I(\lambda) T_i(\lambda, d),$$

the stellar spectral irradiance at the top of the Earth's atmosphere, then the difference in apparent magnitudes for two stars outside the Earth's atmosphere is given by the expression:

$$(m_1 - m_2)_0 = -2.5 \log \frac{\int_{\lambda_a}^{\lambda_b} E_1(\lambda) \cdot S(\lambda) d\lambda}{\int_{\lambda_a}^{\lambda_b} E_2(\lambda) \cdot S(\lambda) d\lambda}.$$

We can define three wavelengths for a photometric system:

$$(1) \quad \lambda_0 = \frac{\int_{\lambda_a}^{\lambda_b} \lambda S(\lambda) d\lambda}{\int_{\lambda_a}^{\lambda_b} S(\lambda) d\lambda},$$

the mean wavelength of the pass-band defined by the response function $S(\lambda)$.

$$(2) \quad E(\lambda_i) \int_{\lambda_a}^{\lambda_b} S(\lambda) d\lambda = \int_{\lambda_a}^{\lambda_b} E(\lambda) S(\lambda) d\lambda,$$

where λ_i is the isophotal wavelength.

$$(3) \quad \lambda_{\text{eff}} = \frac{\int_{\lambda_a}^{\lambda_b} \lambda E(\lambda) S(\lambda) d\lambda}{\int_{\lambda_a}^{\lambda_b} E(\lambda) S(\lambda) d\lambda},$$

the effective wavelength.

$$\lambda_{\text{eff}} - \lambda_0 = \frac{E'(\lambda_0)}{E(\lambda_0)} \cdot \mu^2 \quad \text{and} \quad \lambda_i - \lambda_0 = \frac{1}{2} \mu^2 \frac{E''(\lambda_0)}{E'(\lambda_0)},$$

where

$$\mu^2 = \frac{\int (\lambda - \lambda_0)^2 S(\lambda) d\lambda}{\int S(\lambda) d\lambda}.$$

The color index of a star is defined by

$$C_{AB} \equiv m_A - m_B = -2.5 \log \frac{\int_A E(\lambda) S_A(\lambda) d\lambda}{\int_B E(\lambda) S_B(\lambda) d\lambda} + \text{constant},$$

where A and B represent two different spectral bands.

The relationships between heterochromatic magnitude m_{λ_i} (obtained with a band of mean wavelength λ_0) and monochromatic magnitudes taken at the isophotal wavelength, $m(\lambda_i)$, at the mean wavelength $m(\lambda_0)$ and at the effective wavelength $m(\lambda_{\text{eff}})$:

$$m_{\lambda_i} = m(\lambda_i) + S, \quad \text{where } S = -2.5 \log \int_{\lambda_a}^{\lambda_b} S(\lambda) d\lambda.$$

$$m(\lambda_i) - m(\lambda_0) = -0.543 \mu^2 \left(\frac{E''(\lambda_0)}{E(\lambda_i)} \right).$$

$$m(\lambda_i) - m(\lambda_{\text{eff}}) = -0.543 \left(\frac{\lambda_{\text{eff}} E'(\lambda_{\text{eff}})}{E(\lambda_{\text{eff}})} + 1 \right) \frac{\lambda_0 - \lambda_{\text{eff}}}{\lambda_{\text{eff}}}$$

Standard photometric systemsStandard *U, B, V, R, I* and long wavelength systems

Filter band	$\lambda_0^{(a)}$ (μm)	$\Delta\lambda_0$ (FWHM) (μm)	Absolute spectral irradiance for mag = 0.0	
			$f_\lambda(0)$ ($\text{erg cm}^{-2} \text{s}^{-1} \text{\AA}^{-1}$)	$f_\nu(0)$ ($\text{W m}^{-2} \text{Hz}^{-1}$)
<i>U</i>	0.365	0.068	4.27×10^{-9}	1.90×10^{-23}
<i>B</i>	0.44	0.098	6.61×10^{-9}	$4.27(4.64)^{(b)} \times 10^{-23}$
<i>V</i>	0.55	0.089	3.64×10^{-9}	3.67×10^{-23}
<i>R</i>	0.70	0.22	1.74×10^{-9}	2.84×10^{-23}
<i>I</i>	0.90	0.24	8.32×10^{-10}	2.25×10^{-23}
<i>J</i>	1.25	0.3	3.18×10^{-10}	1.65×10^{-23}
<i>H</i>	1.65	0.4	1.18×10^{-10}	1.07×10^{-23}
<i>K</i>	2.2	0.6	4.17×10^{-11}	6.73×10^{-24}
<i>L</i>	3.6	1.2	6.23×10^{-12}	2.69×10^{-24}
<i>M</i>	4.8	0.8	2.07×10^{-12}	1.58×10^{-24}
<i>N</i>	10.2		1.23×10^{-13}	4.26×10^{-25}

^(a) $\lambda_0 = \int \lambda S(\lambda) d\lambda / \int S(\lambda) d\lambda$, where $S(\lambda)$ is the photometer response function.

^(b) From S. Kleinmann.

U, B, R, I, N values from Allen, C. W., *Astrophysical Quantities*. The Athlone Press (1973). *V, J, H, K, L, M* values from Wamsteker, W., *Astron. Astrophys.*, **97**, 329 (1981).

The spectral irradiance for a star of a given magnitude is given either by:

$$\log f_\lambda(m_x) = -0.4m_x + \log f_\lambda(0),$$


where $f_\lambda(m_x)$ is the spectral irradiance in $\text{erg cm}^{-2} \text{s}^{-1} \text{\AA}^{-1}$ of a star of magnitude (m_x) in the x filter band at the mean wavelength $\lambda_0(x)$, or

$$\log f_\nu(m_x) = -0.4m_x + \log f_\nu(0),$$


where $f_\nu(m_x)$ is the spectral irradiance in $\text{W m}^{-2} \text{Hz}^{-1}$.

The relationships above are for the irradiance at the top of the Earth's atmosphere and are valid for *B* through *M* stars.

Photometer response curves for *UBVRI* and long wavelength systems. (Adapted from Webbink, R. F. & Jeffers, W. Q., *Space Sci. Rev.*, **10**, 191 1969.)



Response curves of photometer plus atmosphere. (Adapted from Webbink, R. F. & Jeffers, W. Q., *Space Sci. Rev.*, **10**, 191, 1969.)



List of UB_V primary standard stars

HD No.	Name	<i>V</i>	<i>B</i> - <i>V</i>	<i>U</i> - <i>B</i>	Spectral type
12 929	α Ari	2.00	+1.151	+1.12	K2 III
18 331	HR 875	5.17	+0.084	+0.05	A1 V
69 267	β Cnc	3.52	+1.480	+1.78	K4 III
74 280	η Hya	4.30	-0.195	-0.74	B3 V
135 742	β Lib	2.61	-0.108	-0.37	B8 V
140 573	α Ser	2.65	+1.168	+1.24	K2 III
143 107	ϵ CrB	4.15	+1.230	+1.28	K3 III
147 394	τ Her	3.89	-0.152	-0.56	B5 IV
214 680	10 Lac	4.88	-0.203	-1.04	O9 V
219 134	HR 8832	5.57	+1.010	+0.89	K3 V

(List taken from Strand, K. A. A., ed., *Basic Astronomical Data*, University of Chicago Press, Chicago, 1963.)


Standard stars for the JHKLM system

Standard	BS	Spectral type	m_{vis}	J	H	K	L	M
0	519	gM 4	5.49	2.117	1.317	1.078	0.890	1.181
1	721	B5 III	4.25	4.548	4.575	4.604	4.601	4.689
2	1195	G5 III	4.17	2.672	2.233	2.119	2.025	2.156
3	2827	B5 Ia	2.44	2.565	2.548	2.557	2.472	2.542
4	4216	G5 III	2.69	1.148	0.744	0.622	0.530	0.673
5	5530	F5 IV	5.16	4.408	4.194	4.156	4.092	4.134
6	7120	K3 III	4.98	2.822	2.197	2.052	1.910	2.103
7	8204	G4 Ibp	3.74	2.346	1.964	1.865	1.794	1.882
8	8502	K3 III	2.85	0.558	-0.077	-0.091	-0.372	-0.199
9	8728	A3 V	1.16	1.075	1.034	1.019	0.998	1.025

(List taken from Wamstecker, W., *Astron. Astrophys.*, **97**, 329, 1981.)

Kron photographic J and F bands

Sensitivity functions $S_j(\lambda)$ for the *J* waveband (left) and the *F* waveband (right). (From Kron, R. G., *Ap. J.*, **43**, 305, 1980.)



The earlier photovisual (m_{pv}) and photographic (m_{pg}) magnitudes are related to the standard *B* and *V* magnitudes by:

$$B \equiv m_B = m_{pg} + 0.11,$$

$$V \equiv m_V = m_{pv} + 0.00.$$

Color index,

$$C = m_{pg} - m_V = B - V - 0.11.$$

Bolometric correction,

$$BC = m_b - m_V = M_b - M_V,$$

where m_b (M_b) is the apparent (absolute) bolometric magnitude, a measure of the total energy output of a star.

$$M_b^{\text{star}} - 4.72 = -2.5 \log(L_{\text{star}}/L_{\odot}),$$

where L_{star} and $L_{\odot} = 3.83 \times 10^{33} \text{ erg s}^{-1}$ are the absolute luminosities of the star and the Sun, respectively.

$$L_{\text{star}} = 2.97 \times 10^{35} \times 10^{-0.4M_b} \text{ erg s}^{-1}.$$

The total irradiance at the top of the Earth's atmosphere is

$$f = 2.48 \times 10^{-5} \times 10^{-0.4m_b} \text{ erg cm}^{-2} \text{ s}^{-1}$$

for a star of apparent bolometric magnitude m_b .

Assuming black-body radiation, the spectral photon irradiance from a star of apparent bolometric magnitude m_b is given by:

$$f(\lambda) = \frac{8.48 \times 10^{34} \times 10^{-0.4m_b}}{T_e^4 \lambda^3 [\exp(1.44 \times 10^8 / \lambda T_e) - 1]} \text{ photons cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$$

λ in \AA ; T_e the effective temperature of the star in K (e.g., A0 star; $m_V = 0$, $T_e = 10800$, $BC = -0.40$, $\lambda = 5000 \text{ \AA}$; $f(\lambda) = 10^3 \text{ photons cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$).

Interstellar reddening

The observed color index is given by

$$C_{ij} = C_{ij}^0 + [A(\lambda_i) - A(\lambda_j)] \equiv C_{ij}^0 + E_{ij},$$

where

$A(\lambda)$ = amount of interstellar absorption at λ ,

C_{ij}^0 = intrinsic color index of the star,

E_{ij} \equiv color excess.

In the UBV system, the color excesses are

$$E(B - V) \equiv (B - V) - (B - V)_0,$$

$$E(U - B) \equiv (U - B) - (U - B)_0$$

(subscript zero denotes intrinsic values).

$$A_V / E(B - V) = 3.2 \pm 0.2 \quad (\text{normal regions})$$

$$\frac{E(U - B)}{E(B - V)} = 0.72 + 0.05E(B - V).$$

Relationship of reddening $E(B - V)$ to the hydrogen column density:

$$\langle N(\text{HI} + \text{H}_2) / E(B - V) \rangle = 5.8 \times 10^{21} \text{ atoms cm}^{-2} \text{ mag}^{-1},$$

$$\langle N(\text{HI}) / E(B - V) \rangle = 4.8 \times 10^{21} \text{ atoms cm}^{-2} \text{ mag}^{-1}$$

(Bohlin *et al.*, *Ap. J.*, **224**, 132, 1978).

Visual extinction to the galactic center:

$$A_V \approx 30 \text{ mag}$$

(Becklin *et al.*, *Ap. J.*, **151**, 145, 1968).

The mean color excess $\bar{E}_{B-V}(b)$ at galactic latitude b for objects outside the absorbing layer can be estimated by:


$$\bar{E}_{B-V}(b) = 0.06 \operatorname{cosec}|b| - 0.06$$

(Woltjer, L., *Astron. Astrophys.*, **42**, 109, 1975).

A more general expression giving an estimate of interstellar absorption can be found in de Vaucouleurs *et al.*, *Second Reference Catalogue of Bright Galaxies*, University of Texas Press, 1976.

The interstellar reddening law

λ (Å)	$1/\lambda$ (μ^{-1})	$A_\lambda/E(B-V)$	λ (Å)	$1/\lambda$ (μ^{-1})	$A_\lambda/E(B-V)$
3200	3.13	5.58	5556	1.80	3.12
3250	3.08	5.47	5840	1.71	2.91
3300	3.03	5.38	6050	1.65	2.80
3350	2.98	5.32	6436	1.55	2.65
3400	2.94	5.26	6790	1.47	2.48
3448	2.90	5.18	7100	1.41	2.35
3509	2.85	5.10	7550	1.32	2.18
3571	2.80	5.06	7780	1.28	2.06
3636	2.75	4.99	8090	1.24	1.98
3862	2.59	4.72	8188	1.22	1.94
4036	2.48	4.56	8370	1.20	1.88
4167	2.40	4.39	8446	1.18	1.85
4255	2.35	4.33	8710	1.15	1.72
4464	2.24	4.12	9700	1.03	1.50
4566	2.19	4.04	9832	1.02	1.48
4780	2.09	3.91	10 256	0.95	1.36
5000	2.00	3.70	10 610	0.94	1.28
5263	1.90	3.41	10 796	0.93	1.26
5480	1.82	3.20	10 870	0.92	1.22



(Courtesy of R. Schild, Center for Astrophysics.)

Absolute magnitude

The absolute magnitude M of a star is the apparent magnitude it would have if placed at a distance of 10 parsecs:

$$m - M = 5 \log D - 5 + A,$$

where D is the distance to the star in parsecs and A is a correction for interstellar absorption expressed in magnitude units. $m - M \equiv$ distance modulus. Solving for D :

$$D = 10^{[1 + (m - M - A) / 5]}.$$

Moon, night sky, sun, and planetary brightness

Moon

$$V(R, \phi) = 0.23 + 5 \log R - 2.5 \log P(\phi),$$

$V(R, \phi)$ = the apparent V magnitude of the Moon,

R = the observer-Moon distance in AU, and

ϕ = the phase angle = angle between the Sun and the Earth as seen from the Moon.

$$P(0^\circ) = 1.000, \quad P(40^\circ) = 0.377, \quad P(80^\circ) = 0.127,$$


$$P(120^\circ) = 0.027, \quad P(160^\circ) = 0.001.$$

Mean lunar distance = 2.570×10^{-3} AU.

The V magnitude of the Moon at the Earth at opposition (full moon) is -12.73 .

(Adapted from Wertz, J. R., *Spacecraft Attitude Determination and Control*, D. Reidel, 1980)

The Moon's phase law.



Night sky

Total brightness (zenith, mean sky) ≈ 1 ($m_v = 22.5$) star arcsec $^{-2}$.

Sun

<i>Apparent magnitude</i>	<i>Color index</i>	<i>Absolute magnitude</i>
$U = -26.06$	$U - B = +0.10$	$M_U = +5.51$
$B = -26.16$	$B - V = +0.62$	$M_B = +5.41$
$V = -26.78$	$BC = -0.07$	$M_V = +4.79$
$m_b = -26.85$		$M_b = +4.72$

Planetary brightness

The change in the brightness of a planet because of the changing distance from the Sun (r) and the Earth (Δ) is given by:

$$V = V(1, 0) + 5 \log(r\Delta) + \alpha(p),$$

where

$V(1, 0)$ = visual magnitude of planet reduced to a distance of 1 AU from both the Sun and the Earth and phase angle $p = 0$,

α = phase law; change of planet brightness with p ,

p = phase angle; angle between Sun and Earth seen from the planet

$$\cos p = \frac{r^2 + \Delta^2 - R^2}{2r\Delta},$$

where R = distance from the Earth to the Sun.

<i>Planet</i>	$V(1, 0)^{(a)}$	$\alpha^{(b)}$
Mercury	-0.42 mag	$+0.027p + 2.2 \times 10^{-13}p^6$
Venus	-4.40	$+0.013p + 4.2 \times 10^{-7}p^3$
Mars	-1.52	$+0.016p$
Jupiter	-9.40	$+0.014p$
Saturn	-8.88	$+0.044L - 2.6 \sin B + 1.2 \sin^2 B$
Uranus	-7.19	$+0.001p$
Neptune	-6.87	$+0.001p$
Pluto	-1.0	

p in degrees; L = Saturnicentric ring longitude difference of Sun and Earth; B = Saturnicentric ring latitude of Earth; $0^\circ < L < 6^\circ$, $0^\circ < |B| < 27^\circ$.

^(a) (from the *The Nautical Almanac*)

^(b) (from Allen, C. W., *Astrophysical Quantities*)

Spherical astronomy

=

Time

The *Julian date (JD)* corresponding to any instant is the interval in *mean solar days* since 4713 BC January 1 at Greenwich mean noon (1200 UT). (Midnight, January 1, 1961 = 0000 UT January 1, 1961 = JD 2,437,300.5.)

A procedure (from Fliegel, H. F. & van Flinders, T. C., *Comm. ACM*, 11, 657, 1968) for finding the Julian date (*JD*) for a given year (*I*), month (*J*), and day of the month (*K*) is given by the following FORTRAN (integer) arithmetic statement:

$$JD = K - 32075 + 1461 * (I + 4800 + (J - 14) / 12) / 4 \\ + 367 * (J - 2 - (J - 14) / 12 * 12) / 12 \\ - 3 * ((I + 4900 + (J - 14) / 12) / 100) / 4.$$

For example, December 25, 1981 ($I = 1981$, $J = 12$, $K = 25$) = *JD* 2,444,964.

One *Besselian year* is the period of a complete circuit of the mean Sun in right ascension beginning at the instant when its right ascension is $18^{\text{h}} 40^{\text{m}}$. The epochs to which stellar coordinates are referred are in Besselian year numbers. (The epoch 1950.0 started December 31, 1949 at 2209 UT.)

A *mean sidereal day* is the interval between two successive upper culminations or transits of the vernal equinox.

The *civil* or *mean solar day* is $\frac{1}{365.2422}$ of a *tropical year*, the interval between two successive passages of the Sun through the vernal equinox.

Sidereal time is the hour angle of the vernal equinox.

Apparent solar time is the local hour angle of the Sun, expressed in hours, plus 12 hours.

Mean solar time is the local hour angle, plus 12 hours, of a fictitious *mean sun* which moves along the equator at a constant rate equal to the average annual rate of the Sun.

Mean solar time at 0° longitude is called *universal time (UT)*, formerly *Greenwich mean time* or GMT).

Ephemeris time (ET) is based on the time interval of a *tropical year*. 1 *ephemeris second* is $\frac{1}{31\,556\,925.9747}$ of the tropical year 1900.

$$ET = UT + \Delta T \quad (\Delta T = +52.5 \text{ s, January 1, 1982}).$$

International atomic time = $ET - 32.18 \text{ s}$.

In 1981:

$$1 \text{ mean solar day} = 1.002\,737\,909\,31 \text{ mean sidereal days} \\ = 24^{\text{h}} 03^{\text{m}} 56^{\text{s}} 555\,36 \text{ of mean sidereal time.}$$

Pacific standard time = $UT - 8^h$.
Mountain standard time = $UT - 7^h$.
Central standard time = $UT - 6^h$.
Eastern standard time = $UT - 5^h$.
Colonial standard time = $UT - 4^h$.
Western European time = UT .
Central European time = $UT + 1^h$.
Eastern European time = $UT + 2^h$.

Notation for time-scales

A summary of the notation for time-scales and related quantities used in the *Astronomical Almanac* is given below. Additional information is given in the *Supplement to the Almanac*.

- UT = UT1; universal time; counted from 0^h at midnight; unit is mean solar day.
 UT0 local approximation to universal time; not corrected for polar motion.
 GMST Greenwich mean sidereal time; GHA of mean equinox of date.
 GAST Greenwich apparent sidereal time; GHA of true equinox of date.
 TAI international atomic time; unit is the SI second.
 UTC coordinated universal time; differs from TAI by an integral number of seconds, and is the basis of most radio time signals and legal time systems.
 $\Delta UT = UT - UTC$; increment to be applied to UTC to give UT.
 DUT = predicted value of ΔUT , rounded to 0.1, given in some radio time signals.
 ET ephemeris time; was used in dynamical theories and in the *Almanac* from 1960–83; but is now replaced by TDT and TDB.
 TDT terrestrial dynamical time; used as time-scale of ephemerides for observations from the Earth's surface. $TDT = TAI + 32^s.184$.
 TDB barycentric dynamical time; used as time-scale of ephemerides referred to the barycenter of the solar system.
 $\Delta T = ET - UT$ (prior to 1984); increment to be applied to UT to give ET.
 $\Delta T = TDT - UT$ (1984 onwards); increment to be applied to UT to give TDT.
 $\Delta T = TAI + 32^s.184 - UT$.
 $\Delta AT = TAI - UTC$; increment to be applied to UTC to give TAI.
 $\Delta ET = ET - UTC$; increment to be applied to UTC to give ET.
 $\Delta TT = TDT - UTC$; increment to be applied to UTC to give TDT.

For most purposes, ET up to 1983 December 31 and TDT from 1984 January 1 can be regarded as a continuous time-scale. Values of ΔT for the years 1620 onwards are given in the *Astronomical Almanac*.

The name Greenwich mean time (GMT) is not used in astronomy since it is ambiguous and is now used, in the sense of UTC in addition to the earlier sense of UT; prior to 1925 it was reckoned for astronomical purposes from Greenwich mean noon (12^h UT).


Relationships with local time and hour angle

The following general relationships are used:

- Local mean solar time = universal time + east longitude.
 Local mean sidereal time = Greenwich mean sidereal time + east longitude.
 Local apparent sidereal time = local mean sidereal time + equation of equinoxes
 = Greenwich apparent sidereal time + east longitude.
 Local hour angle = local apparent sidereal time – apparent right ascension
 = local mean sidereal time – (apparent right ascension – equation of equinoxes).


A further small correction for the effect of polar motion is required in the production of very precise observations.

The celestial sphere



(Adapted from Valley, S. L., ed., *Handbook of Geophysics and Space Environment*, AFCRL, 1965.)

Celestial coordinates




Celestial coordinates

(Unsoeld, A., *The New Cosmos*, Springer-Verlag, 1969, with permission.)

The Zodiac

Path of the Earth around the Sun, seasons, and Zodiac. Perihelion is 2 January and aphelion is 2 July. Right ascension in hours of each constellation is given. (Adapted from Unsoeld, A., *The New Cosmos*, Springer-Verlag, 1969.)



Astronomical coordinate transformations

Horizon-equatorial (celestial) systems

$$\begin{aligned} \cos a \sin A &= +\cos \delta \sin h, \\ \cos a \cos A &= -\sin \delta \cos \varphi + \cos \delta \cos h \sin \varphi, \\ \sin a &= \sin \delta \sin \varphi + \cos \delta \cos h \cos \varphi, \\ \cos \delta \sin h &= \cos a \sin A, \\ \cos \delta \cos h &= \sin a \cos \varphi + \cos a \cos A \sin \varphi, \\ \sin \delta &= \sin a \sin \varphi - \cos a \cos A \cos \varphi, \\ h &= \text{local sidereal time} - \alpha, \\ A &= \text{azimuth, toward West from South,} \\ a &= \text{altitude,} \\ \varphi &= \text{observer's latitude,} \\ h &= \text{local hour angle,} \\ \alpha &= \text{right ascension,} \\ \delta &= \text{declination.} \end{aligned}$$

Ecliptic-equatorial (celestial) systems


$$\begin{aligned} \cos \delta \cos \alpha &= \cos \beta \cos \lambda, \\ \cos \delta \sin \alpha &= \cos \beta \sin \lambda \cos \varepsilon - \sin \beta \sin \varepsilon, \\ \sin \delta &= \cos \beta \sin \lambda \sin \varepsilon + \sin \beta \cos \varepsilon, \\ \cos \beta \cos \lambda &= \cos \delta \cos \alpha, \\ \cos \beta \sin \lambda &= \cos \delta \sin \alpha \cos \varepsilon + \sin \delta \sin \varepsilon, \\ \sin \beta &= \sin \delta \cos \varepsilon - \cos \delta \sin \alpha \sin \varepsilon, \\ \alpha &= \text{right ascension, } \delta = \text{declination,} \\ \lambda &= \text{ecliptic longitude, } \beta = \text{ecliptic latitude,} \\ \varepsilon &= \text{obliquity of the ecliptic} = 23^\circ 27' 8''.26 - 46''.845T \\ &\quad - 0''.0059T^2 + 0''.00181T^3 \\ &\text{where } T \text{ is the time in centuries from 1900.} \end{aligned}$$

Galactic-equatorial (celestial) systems

$$\begin{aligned} \cos b^{\text{II}} \cos(l^{\text{II}} - 33^\circ) &= \cos \delta \cos(\alpha - 282.25^\circ), \\ \cos b^{\text{II}} \sin(l^{\text{II}} - 33^\circ) &= \cos \delta \sin(\alpha - 282.25^\circ) \cos 62.6^\circ + \sin \delta \sin 62.6^\circ, \\ \sin b^{\text{II}} &= \sin \delta \cos 62.6^\circ - \cos \delta \sin(\alpha - 282.25^\circ) \sin 62.6^\circ, \\ \cos \delta \sin(\alpha - 282.25^\circ) &= \cos b^{\text{II}} \sin(l^{\text{II}} - 33^\circ) \cos 62.6^\circ - \sin b^{\text{II}} \sin 62.6^\circ, \\ \sin \delta &= \cos b^{\text{II}} \sin(l^{\text{II}} - 33^\circ) \sin 62.6^\circ + \sin b^{\text{II}} \cos 62.6^\circ, \\ l^{\text{II}} &= \text{new galactic longitude,} \\ b^{\text{II}} &= \text{new galactic latitude,} \\ \alpha &= \text{right descension (1950.0),} \\ \delta &= \text{declination (1950.0),} \\ \text{For, } l^{\text{II}} = b^{\text{II}} = 0: \alpha &= 17^{\text{h}} 42^{\text{m}} 4, \delta = -28^\circ 55' \text{ (1950.0);} \\ b^{\text{II}} = +90.0, \text{ galactic north pole: } \alpha &= 12^{\text{h}} 49^{\text{m}}, \delta = +27^\circ 4 \text{ (1950.0).} \end{aligned}$$


Galactic-equatorial (celestial) systems (cont.)

Chart for conversion of equatorial (1950.0) coordinates into new galactic coordinates ($l^{\text{II}}b^{\text{II}}$) or vice versa.
 (Kraus, J. D., *Radio Astronomy*, 2nd edn., with permission.)




Approximate reduction of astronomical coordinates

Right-ascension precession in seconds of time per year.



Declination precession in seconds of arc per year (*left*) and minutes of arc as a function of interval in years (*right*).



Precession charts: The charts show the precession in right ascension in seconds of time per year and of declination in seconds of arc per year as a function of position as given by the relations

$$\Delta\alpha = 3^m07 + 1^s34 \sin \alpha \tan \delta \quad \text{per year,}$$

$$\Delta\delta = 20'' \cos \alpha \quad \text{per year.}$$

The lower chart also indicates the precession in declination in minutes of arc as a function of the interval in years. For example, to find the precession for an object at RA = 04^h00^m for an 18-year interval one enters the chart at 4 hr, finding a precession of 10 sec of arc per year. Then moving horizontally to the right-hand chart to a point above 18 years the precession is found to be 3 min of arc for the 18-year interval.

The charts are useful for approximate precession determinations for intervals between epochs 1800.0 to 2000.0. (Kraus, J. D., *Radio Astronomy*, 2nd edn., with permission.)

Major ground-based astronomical facilities

Clear aperture	Focal length	Prime focal ratio	In oper.	Latitude	Longitude	Elevation	Facility	
600 cm	236 in	24.0 m	4.0	1976	43° 39' N	41° 26' E	2070 m	Bol'shoi Teleskop Azimutal'nyi Special Astrophysical Observatory Mount Pastukhov, USSR
508	200	16.8	3.3	1948	33° 21' N	116° 52' W	1706	George Ellery Hale Telescope Palomar Observatory
450	177	4.9	2.7	1979	31° 41' N	110° 53' W	2600	Palomar Mountain, California Multiple Mirror Telescope Mount Hopkins Observatory
400	158	11.2	2.8	1976	30° 10' S	70° 48' W	2399	Mount Hopkins, Arizona Cerro-Tololo Inter-American Observatory
389	153	12.7	3.3	1975	31° 17' S	149° 04' E	1164	Cerro Tololo, Chile Anglo-Australian Telescope Anglo-Australian Observatory
381	150	10.7	2.8	1973	31° 58' N	111° 36' W	2064	Siding Spring Mountain, Australia Nicholas U. Mayall Reflector Kitt Peak National Observatory Kitt Peak, Arizona
380	150	9.5	2.5	1979	19° 50' N	155° 28' W	4194	United Kingdom Infrared Telescope Royal Observatory, Edinburgh Mauna Kea, Hawaii
360	142	13.5	3.8	1979	19° 49' N	155° 28' W	4200	Canada-France-Hawaii Telescope Mauna Kea, Hawaii

357	141	10.9	3.0	1976	29° 15' S	70° 44' W	2400	ESO 3.6-meter Telescope European Southern Observatory Cerro La Silla, Chile
350	138	12.2, 13.8	3.5, 4.0	1983	37° 14' N	02° 32' W	2168	Max Planck Institute for Astronomy German-Spanish Astronomical Center Calar Alto, Spain
305	120	15.2	5.0	1959	37° 20' N	121° 39' W	1277	C. Donald Shane Telescope Lick Observatory Mount Hamilton, California

SCHMIDT TELESCOPES

Clear aperture	Mirror diameter	Focal length	Focal ratio	In oper.	Latitude	Longitude	Elevation	Facility	
134 cm	53 in	200 cm	4.0 m	3.0	1960	50° 59' N	11° 43' E	331 m	2-meter Telescope of the Karl Schwarzschild Observatorium near Tautenberg, East Germany
126	49.5	183	3.1	2.4	1948	33° 21' N	116° 52' W	1706	Palomar Schmidt Telescope Palomar Observatory Palomar Mountain, California
124	49	183	3.1	2.5	1973	31° 16' S	149° 04' E	1131	United Kingdom Schmidt Telescope Royal Observatory, Edinburgh
105	41	150	3.3	3.1	1976	35° 48' N	137° 38' E	1130	Siding Spring Mountain, Australia Kiso Station of the Tokyo Astronomical Observatory Kiso, Japan
100	39	162	3.0	3.1	1972	29° 15' S	70° 44' W	2400	ESO 1-meter Schmidt Telescope European Southern Observatory Cerro La Silla, Chile

Major ground-based astronomical facilities (cont.)

SCHMIDT TELESCOPES

Clear aperture	Mirror diameter	Focal length	Focal ratio	In oper.	Latitude	Longitude	Elevation	Facility
100 cm	39 in	152 cm	3.0	1978	8° 47' N	70° 52' W	3600 m	Centro de Investigación de Astronomía 'F. J. Duarte' Llano del Hato, Venezuela
100	39	135	3.0	1963	59° 30' N	17° 36' E	33	Kvistaberg Schmidt Telescope Uppsala University Observatory Kvistaberg, Sweden
100	39	132	2.1	1961	40° 20' N	44° 30' E	1450	3TA-10 Schmidt Telescope Byurakan Astrophysical Observatory
90	35	152	3.5	1981	43° 45' N	6° 56' E	1270	Mount Aragatz, Armenian SSR
84	33	120	2.5	1958	50° 48' N	4° 21' E	105	Télescope de Schmidt Observatoire de Calem Calem, France
								Télescope Combiné de Schmidt Observatoire Royal de Belgique Uccle, Bruxelles, Belgium

REFRACTING TELESCOPES

Clear aperture	Focal length	Prime ratio	In oper.	Latitude	Longitude	Elevation	Facility
101 cm	40 in	19.2	1897	42° 34' N	88° 33' W	334 m	Yerkes Observatory University of Chicago Williams Bay, Wisconsin

89.5	36	17.6	19.7	1888	37° 20' N	121° 39' W	1284	36-inch Refractor Lick Observatory
83	33	16.2	19.5	1889	48° 48' N	2° 14' E	162	Mount Hamilton, California Observatoire de Paris
80	31	12.0	15.0	1899	52° 23' N	13° 04' E	107	Meudon, France Zentralinstitut für Astrophysik
76	30	14.1	18.6	1914	40° 29' N	80° 01' W	370	Telegrafenberg, Potsdam, E. Germany The Thaw Refractor
74	29	17.9	24.2	1886	43° 43' N	7° 18' E	376	Allegheny Observatory River View Park, Pittsburgh, PA Lunette Bischoffsheim
71	28	8.5	11.9	1894	51° 29' N	0° 00'	47	Observatoire de Nice Mont Gros, France 28-inch Visual Refractor
68	27	21.0	30.9	1896	52° 29' N	13° 29' E	41	Old Royal Greenwich Observatory Greenwich, England Grosser Refraktor
67	26	10.5	15.7	1880	48° 14' N	16° 20' E	240	Archenhold-Sternwarte Alt Treptow, Berlin, E. Germany Grosser Refraktor
67	26	10.9	16.3	1925	26° 11' S	28° 04' E	1806	Institut für Astronomie Vienna, Austria The Innes Telescope Outstation South African Astronomy Observatory
67	26.3	9.9	14.9	1883	38° 02' N	78° 31' W	259	Johannesburg, South Africa Leander McCormick Observatory Mount Jefferson Charlottesville, Virginia

Major ground-based astronomical facilities (cont.)

REFRACTING TELESCOPES

Clear aperture	Focal length	Prime ratio	In oper.	Latitude	Longitude	Elevation	Facility
66 cm	9.9 m	15.0	1873	38° 55' N	77° 04' W	92 m	26-inch Equatorial United States Naval Observatory Washington, DC
66	6.9	10.4	1899	50° 52' N	0° 20' E	34	The Thompson Refractor Royal Greenwich Observatory Herstmonceux Castle, Hailsham, UK
66	10.8	16.4	1925	35° 19' S	149° 00' E	769	Yale-Columbia Refractor Mt. Stromlo and Siding Springs Observatory, Mt. Stromlo, ACT, Australia

RADIO TELESCOPES/ANTENNAE

Aperture	Frequency range (MHz)	Latitude	Longitude	Elevation	Facility
305 m dish (non-steerable)	50-3000	18° 21' N	66° 45' W	365 m	Arecibo Observatory Arecibo, Puerto Rico
100 m dish (steerable)	600-15 000	50° 32' N	6° 53' E	366	MPI für Radioastronomie Effelsberg, Federal Republic of Germany
14 25 m dishes (syn. tel.)	610, 1415, 4995	52° 55' N	6° 36' E	5	Westerbork Radio Observatory Hooghalen, The Netherlands
64 m dish (steerable)	500-5000	33° 00' S	148° 16' E	392	Australian National Radio Astronomy Observatory, Parkes, New South Wales, Australia

Rectangular arrays	1415	33° 52' S	150° 46' E	Radio Astronomy Observatory Fleurs, New South Wales, Australia
Radioheliograph	80, 160, 327	30° 19' S	150° 46' E	CSIRO Solar Radio Observatory Culgoora, New South Wales, Australia
76 m dish (steerable)	0.3–4000	53° 14' N	2° 18' W	Nuffield Radio Astronomy Laboratories Jodrell Bank, Macclesfield Cheshire, England
8 element interferometer (5 km baseline)	2695, 5000, 15 375	52° 10' N	0° 03' E	Mullard Radio Astronomy Observatory Cambridge, England
40 m dish (steerable)	500–12 000 18 000–24 000	37° 14' N	118° 17' W	Owens Valley Radio Observatory Big Pine, California
91.4 m dish (steerable)†	0.3–5000	38° 26' N	79° 50' W	National Radio Astronomy Observatory Greenbank, West Virginia
45.7 m dish (steerable)	400–10 000	45° 57' N	78° 04' W	Algonquin Radio Observatory Lake Traverse, Ontario, Canada
V(ery) L(arge) A(rray)	1420, 1700, 5000, 15 000, 24 010	34° 05' N	107° 37' W	National Radio Astronomy Observatory Plains of San Augustin, New Mexico
27–25 m dishes (steerable)	$\lambda_{\min} = 1.4$ mm	37° 02' N	03° 12' W	Institut de Radio Astronomie Millimétrique Lomo de Dilar, Granada, Spain
30 m dish				Science Research Council Nagano, Japan
45 m dish	$\lambda_{\min} = 3$ mm	36° N	138° E	

(Adapted from Classen, J. & Sperling, N., *Sky and Telescope*, **61**, 303, 1981.)

† Note added in proof: Structure collapsed on November 15, 1988.

The Hubble Space Telescope

(Material for this section was taken from STScI document SC-02, 'Call for proposals', October, 1985.)

The Hubble Space Telescope (HST) is managed by the Space Telescope Science Institute (STScI):


Space Telescope Science Institute
3700 San Martin Drive
Baltimore, MD 21218
USA
Tel.: (301) 338-4700

To allow European astronomers to make use of the Hubble Space Telescope, the European Space Agency (ESA) has established the Space Telescope-European Coordinating Facility (ECF). The address of the ST-ECF:

Space Telescope-European Coordinating Facility
European Southern Observatory
Karl-Schwarzschild-Str. 2
D-8046 Garching bei München
Federal Republic of Germany
Tel.: (89) 32-006-291
Telex: 05-28282-22 EO D
Telefax: +49-89-3202362

An overview of the HST dealing with the performance of the telescope and instruments and plans for observations and data reduction and analysis can be found in the NASA publication, *The Space Telescope Observatory*, special session of commission 44, IAU 18th General Assembly, 1982, NASA CP-2244.

Description of the Hubble Space Telescope



HST nominal orbital parameters


Altitude	500 km
Inclination	28.5
Orbital period	95 min
Orbital precession period	54 days

Optical characteristics of HST


Optical design	Ritchey-Chrétien
Aperture	2.4 m
Collecting area	38 993 cm ²
Wavelength coverage (MgF ₂ -overcoated aluminum)	1150 Å–1 mm
Focal ratio	f/24
Plate scale	3.58 arcsec mm ⁻¹
Predicted FWHM of on-axis images, including 0.007 rms pointing jitter:	
1220 Å	0.023 arcsec
2000 Å	0.025 arcsec
3500 Å	0.035 arcsec
4500 Å	0.041 arcsec
6328 Å	0.056 arcsec
Predicted radius of 70% encircled energy (at 6328 Å)	0.10 arcsec

Description of the Hubble Space Telescope (cont.)

Predicted HST throughput.



Predicted on-axis HST encircled energy at various wavelengths.



The following six scientific instruments will be available on HST:

- Wide Field and Planetary Camera (WF/PC)
- Faint Object Camera (FOC)
- Faint Object Spectrograph (FOS)
- High Resolution Spectrograph (HRS)

Description of the Hubble Space Telescope (cont.)

- High Speed Photometer (HSP)
- Fine Guidance Sensors (FGS)

*HST instrument capabilities*Direct imaging⁽¹⁾

Instrument	Field of view	Projected pixel spacing on sky	Wavelength range (Å)	Magnitude limit ⁽²⁾
WFC	154" × 154"	0".10	1150–11 000	28
PC	66" × 66"	0".043	1150–11 000	28
FOC <i>f</i> /48	44" × 44"	0".043	1150–6500	27
<i>f</i> /96	22" × 22"	0".022	1150–6500	27
<i>f</i> /288	7".3 × 7".3	0".0072	1150–6500	26

Astrometry

Instrument	Field of view	Positional accuracy	Wavelength range (Å)	Magnitude limit ⁽³⁾
FGS	69 (arcmin) ²	± 0".0016 ⁽⁴⁾	4700–7100	17

Photometry

Instrument	Aperture diameters	Time resolution	Wavelength range (Å)	Magnitude limit ⁽⁵⁾
HSP ⁽⁶⁾	0".4, 1".0	10 μs	1200–9000	25

Slitless spectroscopy⁽⁷⁾

Instrument	Projected pixel spacing on sky	Resolving power ($\lambda/\Delta\lambda$)	Wavelength range (Å)	Magnitude limit
WFC	0".10	100	1600–4000	22 ⁽⁸⁾
		40	1300–2000	21 ⁽⁸⁾
		45	3000–6000	23 ⁽²⁾
		35	6000–10 000	23 ⁽²⁾
FOC <i>f</i> /48	0".043	50 at 1500 Å	1150–6000	22 ⁽⁹⁾
	<i>f</i> /96	0".022	50 at 1500 Å	1150–6000

*Description of the Hubble Space Telescope (cont.)**HST instrument capabilities (cont.)*

Slit spectroscopy

Instrument	Projected aperture size	Resolving power ($\lambda/\Delta\lambda$)	Time resolution	Wave-length range (Å)	Magnitude limit ⁽²⁾
FOC $f/48$ ⁽¹⁰⁾	0".1 × 20"	2000		1150–1325	18
		2000		1167–1767	18
		2000		1750–2650	20
		2000		3500–5300	21
FOS ⁽¹¹⁾	0".1–4".3	1300	20 ms	1150–8500	18–22–22
		250	20 ms	1150–8500	21–26–23
HRS	0".25, 2".0	100 000	50 ms	1150–3200	11–14
		20 000	50 ms	1150–3200	13–16
		2000	50 ms	1150–1800	17

Notes to tables:

⁽¹⁾ The cameras also have coronagraphic and polarimetric imaging capabilities.

⁽²⁾ Predicted limiting V magnitude for an unreddened A0 V star in order to achieve a S/N ratio of 5 in an exposure time of 1 hour. Single entries refer to wavelengths near the center of the indicated wavelength range. For FOC direct imaging, the F342W filter was assumed. When two values are given, the first refers to wavelengths near the short-wavelength limit, and the second to those near the long-wavelength limit. For FOS spectroscopy, three values are given, corresponding to 1300, 3200, and 6000 Å, respectively.

⁽³⁾ For $S/N = 1$ in fine lock.

⁽⁴⁾ Within the central area of 4' × 5'. In-orbit performance for astrometry is very uncertain; figure quoted is design goal.

⁽⁵⁾ Predicted limiting V magnitude for a 1-hour exposure (half on star, half on sky) on an A0 V star at $S/N = 5$ with the F450W filter.

⁽⁶⁾ The HSP can also perform polarimetry.

⁽⁷⁾ Using 'objective' gratings or prisms.

⁽⁸⁾ Predicted limiting V magnitude for an unreddened B0 V star in order to achieve a S/N ratio of 5 in an exposure time of 1 hour.

⁽⁹⁾ Predicted limiting V magnitude for a QSO with a ν^{-2} spectrum in order to achieve a S/N ratio of 10 in an exposure time of 1 hour with the far-UV objective prism (PRISM2).

⁽¹⁰⁾ Two-dimensional spectroscopy with the long slit and grating.

⁽¹¹⁾ The FOS can also perform spectropolarimetry.

Glossary of astronomical terms

aberration: the apparent angular displacement of the observed position of a celestial object from its *geometric position*, caused by the finite velocity of light in combination with the motions of the observer and of the observed object. (See *aberration, planetary*.)

- aberration, annual*: the component of stellar aberration (see *aberration, stellar*) resulting from the motion of the Earth about the Sun.
- aberration, diurnal*: the component of stellar aberration (see *aberration, stellar*) resulting from the observer's diurnal motion about the center of the Earth.
- aberration, E-terms of*: terms of annual aberration (see *aberration, annual*) depending on the *eccentricity* and longitude of *perihelion* of the Earth.
- aberration, elliptic*: see *aberration, E-terms of*.
- aberration, planetary*: the apparent angular displacement of the observed position of a celestial body produced by motion of the observer (see *aberration, stellar*) and the actual motion of the observed object (see *correction for light-time*).
- aberration, secular*: the component of stellar aberration (see *aberration, stellar*) resulting from the essentially uniform and rectilinear motion of the entire solar system in space. Secular aberration is usually disregarded.
- aberration, stellar*: the apparent angular displacement of the observed position of a celestial body resulting from the motion of the observer. Stellar aberration is divided into diurnal, annual and secular components (see *aberration, diurnal*; *aberration, annual*; *aberration, secular*).
- altitude*: the angular distance of a celestial body above or below the *horizon*, measured along the great circle passing through the body and the *zenith*. Altitude is 90° minus *zenith distance*.
- aphelion*: the point in a planetary *orbit* that is at the greatest distance from the Sun.
- apparent place*: the position on a *celestial sphere*, centered at the Earth, determined by removing from the directly observed position of a celestial body the effects that depend on the *topocentric* location of the observer; i.e., *refraction*, diurnal aberration (see *aberration, diurnal*) and geocentric (diurnal) *parallax*. Thus the position at which the object would actually be seen from the center of the Earth, displaced by planetary aberration (except the diurnal part – see *aberration, planetary* and *aberration, diurnal*) and referred to the *true equator and equinox*.
- apparent solar time*: the measure of time based on the diurnal motion of the true Sun. The rate of diurnal motion undergoes seasonal variation because of the *obliquity* of the *ecliptic* and because of the *eccentricity* of the Earth's *orbit*. Additional small variations result from irregularities in the rotation of the Earth on its axis.

- astrometric ephemeris*: an ephemeris of a solar system body in which the tabulated positions are essentially comparable to catalog *mean places* of stars at a *standard epoch*. An astrometric position is obtained by adding to the *geometric position*, computed from gravitational theory, the correction for *light-time*. Prior to 1984, the *E-terms* of annual aberration (see *aberration, annual* and *aberration, E-terms of*) were also added to the geometric position.
- astronomical coordinates*: the longitude and latitude of a point on the Earth relative to the *geoid*. These coordinates are influenced by local gravity anomalies (see *zenith*.)
- astronomical unit (AU)*: the radius of a circular orbit in which a body of negligible mass, and free of perturbations, would revolve around the Sun in $2\pi/k$ days, where k is the *Gaussian gravitational constant*. This is slightly less than the *semimajor axis* of the Earth's orbit.
- atomic second*: see *second, Système International*.
- augmentation*: the amount by which the apparent *semidiameter* of a celestial body, as observed from the surface of the Earth, is greater than the semidiameter that would be observed from the center of the Earth.
- azimuth*: the angular distance measured clockwise along the *horizon* from a specified reference point (usually north) to the intersection with the great circle drawn from the *zenith* through a body on the *celestial sphere*.
- barycenter*: the center of mass of a system of bodies; e.g., the center of mass of the solar system or the Earth–Moon system.
- barycentric dynamical time (TDB)*: the independent argument of ephemerides and equations of motion that are referred to the *barycenter* of the solar system. A family of time scales results from the transformation by various theories and metrics of relativistic theories of *terrestrial dynamical time (TDT)*. TDB differs from TDT only by periodic variations. In the terminology of the general theory of relativity, TDB may be considered to be a coordinate time. (See *dynamical time*.)
- catalog equinox*: the intersection of the *hour circle* of zero *right ascension* of a star catalog with the *celestial equator*. (See *dynamical equinox* and *equator*.)
- celestial ephemeris pole*: the reference pole for *nutation* and *polar motion*; the axis of figure for the mean surface of a model Earth in which the free motion has zero amplitude. This pole has no nearly-diurnal nutation with respect to a space-fixed or Earth-fixed coordinate system.

- celestial equator*: the projection onto the *celestial sphere* of the Earth's equator. (See *mean equator and equinox* and *true equator and equinox*.)
- celestial pole*: either of the two points projected onto the *celestial sphere* by the extension of the Earth's axis of rotation to infinity.
- celestial sphere*: an imaginary sphere of arbitrary radius upon which celestial bodies may be considered to be located. As circumstances require, the celestial sphere may be centered at the observer, at the Earth's center, or at any other location.
- conjunction*: the phenomenon in which two bodies have the same apparent celestial longitude (see *longitude, celestial*) or *right ascension* as viewed from a third body. Conjunctions are usually tabulated as *geocentric* phenomena, however. For Mercury and Venus, geocentric inferior conjunction occurs when the planet is between the Earth and Sun, and superior conjunction occurs when the Sun is between the planet and Earth.
- constellation*: a grouping of stars, usually with pictorial or mythical associations, that serves to identify an area of the *celestial sphere*. Also one of the precisely defined areas of the celestial sphere, associated with a grouping of stars, that the International Astronomical Union has designated as a constellation.
- coordinated universal time (UTC)*: the time scale available from broadcast time signals. UTC differs from TAI (see *international atomic time*) by an integral number of seconds; it is maintained within ± 0.90 seconds of UT1 (see *universal time*) by the introduction of one second steps (leap seconds).
- culmination*: passage of a celestial object across the observer's *meridian*; also called 'meridian passage'. More precisely, culmination is the passage through the point of greatest *altitude* in the diurnal path. Upper culmination (also called 'culmination above pole' for circumpolar stars and the Moon) or transit is the crossing closer to the observer's *zenith*. Lower culmination (also called 'culmination below pole' for circumpolar stars and the Moon) is the crossing farther from the zenith.
- day*: an interval of 86 400 SI seconds (see *second, Système International*), unless otherwise indicated.
- day numbers*: quantities that facilitate hand calculations of the reduction of *mean place* to *apparent place*. Besselian day numbers depend solely on the Earth's position and motion; second-order day numbers, used in high precision reductions, depend on the positions of both the Earth and the star.

- declination*: angular distance on the *celestial sphere* north or south of the *celestial equator*. It is measured along the *hour circle* passing through the celestial object. Declination is usually given in combination with *right ascension* or *hour angle*.
- defect of illumination*: the angular amount of the observed lunar or planetary disk that is not illuminated to an observer on the Earth.
- deflection of light*: the angle by which the apparent path of a photon is altered from a straight line by the gravitational field of the Sun. The path is deflected radially away from the Sun by up to 1".75 at the Sun's limb. Correction for this effect, which is independent of wavelength, is included in the reduction from *mean place* to *apparent place*.
- deflection of the vertical*: the angle between the astronomical vertical and the geodetic vertical (see *zenith*; *astronomical coordinates*; *geodetic coordinates*.)
- Delta T (ΔT)*: the difference between *dynamical time* and *universal time*; specifically the difference between *terrestrial dynamical time* (TDT) and UT1: $\Delta T = \text{TDT} - \text{UT1}$.
- direct motion*: for orbital motion in the solar system, motion that is counterclockwise in the orbit as seen from the north pole of the *ecliptic*; for an object observed on the celestial sphere, motion that is from west to east, resulting from the relative motion of the object and the Earth.
- DUT1*: the predicted value of the difference between UT1 and UTC, transmitted in code on broadcast time signals: $\text{DUT1} = \text{UT1} - \text{UTC}$. (See *universal time* and *coordinated universal time*.)
- dynamical equinox*: the ascending *node* of the Earth's mean orbit on the Earth's equator; i.e., the intersection of the *ecliptic* with the *celestial equator* at which the Sun's *declination* is changing from south to north. (See *catalog equinox* and *equinox*.)
- dynamical time*: the family of time scales introduced in 1984 to replace *ephemeris time* as the independent argument of dynamical theories and ephemerides. (See *barycentric dynamical time* and *terrestrial dynamical time*.)
- eccentric anomaly*: in undisturbed elliptic motion, the angle measured at the center of the ellipse from *pericenter* to the point on the circumscribing auxiliary circle from which a perpendicular to the major axis would intersect the orbiting body. (See *mean anomaly* and *true anomaly*.)
- eccentricity*: a parameter that specifies the shape of a conic section; one

- of the standard elements used to describe an elliptic *orbit* (see *elements, orbital*).
- eclipse*: the obscuration of a celestial body caused by its passage through the shadow cast by another body.
- eclipse, annular*: a solar *eclipse* (see *eclipse, solar*) in which the solar disk is never completely covered but is seen as an annulus or ring at maximum eclipse. An annular eclipse occurs when the apparent disk of the Moon is smaller than that of the Sun.
- eclipse, lunar*: an *eclipse* in which the Moon passes through the shadow cast by the Earth. The eclipse may be total (the Moon passing completely through the Earth's *umbra*), partial (the Moon passing partially through the Earth's *umbra* at maximum eclipse), or penumbral (the Moon passing only through the Earth's *penumbra*).
- eclipse, solar*: an *eclipse* in which the Earth passes through the shadow cast by the Moon. It may be total (observer in the Moon's *umbra*), partial (observer in the Moon's *penumbra*), or annular (see *eclipse, annular*).
- ecliptic*: the mean plane of the Earth's *orbit* around the Sun.
- elements, Besselian*: quantities tabulated for the calculation of accurate predictions of an *eclipse* or *occultation* for any point on or above the surface of the Earth.
- elements, orbital*: parameters that specify the position and motion of a body in *orbit* (see *osculating elements* and *mean elements*.)
- elongation, greatest*: the instants when the *geocentric* angular distances of Mercury and Venus are at a maximum from the Sun.
- elongation (planetary)*: the *geocentric* angle between a planet and the Sun, measured in the plane of the planet, Earth and Sun. Planetary elongations are measured from 0° to 180°, east or west of the Sun.
- elongation (satellite)*: the *geocentric* angle between a satellite and its primary, measured in the plane of the satellite, planet and Earth. Satellite elongations are measured from 0° east or west of the planet.
- ephemeris hour angle*: an *hour angle* referred to the *ephemeris meridian*.
- ephemeris longitude*: longitude (see *longitude, terrestrial*) measured eastward from the *ephemeris meridian*.
- ephemeris meridian*: a fictitious *meridian* that rotates independently of the Earth at the uniform rate implicitly defined by *terrestrial dynamical time* (TDT). The ephemeris meridian is $1.002\,738\Delta T$ east of the Greenwich meridian, where $\Delta T = \text{TDT} - \text{UT1}$.
- ephemeris time* (ET): the time scale used prior to 1984 as the independent variable in gravitational theories of the solar system. In 1984, ET was replaced by *dynamical time*.

- ephemeris transit*: the passage of a celestial body or point across the *ephemeris meridian*.
- equation of center*: in elliptic motion the *true anomaly* minus the *mean anomaly*. It is the difference between the actual angular position in the elliptic orbit and the position the body would have if its angular motion were uniform.
- equation of the equinoxes*: the *right ascension* of the mean equinox (see *mean equator and equinox*) referred to the *true equator and equinox*; apparent *sidereal time* minus mean *sidereal time*. (See *apparent place and mean place*.)
- equation of time*: the *hour angle* of the true Sun minus the hour angle of the *fictitious mean sun*; alternatively, *apparent solar time* minus *mean solar time*.
- equator*: the great circle on the surface of a body formed by the intersection of the surface with the plane passing through the center of the body perpendicular to the axis of rotation. (See *celestial equator*.)
- equinox*: either of the two points on the *celestial sphere* at which the *ecliptic* intersects the *celestial equator*; also the time at which the Sun passes through either of these intersection points: i.e., when the apparent longitude (see *apparent place and longitude, celestial*) of the Sun is 0° or 180° . (See *catalog equinox* and *dynamical equinox* for precise usage.)
- fictitious mean sun*: an imaginary body introduced to define *mean solar time*; essentially the name of a mathematical formula that defined mean solar time. This concept is no longer used in high precision work.
- flattening*: a parameter that specifies the degree by which a planet's figure differs from that of a sphere; the ratio $f = (a - b)/a$, where a is the equatorial radius and b is the polar radius.
- Gaussian gravitational constant* ($k = 0.017\ 202\ 098\ 95$): the constant defining the astronomical system of units of length (*astronomical unit*), mass (solar mass) and time (*day*), by means of Kepler's third law. The dimensions of k^2 are those of Newton's constant of gravitation: $L^3 M^{-1} T^{-2}$.
- geocentric*: with reference to, or pertaining to, the center of the Earth.
- geocentric coordinates*: the latitude and longitude of a point on the Earth's surface relative to the center of the Earth; also celestial coordinates given with respect to the center of the Earth. (See *zenith; latitude, terrestrial; longitude, terrestrial*.)

- geodetic coordinates*: the latitude and longitude of a point on the Earth's surface determined from the geodetic vertical (normal to the specified spheroid). (See *zenith*; *latitude*; *terrestrial*; *longitude*, *terrestrial*.)
- geoid*: an equipotential surface that coincides with mean sea level in the open ocean. On land it is the level surface that would be assumed by water in an imaginary network of frictionless channels connected to the ocean.
- geometric position*: the *geocentric* position of an object on the *celestial sphere* referred to the *true equator and equinox*, but without the displacement due to planetary aberration. (See *apparent place*; *mean place*; *aberration*, *planetary*.)
- Greenwich sidereal date* (GSD): the number of *sidereal days* elapsed at Greenwich since the beginning of the Greenwich sidereal day that was in progress at *Julian date* 0.0.
- Greenwich sidereal day number*: the integral part of the *Greenwich sidereal date*.
- Gregorian calendar*: the calendar introduced by Pope Gregory XIII in 1582 to replace the *Julian calendar*; the calendar now used as the civil calendar in most countries. Every year that is exactly divisible by four is a leap year, except for centurial years, which must be exactly divisible by 400 to be leap years. Thus 2000 is a leap year, but 1900 and 2100 are not leap years.
- heliocentric*: with reference to, or pertaining to, the center of the Sun.
- horizon*: a plane perpendicular to the line from an observer to the *zenith*. The great circle formed by the intersection of the *celestial sphere* with a plane perpendicular to the line from an observer to the *zenith* is called the astronomical horizon.
- horizontal parallax*: the difference between the *topocentric* and *geocentric* positions of an object, when the object is on the astronomical *horizon*.
- hour angle*: angular distance on the *celestial sphere* measured westward along the *celestial equator* from the *meridian* to the *hour circle* that passes through a celestial object.
- hour circle*: a great circle on the *celestial sphere* that passes through the *celestial poles* and is therefore perpendicular to the *celestial equator*.
- inclination*: the angle between two planes or their poles; usually the angle between an orbital plane and a reference plane; one of the standard orbital elements (see *elements*, *orbital*) that specifies the orientation of an *orbit*.

international atomic time (TAI): the continuous scale resulting from analyses by the Bureau International des Poids et Mesures of atomic time standards in many countries. The fundamental unit of TAI is the SI second (see *second*, *Système International*), and the epoch is 1958 January 1.

invariable plane: the plane through the center of mass of the solar system perpendicular to the angular momentum vector of the solar system.

irradiation: an optical effect of contrast that makes bright objects viewed against a dark background appear to be larger than they really are.

Julian calendar: the calendar introduced by Julius Caesar in 46 BC to replace the Roman calendar. In the Julian calendar a common year is defined to comprise 365 days, and every fourth year is a leap year comprising 366 days. The Julian calendar was superseded by the *Gregorian calendar*.

Julian date (JD): the interval of time in days and fraction of a day since 1 January 4713 BC, Greenwich noon, *Julian proleptic calendar*. In precise work the time scale, e.g., *dynamical time* or *universal time*, should be specified.

Julian date, modified (MJD): the Julian date minus 240 0000.5.

Julian day number (JD): the integral part of the *Julian date*.

Julian proleptic calendar: the calendric system employing the rules of the *Julian calendar*, but extended and applied to dates preceding the introduction of the Julian calendar.

Julian year: a period of 365.25 days. This period served as the basis for the *Julian calendar*.

Laplacian plane: for planets see *invariable plane*; for a system of satellites, the fixed plane relative to which the vector sum of the disturbing forces has no orthogonal component.

latitude, celestial: angular distance on the *celestial sphere* measured north or south of the *ecliptic* along the great circle passing through the poles of the ecliptic and the celestial object.

latitude, terrestrial: angular distance on the Earth measured north or south of the *equator* along the *meridian* of a geographic location.

librations: variations in the orientation of the Moon's surface with respect to an observer on the Earth. Physical librations are due to variations in the rate at which the Moon rotates on its axis. The much larger optical librations are due to variations in the rate of the Moon's orbital motion, the *obliquity* of the Moon's *equator* to its

orbital plane, and the diurnal changes of geometric perspective of an observer on the Earth's surface.

light-time: the interval of time required for light to travel from a celestial body to the Earth. During this interval the motion of the body in space causes an angular displacement of its *apparent place* from its *geometric position* (see *aberration*, *planetary*).

light year: the distance that light traverses in a vacuum during one year.

local sidereal time: the local *hour angle* of a *catalog equinox*.

longitude, celestial: angular distance on the *celestial sphere* measured eastward along the *ecliptic* from the *dynamical equinox* to the great circle passing through the poles of the ecliptic and the celestial object.

longitude, terrestrial: angular distance measured along the Earth's *equator* from the *Greenwich meridian* to the meridian of a geographic location.

lunar phases: cyclically recurring apparent forms of the Moon. New Moon, First Quarter, Full Moon and Last Quarter are defined as the times at which the excess of the apparent celestial longitude (see *longitude, celestial*) of the Moon over that of the Sun is 0° , 90° , 180° and 270° , respectively.

lunation: the period of time between two consecutive New Moons.

magnitude, stellar: a measure on a logarithmic scale of the brightness of a celestial object considered as a point source.

magnitude of a lunar eclipse: the fraction of the lunar diameter obscured by the shadow of the Earth at the greatest phase of a lunar eclipse (see *eclipse, lunar*), measured along the common diameter.

magnitude of a solar eclipse: the fraction of the solar diameter obscured by the Moon at the greatest phase of a solar eclipse (see *eclipse, solar*), measured along the common diameter.

mean anomaly: in undisturbed elliptic motion, the product of the *mean motion* of an orbiting body and the interval of time since the body passed *pericenter*. Thus the mean anomaly is the angle from pericenter of a hypothetical body moving with a constant angular speed that is equal to the mean motion. (See *true anomaly* and *eccentric anomaly*.)

mean distance: the *semimajor axis* of an elliptic orbit.

mean elements: elements of an adopted reference orbit (see *elements, orbital*) that approximates the actual, perturbed orbit. Mean elements may serve as the basis for calculating *perturbations*.

mean equator and equinox: the celestial reference system determined by ignoring small variations of short period in the motions of the *celestial equator*. Thus the mean equator and equinox are affected only by *precession*. Positions in star catalogs are normally referred to the mean catalog equator and equinox (see *catalog equinox*) of a *standard epoch*.

mean motion: in undisturbed elliptic motion, the constant angular speed required for a body to complete one revolution in an *orbit* of a specified *semimajor axis*.

mean place: the coordinates, referred to the *mean equator and equinox* of a *standard epoch*, of an object on the *celestial sphere* centered at the Sun. A mean place is determined by removing from the directly observed position the effects of *refraction*, *geocentric* and *stellar parallax*, and *stellar aberration* (see *aberration, stellar*), and by referring the coordinates to the mean equator and equinox of a standard epoch. In compiling star catalogs it has been the practice not to remove the secular part of stellar aberration (see *aberration, secular*). Prior to 1984, it was additionally the practice not to remove the elliptic part of annual aberration (see *aberration, annual* and *aberration, E-terms of*).

mean solar time: a measure of time based conceptually on the diurnal motion of the *fictitious mean sun*, under the assumption that the Earth's rate of rotation is constant.

meridian: a great circle passing through the *celestial poles* and through the *zenith* of any location on Earth. For planetary observations a meridian is half the great circle passing through the planet's poles and through any location on the planet.

moonrise, moonset: the times at which the apparent upper limb of the Moon is on the astronomical *horizon*; i.e., when the true *zenith distance*, referred to the center of the Earth, of the central point of the disk is $90^\circ 34' + s - \pi$, where s is the Moon's semidiameter, π is the *horizontal parallax*, and $34'$ is the adopted value of *horizontal refraction*.

nadir: the point on the *celestial sphere* diametrically opposite to the *zenith*.

node: either of the points on the *celestial sphere* at which the plane of an *orbit* intersects a reference plane. The position of a node is one of the standard orbital elements (see *elements, orbital*) used to specify the orientation of an orbit.

nutation: the short-period oscillations in the motion of the pole of rotation of a freely rotating body that is undergoing torque from

external gravitational forces. Nutation of the Earth's pole is discussed in terms of components in *obliquity* and longitude (see *longitude, celestial*).

obliquity: in general the angle between the equatorial and orbital planes of a body or, equivalently, between the rotational and orbital poles. For the Earth the obliquity of the *ecliptic* is the angle between the planes of the *equator* and the ecliptic.

occultation: the obscuration of one celestial body by another of greater apparent diameter; especially the passage of the Moon in front of a star or planet, or the disappearance of a satellite behind the disk of its primary. If the primary source of illumination of a reflecting body is cut off by the occultation, the phenomenon is also called an *eclipse*. The occultation of the Sun by the Moon is a solar eclipse (see *eclipse, solar*).

opposition: a configuration of the Sun, Earth and a planet in which the apparent *geocentric* longitude (see *longitude, celestial*) of the planet differs by 180° from the apparent geocentric longitude of the Sun.

orbit: the path in space followed by a celestial body.

osculating elements: a set of parameters (see *elements, orbital*) that specifies the instantaneous position and velocity of a celestial body in its perturbed *orbit*. Osculating elements describe the unperturbed (two-body) orbit that the body would follow if *perturbations* were to cease instantaneously.

parallax: the difference in apparent direction of an object as seen from two different locations; conversely the angle at the object that is subtended by the line joining two designated points. Geocentric (diurnal) parallax is the difference in direction between a *topocentric* observation and a hypothetical *geocentric* observation. Heliocentric or annual parallax is the difference between hypothetical geocentric and *heliocentric* observations; it is the angle subtended at the observed object by the *semimajor axis* of the Earth's *orbit*. (See also *horizontal parallax*.)

parsec: the distance at which one *astronomical unit* subtends an angle of one second of arc; equivalently the distance to an object having an annual *parallax* of one second of arc.

penumbra: the portion of a shadow in which light from an extended source is partially but not completely cut off by an intervening body; the area of partial shadow surrounding the *umbra*.

pericenter: the point in an *orbit* that is nearest to the center of force. (See *perigee* and *perihelion*.)

perigee: the point at which a body in *orbit* around the Earth most

closely approaches the Earth. Perigee is sometimes used with reference to the apparent orbit of the Sun around the Earth.

perihelion: the point at which a body in *orbit* around the Sun most closely approaches the Sun.

period: the interval of time required to complete one revolution in an *orbit* or one cycle of a periodic phenomenon, such as a cycle of *phases*.

perturbations: deviations between the actual *orbit* of a celestial body and an assumed reference orbit; also the forces that cause deviations between the actual and reference orbits. Perturbations, according to the first meaning, are usually calculated as quantities to be added to the coordinates of the reference orbit to obtain the precise coordinates.

phase: the ratio of the illuminated area of the apparent disk of a celestial body to the area of the entire apparent disk taken as a circle. For the Moon, phase designations (see *lunar phases*) are defined by specific configurations of the Sun, Earth and Moon. For eclipses phase designations (total, partial, penumbral, etc.) provide general descriptions of the phenomena (see *eclipse, solar; eclipse, annular; eclipse, lunar*.)

phase angle: the angle measured at the center of an illuminated body between the light source and the observer.

polar motion: the irregularly varying motion of the Earth's pole of rotation with respect to the Earth's crust. (See *celestial ephemeris pole*.)

precession: the uniformly progressing motion of the pole of rotation of a freely rotating body undergoing torque from external gravitational forces. In the case of the Earth, the component of precession caused by the Sun and Moon acting on the Earth's equatorial bulge is called *lunisolar precession*; the component caused by the action of the planets is called *planetary precession*. The sum of lunisolar and planetary precession is called *general precession*. (See *nutation*.)

proper motion: the projection onto the *celestial sphere* of the space motion of a star relative to the solar system; thus the transverse component of the space motion of a star with respect to the solar system. Proper motion is usually tabulated in star catalogs as changes in *right ascension* and *declination* per year or century.

quadrature: a configuration in which two celestial bodies have apparent longitudes (see *longitude, celestial*) that differ by 90° as viewed from a third body. Quadratures are usually tabulated with respect to the Sun as viewed from the center of the Earth.

- radial velocity*: the rate of change of the distance to an object.
- refraction, astronomical*: the change in direction of travel (bending) of a light ray as it passes obliquely through the atmosphere. As a result of refraction the observed *altitude* of a celestial object is greater than its geometric altitude. The amount of refraction depends on the altitude of the object and on atmospheric conditions.
- retrograde motion*: for orbital motion in the solar system, motion that is clockwise in the *orbit* as seen from the north pole of the *ecliptic*; for an object observed on the *celestial sphere*, motion that is from east to west, resulting from the relative motion of the object and the Earth. (See *direct motion*.)
- right ascension*: angular distance on the *celestial sphere* measured eastward along the *celestial equator* from the *equinox* to the *hour circle* passing through the celestial object. Right ascension is usually given in combination with *declination*.
- second, Système International (SI)*: the duration of 9 192 631 770 cycles of radiation corresponding to the transition between two hyperfine levels of the ground state of cesium 133.
- selenocentric*: with reference to, or pertaining to, the center of the Moon.
- semidiameter*: the angle at the observer subtended by the equatorial radius of the Sun, Moon or a planet.
- semimajor axis*: half the length of the major axis of an ellipse; a standard element used to describe an elliptical *orbit* (see *elements, orbital*).
- sidereal day*: the interval of time between two consecutive *transits* of the *catalog equinox*. (See *sidereal time*.)
- sidereal hour angle*: angular distance on the *celestial sphere* measured westward along the *celestial equator* from the *catalog equinox* to the *hour circle* passing through the celestial object. It is equal to 360° minus *right ascension* in degrees.
- sidereal time*: the measure of time defined by the apparent diurnal motion of the *catalog equinox*; hence a measure of the rotation of the Earth with respect to the stars rather than the Sun.
- solstice*: either of the two points on the *ecliptic* at which the apparent longitude (see *longitude, celestial*) of the Sun is 90° or 270° ; also the time at which the Sun is at either point.
- standard epoch*: a date and time that specifies the reference system to which celestial coordinates are referred. Prior to 1984 coordinates of star catalogs were commonly referred to the *mean equator and equinox* of the beginning of a Besselian year (see *year, Besselian*).

Beginning with 1984 the *Julian year* has been used, as denoted by the prefix J, e.g., J2000.0.

stationary point (of a planet): the position at which the rate of change of the apparent *right ascension* (see *apparent place*) of a planet is momentarily zero.

sunrise, sunset: the times at which the apparent upper limb of the Sun is on the astronomical *horizon*; i.e., when the true *zenith distance*, referred to the center of the Earth, of the central point of the disk is $90^{\circ} 50'$, based on adopted values of $34'$ for horizontal *refraction* and $16'$ for the Sun's *semidiameter*.

surface brightness (of a planet): the visual magnitude of an average square arc-second area of the illuminated portion of the apparent disk.

synodic period: for planets, the mean interval of time between successive *conjunctions* of a pair of planets, as observed from the Sun; for satellites, the mean interval between successive conjunctions of a satellite with the Sun, as observed from the satellite's primary.

terrestrial dynamical time (TDT): the independent argument for apparent *geocentric* ephemerides. At 1977 January $1^{\text{d}} 00^{\text{h}} 00^{\text{m}} 00^{\text{s}}$ TAI, the value of TDT was exactly 1977 January $1^{\text{d}} 000 3725$. The unit of TDT is 86 400 SI seconds at mean sea level. For practical purposes $\text{TDT} = \text{TAI} + 32^{\text{s}}.184$. (See *barycentric dynamical time*; *dynamical time*; *international atomic time*.)

terminator: the boundary between the illuminated and dark areas of the apparent disk of the Moon, a planet or a planetary satellite.

topocentric: with reference to, or pertaining to, a point on the surface of the Earth, usually with reference to a coordinate system.

transit: the passage of a celestial object across a *meridian*; also the passage of one celestial body in front of another of greater apparent diameter (e.g., the passage of Mercury or Venus across the Sun or Jupiter's satellites across its disk); however, the passage of the Moon in front of the larger apparent Sun is called an annular eclipse (see *eclipse, annular*). The passage of a body's shadow across another body is called a shadow transit; however, the passage of the Moon's shadow across the Earth is called a solar eclipse (see *eclipse, solar*).

true anomaly: the angle, measured at the focus nearest the *pericenter* of an elliptical *orbit*, between the pericenter and the radius vector from the focus to the orbiting body; one of the standard orbital elements (see *elements, orbital*). (See also *eccentric anomaly, mean anomaly*.)

true equator and equinox: the celestial coordinate system determined by the instantaneous positions of the *celestial equator* and *ecliptic*. The

- motion of this system is due to the progressive effect of *precession* and the short-term, periodic variations of *nutation*. (See *mean equator and equinox*.)
- twilight*: the interval of time preceding sunrise and following sunset (see *sunrise, sunset*) during which the sky is partially illuminated. Civil twilight comprises the interval when the *zenith distance*, referred to the center of the Earth, of the central point of the Sun's disk is between $90^{\circ} 50'$ and 96° , nautical twilight comprises the interval from 96° to 102° , astronomical twilight comprises the interval from 102° to 108° .
- umbra*: the portion of a shadow cone in which none of the light from an extended light source (ignoring *refraction*) can be observed.
- universal time (UT)*: a measure of time that conforms, within a close approximation, to the mean diurnal motion of the Sun and serves as the basis of all civil timekeeping. UT is formally defined by a mathematical formula as a function of *sidereal time*. Thus UT is determined from observations of the diurnal motions of the stars. The time scale determined directly from such observations is designated UT0; it is slightly dependent on the place of observation. When UT0 is corrected for the shift in longitude of the observing station caused by *polar motion*, the time scale UT1 is obtained.
- vernal equinox*: the ascending *node* of the *ecliptic* on the *celestial equator*; also the time at which the apparent longitude (see *apparent place and longitude, celestial*) of the Sun is 0° . (See *equinox*.)
- vertical*: apparent direction of gravity at the point of observation (normal to the plane of a free level surface).
- year*: a period of time based on the revolution of the Earth around the Sun. The calendar year (see *Gregorian calendar*) is an approximation to the tropical year (see *year, tropical*). The anomalistic year is the mean interval between successive passages of the Earth through *perihelion*. The sidereal year is the mean period of revolution with respect to the background stars. (See *Julian year* and *year, Besselian*.)
- year, Besselian*: the period of one complete revolution in *right ascension* of the *fictitious mean sun*, as defined by Newcomb. The beginning of a Besselian year, traditionally used as a *standard epoch*, is denoted by the suffix '.0'. Since 1984 standard epochs have been defined by the *Julian year* rather than the Besselian year. For distinction, the beginning of a Besselian year is now identified by the prefix B (e.g., B1950.0).

year, tropical: the period of one complete revolution of the mean longitude of the Sun with respect to the *dynamical equinox*. The tropical year is longer than the Besselian year (see *year, Besselian*) by $0^{\circ}.148T$, where T is centuries from B1900.0.

zenith: in general, the point directly overhead on the *celestial sphere*. The astronomical zenith is the extension to infinity of a plumb line. The geocentric zenith is defined by the line from the center of the Earth through the observer. The geodetic zenith is the normal to the geodetic ellipsoid or spheroid at the observer's location. (See *deflection of the vertical*.)

zenith distance: angular distance on the *celestial sphere* measured along the great circle from the *zenith* to the celestial object. Zenith distance is 90° minus *altitude*.

(From *The Astronomical Almanac for the Year 1987*, US Government Printing Office, with permission.)

Bibliography

- Astrophysical Quantities*, Allen, C. W., The Athlone Press, University of London, 1973.
- Astrophysical Formulae*, Lang, K. R., Springer-Verlag, 1974.
- Astronomy Handbook*, Roth, G. D., ed., Springer-Verlag, 1975.
- An Introduction to Astronomical Photometry*, Golay, M., D. Reidel Publishing Company, 1974.
- The Astronomical Almanac*, US Government Printing Office.
- Astronomy and Astrophysics*, Landolt-Börnstein, Springer-Verlag, 1981.
- Computational Spherical Astronomy*, Taff, L. G., John Wiley and Sons, 1981.

Chapter 3

Radio astronomy

	<i>page</i>
The major radio surveys	142
Microwave background	143
Selected discrete radio sources	145
The brightest radio sources visible in the northern hemisphere	148
Radio spectra	150
Radio flux calibrators	151
Radio propagation effects	153
<i>The Earth's atmosphere</i>	153
<i>Interplanetary medium</i>	153
<i>Interstellar medium (delay and Faraday rotation)</i>	154
<i>Interstellar medium (scintillation)</i>	155
Atmospheric opacity	155
Ionospheric electron density	156
Spectral features	157
Bibliography	157

The major radio surveys


Observatory	Survey	ν (MHz)	Sources	Limit (Jy) ^(a)
Cambridge	3C	159	471	8
	3CR	178	—	9
	4C	178	4843	2
	5C	408	276	0.025
	WKB	38	1069	14
	RN	178	87	0.25
	NB	81.5	558	1
Mills Cross	MSH	86	2270	7
Parkes	PKS	408, 1410, 2650	297	4
	PKS	408, 1410	247	0.5
	PKS	408, 1410	564	0.3
	PKS	408, 1410	628	0.4
	PKS	635, 1410, 2650	397	1.5
Owens Valley	CTA	906	106	—
	CTB, CTBR	960	110	—
	CTD	1421	—	1.15
National Radio Observatory	NRAO	750, 1400	726	(3C and 3CR)
	NRAO	750, 1400	458	0.5
Bologna	B1	408	629	1
	B2	408	3235	0.2
Ohio State	O	1415	128	2, 0.5
	O	1415	236	0.37
	O	1415	1199	0.3
	O	1415	2101	0.2
Vermillion River	VRO	610.5	239	0.8
	VRO	610.5	625	0.8
Dominion Radio Observatory	DA	1420	615	2
Dwingeloo- NRAO	DW	1417	188	2.3
Arecibo	AO	430	25	—

^(a) 1 jansky (Jy) = $10^{-26} \text{ W m}^{-2} \text{ Hz}^{-1}$.


(Adapted from Weinberg, S., *Gravitation and Cosmology*, John Wiley and Sons, 1972.)

Microwave background


Measurement of the surface brightness, B_ν , of the cosmic background radiation. The spectrum of a 2.75 K blackbody emitter is also shown. (Courtesy of T. Wilkinson, Princeton University.)



Free space microwave window. (From NASA SP-419, *The Search for Extraterrestrial Intelligence, SETI.*)



Microwave background (cont.)

Terrestrial microwave window. (From NASA SP-419, *op. cit.*)

Selected discrete radio sources

Source	α 1950	δ 1950	Spectral flux density (fu) ^a				Size (arcmin)	Log distance (pc)	Identification
			100	1000	10 000				
Cas B	00 ^h 23 ^m	+63 52'	250	56		7	3.5	Tycho SN I, 1572	
And A	00 40	+41 00	190	60		140	5.8	Andr. Galaxy, M31	
	00 54	-73	400	100					
Per A	02 22	+61 51	100	100		2	7.9	Multi. H II region OH em	
	03 16	+41 19	130	20				Seyfert Galaxy, NGC 1275	
For A	03 20	-37 22	400	120		Large		Pec. Galaxy, NGC 1316?	
Per 3C 123	04 34	+29 34	280	70		1			
	04 58	+46 26	120	150		60 + <i>h</i> (<i>l</i>)			
Pic A	05 18	-45 49	400	80		Large		Gal. Nebula SN II	
	05 21	-69	3000	700					
Tau A	05 32	+21 59	1700	955	560	5	3.3	Crab Nebula SN I 1054	
Ori Neb.	05 33	-05 24	40	340	400	10	2.7	Orion Nebula, M42	
Gem 3C 157	06 15	+22 38	400	180		30 + <i>h</i>	3.1	IC 443, SN II	
Mon	06 29	+04 54	400	250		70	3.0	Rosette Nebula	
Pup A	08 21	-42 58	600	150		40 + <i>h</i>	2.7		
	08 32	-45 37	500	200				Vela X (?)	
Hya A	09 16	-11 53	400	60	10	1	8.4	Pec. Galaxy	
Car	10 43	-59 30	500	800			3.1	Carina Nebula	
3C 273	12 27	+02 19	140	50		1		Quasar	
Vir A	12 28	+12 40	1800	263	40	5	7.1	Pec. Jet Galaxy, M87	
Cen A	13 22	-42 46	3000	2000		5 + <i>h</i>	6.8	Pec. Galaxy, NGC 5128	
Cen B	13 30	-60	600	80					

Selected discrete radio sources (*cont.*)

Source	α 1950	δ 1950	Spectral flux density (fu) ^(a)				Size (arcmin)	Log dist. (pc)	Identification
			100	1000	10 000	(MHz)			
Boo 3C 295	14 ^h 10 ^m	+52 26'	100	30		1		Distant galaxy	
Tr A	16 10	-60 47	800	80					
3C 338	16 27	+39 39	80	7		1		4 Galaxies, NGC 6161	
Her A	16 48	+05 04	700	70	8	3	8.6	Pec. Galaxy	
	17 11	-38 25	400	100					
2C 1473	17 16	-00 55	400	80	10	4		Galaxy	
	17 22	-34 14	400	400	500				
2C 1485	17 28	-21 20	80	20		1		Kepler SN I, 1604	
Sgr A	17 43	-28 56	4000	2000	200	70	2.9	Galactic center	
Trifid	17 58	-23 24	800	300			3.0	Galactic nebula, M20	
Lagoon	18 01	-24 22	70	150			3.1	Galactic nebula, M8	
	18 02	-21 30	200	150					
Omega	18 18	-16 10	200	800	500	10	3.2	Galactic nebula, M17	
	18 45	-02 06	500	300	250				
3C 392	18 54	+01 16	500	210		16		Shell source, SN	
3C 398	19 08	+09 01	40	70		3		SN II region, OH em	
3C 400	19 21	+14 20	400	400		60			
Cyg A	19 58	+40 35	13 800	2340	163	1.2	8.5	Radio galaxy	
Cyg X	20 21	+40 12	200	400		60			

Cyg X	20 34	+41 40	150	500	50	40	3.1	? γ Cyg complex
2C 1725	20 44	+50	400	150		100		SN II
Cyg Loop	20 49	+30	400	200		150	2.7	Loops SN II
America	20 52	+43 54	700	500		150	2.9	Galactic nebula
3C 446	22 23	-05 12	30	6				Quasar
Cas A	23 21	+58 32	19 500	3300	490	4	3.4	Galactic nebula SN II

^(a) Flux unit, $\text{fu} = 10^{-26} \text{W m}^{-2} \text{Hz}^{-1}$.

(After Allen, C. W., *Astrophysical Quantities*, The Athlone Press, 1973.)

The brightest radio sources visible in the northern hemisphere (based on observations at the 20 cm wavelength)

Name	α 1950	δ 1950	Intensity (fu)	Identification
3C 10	00 ^h 22 ^m 37 ^s	63° 51' 41"	44	Supernova remnant ^(a) - Tycho's supernova
3C 20	00 40 20	51 47 10	12	Galaxy
3C 33	01 06 13	13 03 28	13	Elliptical Galaxy
3C 48	01 34 50	32 54 20	16	Quasar
3C 58	02 01 52	64 35 17	34	Supernova remnant ^(a)
3C 84	03 16 30	41 19 52	14	Seyfert Galaxy
Fornax A	03 20 42	-37 25 00	115	Spiral Galaxy
NRAO 1560	04 00 00	51 08 00	26	
NRAO 1650	04 07 08	50 58 00	19	
3C 111	04 15 02	37 54 29	15	
3C 123	04 33 55	29 34 14	47	Galaxy
Pictor A	05 18 18	-45 49 39	66	D Galaxy ^(b)
3C 139.1	05 19 21	33 25 00	40	Emission nebula ^(a)
NRAO 2068	05 21 13	-36 30 19	19	N Galaxy ^(c)
3C 144	05 31 30	21 59 00	875	Supernova remnant ^(a) - Crab Nebula Taurus A
3C 145	05 32 51	-05 25 00	520	Emission nebula ^(a) Orion A NGC 1976
3C 147	05 38 44	49 49 42	23	Quasar
3C 147.1	05 39 11	-01 55 42	65	Emission nebula ^(a) - Orion B-NGC 2024
3C 153.1	06 06 53	20 30 40	29	Emission nebula ^(a)
3C 161	06 24 43	-05 51 14	19	
3C 196	08 09 59	48 22 07	14	Quasar
3C 218	09 15 41	-11 53 05	43	D Galaxy ^(b)
3C 270	12 16 50	06 06 09	18	Elliptical Galaxy
3C 273	12 26 33	02 19 42	46	Quasar
3C 274	12 28 18	12 40 02	198	Elliptical Galaxy - M87-Virgo A
3C 279	12 53 36	-5 31 08	11	Quasar
Centaurus A	13 22 32	-42 45 24	1330	Elliptical Galaxy NGC 5128
3C 286	13 28 50	30 45 58	15	Quasar
3C 295	14 09 33	52 26 13	23	D Galaxy ^(b)
3C 348	16 48 41	05 04 36	45	D Galaxy ^(b)
3C 353	17 17 56	-00 55 53	57	D Galaxy ^(b)
3C 358	17 27 41	-21 27 11	15	Supernova remnant ^(a) - Kepler's supernova
3C 380	18 28 13	48 42 41	14	Quasar
NRAO 5670	18 28 51	-02 06 00	12	
NRAO 5690	18 32 41	-07 22 00	90	
NRAO 5720	18 35 33	-06 50 18	30	
3C 387	18 38 35	-05 11 00	51	

The brightest radio sources visible in the northern hemisphere (cont.)

Name	α 1950	δ 1950	Intensity (fu)	Identification
NRAO 5790	18 ^h 43 ^m 30 ^s	-02 46' 39"	19	
3C 390.2	18 44 25	-02 33 00	80	
3C 390.3	18 45 53	79 42 47	12	N Galaxy
3C 391	18 46 49	-00 58 58	21	
NRAO 5840	18 50 52	01 08 18	15	
3C 392	18 53 38	01 15 10	171	Supernova remnant ^(a)
NRAO 5890	18 59 16	01 42 31	14	
3C 396	19 01 39	05 21 54	14	
3C 397	19 04 57	07 01 50	29	
NRAO 5980	19 07 55	08 59 09	47	
3C 398	19 08 43	08 59 49	33	
NRAO 6010	19 11 59	11 03 30	10	
NRAO 6020	19 13 19	10 57 00	35	
NRAO 6070	19 15 47	12 06 00	11	
3C 400	19 20 40	14 06 00	576	
NRAO 6107	19 32 20	-46 27 32	13	
3C 403.2	19 52 19	32 46 00	75	
3C 405	19 57 44	40 35 46	1495	D Galaxy ^(b) - Cygnus A
NRAO 6210	19 59 49	33 09 00	55	
3C 409	20 12 18	23 25 42	14	
3C 410	20 18 05	29 32 41	10	
NRAO 6365	20 37 14	42 09 07	20	Emission nebula ^(a)
NRAO 6435	21 04 25	-25 39 06	12	Elliptical Galaxy
NRAO 6500	21 11 06	52 13 00	46	
3C 433	21 21 31	24 51 18	12	D Galaxy ^(b)
3C 434.1	21 23 26	51 42 14	12	
NRAO 6620	21 27 41	50 35 00	37	
NRAO 6635	21 34 05	00 28 26	10	Quasar
3C 452	22 43 33	39 25 28	11	Elliptical Galaxy
3C 454.3	22 51 29	15 52 54	11	Quasar
3C 461	23 21 07	58 32 47	2477	Supernova remnant ^(a) - Cassiopeia A

fu = $10^{-26} \text{ W m}^{-2} \text{ Hz}^{-1}$.

^(a) Supernova remnants and emission nebulae lie within our own galaxy.


^(b) Galaxy refers to a Dumbell-shaped galaxy.

^(c) N Galaxy refers to a galaxy with a bright nucleus.

(Adapted from Verschuur, G. L., *The Invisible Universe*, Springer-Verlag, 1974.)

Radio spectra

Spectra of typical radio sources. (Adapted from Kraus, J. D., *Radio Astronomy*, McGraw-Hill Co., 1966.)



Radio flux calibrators*

(a) Spectral parameters of telescope calibrators

Source	Frequency interval	Spectral parameters $\log S[\text{Jy}] = a + b \cdot \log \nu [\text{MHz}] + c \cdot \log^2 \nu [\text{MHz}]$		
		a	b	c
3C 48	405 MHz ... 15 GHz	2.345	+0.071	0.138
3C 123	405 MHz ... 15 GHz	2.921	-0.002	-0.124
3C 147	405 MHz ... 15 GHz	1.766	+0.447	-0.184
3C 161	405 MHz ... 10.7 GHz	1.633	+0.498	-0.194
3C 218	405 MHz ... 10.7 GHz	4.497	-0.910	-
3C 227	405 MHz ... 15 GHz	3.460	-0.827	-
3C 249.1	405 MHz ... 15 GHz	1.230	+0.288	-0.176
3C 286	405 MHz ... 15 GHz	1.480	+0.292	-0.124
3C 295	405 MHz ... 15 GHz	1.485	+0.759	-0.255
3C 348	405 MHz ... 10.7 GHz	4.963	-1.052	-
3C 353	405 MHz ... 10.7 GHz	2.944	-0.034	-0.109
DR 21	7 GHz ... 31 GHz	1.81	-0.122	-
NGC 7027	10 GHz ... 31 GHz	1.32	-0.127	-
		± 0.030	± 0.001	± 0.001
		± 0.025	± 0.0001	± 0.001
		± 0.017	± 0.006	± 0.001
		± 0.016	± 0.008	± 0.001
		± 0.038	± 0.011	± 0.001
		± 0.055	± 0.016	± 0.003
		± 0.027	± 0.007	± 0.001
		± 0.018	± 0.006	± 0.001
		± 0.013	± 0.009	± 0.001
		± 0.045	± 0.014	± 0.001
		± 0.031	± 0.001	± 0.001
		± 0.05	± 0.010	± 0.001
		± 0.08	± 0.012	± 0.001

(b) Characteristics, position, and flux densities of telescope calibrators

Source	α 1950 [^h m s]	δ 1950 [[°] '"]	b^H []	$S_{14.00}$ [Jy]	$S_{1.665}$ [Jy]	$S_{2.700}$ [Jy]	$S_{5.000}$ [Jy]	$S_{8.000}$ [Jy]	$S_{10.700}$ [Jy]	$S_{15.000}$ [Jy]	$S_{22.235}$ [Jy]	Spec.	Ident.	Polar. (at 5 GHz) [" ₀]	Ang. size (at 1.4GHz) ["]
3C 48	01 34 49.8	+32 54 20	-29	15.9	13.9	9.20	5.24	3.31	2.46	1.72	1.11	C	QSS	5	<1
3C 123	04 33 55.2	+29 34 14	-12	48.7	42.4	28.5	16.5	10.6	7.94	5.63	3.71	C	GAL	2	20
3C 147	05 38 43.5	+49 49 42	+10	22.4	19.8	13.6	7.98	5.10	3.80	2.65	1.71	C	QSS	<1	<1
3C 161	06 24 43.1	-05 51 14	-8	19.0	16.8	11.4	6.62	4.18	3.09	2.14		C	GAL	5	<3
3C 218	09 15 41.5	-11 53 06	+25	43.1	36.8	23.7	13.5	8.81	6.77			S	GAL	1	core 25 halo 200
3C 227	09 45 07.8	+07 39 09	-42	7.21	6.25	4.19	2.52	1.71	1.34	1.02	0.73	S	GAL	7	180
3C 249.1	11 00 25.0	+77 15 11	+39	2.48	2.14	1.40	0.77	0.47	0.34	0.23		S	QSS		15
3C 274	12 28 17.7	+12 39 55	+74	214	184	122	71.9	48.1	37.5	28.1	20.0	S	GAL	1	halo 400 ^a
3C 286	13 28 49.7	+30 45 58	+81	14.8	13.6	10.5	7.30	5.38	4.40	3.44	2.55	C	QSS	11	<5
3C 295	14 09 33.5	+52 26 13	+61	22.3	19.2	12.2	6.36	3.65	2.53	1.61	0.92	C	GAL	0.1	4
3C 348	16 48 40.1	+05 04 28	+29	45.0	37.5	22.6	11.8	7.19	5.30			S	GAL	8	115 ^b
3C 353	17 17 54.6	-00 55 55		57.3	50.5	35.0	21.2	14.2	10.9			C	GAL	5	150
DR 21	20 37 14.2	+42 09 07	+1					21.6	20.8	20.0	19.0	Th	HII		20 ^c
NGC 7027 ^d	21 05 09.4	+42 02 03	-3	1.35	1.65	3.5	5.7		6.43	6.16	5.86	Th	PN	<1	10

^a Halo has steep spectral index, so for $\lambda \lesssim 6$ cm, more than 90% of the flux is in the core.

^b Angular distance between the two components.

^c Angular size at 2 cm, but consists of 5 smaller components.

^d Data up to 5 GHz are the direct measurements, not calculated from fit.

Systematic errors over the frequency range 0.4–15 GHz are less than 4%.

(Taken from Baars, J. W. M., Genzel, R., Pauliny-Toth, I. I. K., Witzel, A., *Astron. Astrophys.* **61**, 99 (1977).)

Radio propagation effects

The Earth's atmosphere (> 25 MHz)

$$S(z) = S_0 d^{-X(z)}$$

where

$S(z)$ = the flux measured at zenith distance z ,

S_0 = flux outside the atmosphere,

d = the attenuation at the zenith for airmass 1,

$X(z)$ = relative airmass in units of the airmass at the zenith.

For a plane parallel atmospheric model:

$$X(z) = \sec z = \frac{1}{\cos z}$$

Taking into account the Earth's and troposphere's curvature:

$$X(z) = \frac{1}{H} \int_R^{R+H} \frac{\rho(r)/\rho(R)}{\left[1 - \left(\frac{R}{r} \frac{n_0}{n}\right)^2 \sin^2 z\right]^{1/2}} dr$$

where

R = radius of Earth,

H = height of atmosphere,

$\rho(r)$ = gas density of the atmosphere at radius r ,

n = index of refraction at radius r .

n_0 = index of refraction at radius R .

Up to $X = 5.2$ the following equation gives $X(z)$, with an error less than 6.4×10^{-4} .

$$X(z) = -0.0045 + 1.00672 \sec z - 0.002234 \sec^2 z - 0.0006247 \sec^3 z.$$

(After K. Rohlfs, *Tools of Radio Astronomy*, Springer-Verlag, 1986.)

Interplanetary medium

The electron density as a function of distance from the Sun:

$$N_e = [1.55r^{-6} + 2.99r^{-16}] \times 10^{14} \quad (\text{m}^{-3}), \quad r < 4,$$

where r is the radial distance from the Sun in units of the solar radius,

$$N_e = 5 \times 10^{11} r^{-2} \quad (\text{m}^{-3}), \quad 4 < r < 20.$$

The scattering angle due to the interplanetary medium may be approximated by:

$$\theta_s \simeq 50 \left(\frac{\lambda}{p} \right)^2 \quad (\text{arcmin}),$$

where λ is in meters and p , the solar impact parameter, is in solar radii.

Interstellar medium (delay and Faraday rotation)

The smooth, ionized component of the interstellar medium of the Galaxy affects propagation by introducing delay and Faraday rotation. The time of arrival of a pulse of radiation is:

$$t_p = \int_0^L \frac{dz}{v_g},$$

where L is the propagation path, v_g is the group velocity;

$$\frac{dt_p}{dv} \simeq - \frac{e^2}{4\pi\epsilon_0 mc v^3} \int_0^L N_e dz.$$

The integral of N_e over the path length is called the *dispersion measure*,

$$D_m = \int_0^L N_e dz,$$

The magnetic field of the Galaxy causes Faraday rotation of the polarization plane of radiation from extragalactic radio sources.

$$\Delta\psi = \lambda^2 R_m,$$

where R_m is the *rotation measure* given by

$$R_m = 8.1 \times 10^5 \int N_e B_{\parallel} dz.$$

Here R_m is in radians per square meter, λ is in meters, B_{\parallel} is the longitudinal component of magnetic field in gauss (1 gauss = 10^{-4} tesla), N_e is in centimeters⁻³, dz is in parsecs (pc) (1 pc = 3.1×10^{16} m), and $\Delta\psi$ is the change in position angle of the electric field:

$$B_{\parallel} \simeq 2\mu G,$$

$$R_m \simeq -18 |\cot b| \cos(l - 94^\circ),$$

where l and b are the galactic longitude and latitude, respectively.

Interstellar medium (scintillation)

Approximate formulae for interstellar scintillation:

$$\begin{aligned} \theta_s &\simeq 7.5\lambda^{11.5} && (\text{arcsec}), && |b| \leq .6 \\ &\simeq 0.5|\sin b|^{-3.5}\lambda^{11.5} && (\text{arcsec}), && 0. < |b| < 3.5 \\ &\simeq 13|\sin b|^{-3.5}\lambda^{11.5} && (\text{milliarcsec}), && |b| \geq 3.5 \\ &\simeq \frac{15}{\sqrt{(|\sin b|)}} \lambda^2 && (\text{milliarcsec}), && |b| > 15. \end{aligned}$$

where b is the galactic latitude and λ is the wavelength in meters.


The accuracy of the approximations above decreases with decreasing $|b|$. In particular, the scattering angle at low latitudes, $|b| < 1$, can take on a large range of values.

(Adapted from Thompson, A. R., Moran, J. M. & Swenson, G. W., *Interferometry and Synthesis in Radio Astronomy*, Wiley-Interscience, 1986.)

(For propagation effects in the neutral lower atmosphere, see Thompson *et al.*, *op. cit.*)


Atmospheric opacity

Atmospheric zenith opacity. The absorption from narrow ozone lines has been omitted. (From Thompson, A. R. *et al.*, *op. cit.*)



Ionospheric electron density

Idealized electron density distribution in the Earth's ionosphere. The curves drawn indicate the densities to be expected at sunspot maximum in temperate latitudes. (From Evans & Hagfors, 1968.)



An incomplete list of astrophysically important spectral radio features†

Chemical name	Chemical formula	Transition	Frequency ^(a) (GHz)
Deuterium ^(b)	D		0.327
Hydrogen	H	$1^2S_{1/2}, F = 1-0$	1.420
Hydroxyl radical	OH	$^2\Pi_{3/2}, J = 3, 2, F = 1-2$	1.612
	OH	$^2\Pi_{3/2}, J = 3, 2, F = 1-1$	1.665
	OH	$^2\Pi_{3/2}, J = 3, 2, F = 2-2$	1.667
	OH	$^2\Pi_{3/2}, J = 3, 2, F = 2-1$	1.721
Formaldehyde	H ₂ CO	$1_{10}-1_{11}$, hyperfine structure	4.830
Water	H ₂ O	$6_{16}-5_{23}$, hyperfine structure	22.235
Ammonia	NH ₃	1, 1-1, 1, hyperfine structure	23.694
	NH ₃	2, 2-2, 2, hyperfine structure	23.723
	NH ₃	3, 3-3, 3, hyperfine structure	23.870
Silicon monoxide	SiO	$r = 2, J = 1-0$	42.820
	SiO	$r = 1, J = 1-0$	43.122
	SiO	$r = 1, J = 2-1$	86.243
Hydrogen cyanide	HCN	$J = 1-0$, hyperfine structure	88.631
Formyl ion	HCO ⁺	$J = 1-0$	89.189
Carbon monoxide	¹² C ¹⁸ O	$J = 1-0$	109.782
Carbon monoxide	¹³ C ¹⁶ O	$J = 1-0$	110.201
Carbon monoxide	¹² C ¹⁶ O	$J = 1-0$	115.271
Carbon monoxide	¹² C ¹⁶ O	$J = 2-1$	230.538
Carbon	C	$^3P_1-^3P_0$	492.162

^(a) λ (m) = 0.3/f (GHz).

^(b) Not yet found in radio spectra.

† Many hundreds of lines have been detected for more than 70 molecules.

(Adapted from Thompson, A. R., Moran, J. M. & Swenson, G. W., *Interferometry and Synthesis in Radio Astronomy*, Wiley-Interscience, 1986.)

Bibliography

A Handbook of Radio Sources, Pacholczyk, A. G., Pachart Publishing House, 1977.

Interferometry and Synthesis in Radio Astronomy, Thompson, A. R., Moran, J. M. & Swenson, G. W., Wiley-Interscience, 1986.

Radio Astronomy, 2nd edn., Kraus, J. O.

Tools of Radio Astronomy, Rohlfs, K., Springer Verlag, 1986.

Loves, F. J. et al., 1979, *Ap. J. (Suppl.)*, **41**, 451.

Chapter 4

Infrared astronomy

	<i>page</i>
Infrared sources	160
<i>Stars</i>	160
<i>Planetary nebula</i>	160
<i>H II regions</i>	160
<i>Molecular clouds</i>	161
<i>Galactic nucleus</i>	161
<i>Galactic nuclei</i>	161
<i>QSO</i>	161
Infrared background	162
Atmospheric transmission	164
Source temperatures	165
<i>Brightness temperature</i>	165
<i>Color temperature</i>	165
<i>Effective temperature</i>	165
Blackbody flux densities	165
Bibliography	165

Infrared sources

The brightest members of the seven presently known classes of infrared sources ($\text{Jy} = \text{Jansky} = \text{fu} = 10^{-26} \text{ W m}^{-2} \text{ Hz}^{-1}$).

1. STARS			
α 1950	δ 1950	20μ flux den. (Jy)	Name
1 ^h 04 ^m	12° 19'	0.83×10^3	CIT 3
2 17	-3 12	1.9	δ Cet
3 51	11 14	1.3	NML Tau
4 57	56 07	1.0	TY Cam
5 52	7 25	2.1	α Ori
7 21	-25 41	11	VY C Ma
9 45	11 39	1.1	R Leo
9 45	13 30	~25	IRC + 10216
10 13	30 49	0.83	RW LMi
10 43	-59 25	~50	η Car
12 01	-32 04	2.7	IRC - 30187
13 46	-28 07	1.9	W Hya
16 26	-26 19	1.0	α Sco
18 05	-22 16	2.1	VY Sgr
18 36	-6 51	2.5	EW Sct
18 45	-2 03	2.5	AB Aql
19 24	11 16	3.3	IRC + 10420
20 08	-6 25	1.4	IRC - 10529
20 20	37 22	1.1	BC Cyg
20 45	39 56	4.8	NML Cyg
21 42	58 33	0.69	μ Cep
2. PLANETARY NEBULA			
α 1950	δ 1950	20μ flux den. (Jy)	Name
21 ^h 00 ^m	36° 30'	2.5×10^3	'Egg Nebula'
21 05	42 02	0.6	NGC 7027
3. H II REGIONS			
α 1950	δ 1950	20μ flux den. (Jy)	Name
2 ^h 22 ^m	61° 52'	5.6×10^3	W3
5 33	-5 27	5.9	M42
5 39	-1 57	3.3	NGC 2024
17 44	-28 33	0.7	Sgr B2
18 01	-24 21	3.3	M8
18 06	-20 19	3.0	W31
18 11	-17 58	1.4	HFE50
18 16	-13 46	1.4	M16
18 18	-16 13	20	M17
18 43	-2 42	1.1	HFE 56
18 59	1 08	1.0	W48
19 08	9 02	1.7	HFE 58
19 11	10 48	1.2	?
19 20	13 59	1.2	HFE 59

Infrared sources (cont.)**3. H II REGIONS**

α 1950	δ 1950	20μ flux den. (Jy)	Name
19 ^h 21 ^m	14 24'	1.7×10^4	HFE 60
20 00	33 25	1.4	NGC 6857
20 26	37 13	1.2	Sharp 106
23 12	61 12	3.6	NGC 7358

4. MOLECULAR CLOUDS

α 1950	δ 1950	100μ flux den. (Jy)	Name
2 ^h 22 ^m	61 52'	$\sim 10^4$	W3 IRS 5
5 33	-5 27	1×10^5	KL Neb
6 05	-6 23	5×10^4	Mon R2
16 23	-24 17	3×10^4	ρ Oph DK.Cl.
20 37	42 12	$\sim 10^4$	W75 S OH

5. GALACTIC NUCLEUS

α 1950	δ 1950	20μ flux den. (Jy)	Name
17 ^h 43 ^m	-28 54'	2.6×10^3	Sgr A

6. GALACTIC NUCLEI

α 1950	δ 1950	20μ flux den. (Jy)	Name
00 ^h 45 ^m	-25 34'	30	NGC 253
2 40	00 20	60	NGC 1068
9 52	69 55	100	M 82
12 55	56 15	6	MK 231


7. QSO

α 1950	δ 1950	10μ flux den. (Jy)	Name
08 ^h 53 ^m	20 15'	0.04 \rightarrow 0.07	OJ 287
12 27	2 20	0.2 \rightarrow 0.5	3C 273
22 01	42 12	0.2 \rightarrow 0.7	BL Lac

(Adapted from Low, F. in *Symposium on Infrared and Submillimeter Astronomy*, G. G. Fazio, ed., D. Reidel Publishing Company, 1977.)


Infrared background

Infrared background fluxes. (Adapted from Rieke, G. H. *et al.*, *Science*, **231**, 807, 1986.)




Infrared background (cont.)

Predicted diffuse far-infrared radiation fluxes. The curve marked NR is an estimate of the contribution from rich clusters neglecting redshift and other relativistic effects. (Adapted from Stecker, F. W., Puget, J. L. & Fazio, G. G., *Ap. J. (Lett.)*, **214**, L51, 1977.)



Atmospheric transmission

Transmission of the atmosphere at infrared wavelengths at four altitudes. (Adapted from Fazio, G. G. in *Frontiers of Astrophysics*, E. H. Avrett, ed., Harvard University Press, Cambridge, 1976.)



Source temperatures

Brightness temperature

$$T_b(\nu) = \frac{c^2 B_\nu}{2\nu^2 k} \quad (\text{Rayleigh-Jeans approximation}).$$

T_b is the temperature of a blackbody which would have the same spectral radiance B_ν at frequency ν as the source.

Color temperature

$$\frac{B_{\lambda_1}}{B_{\lambda_2}} = \left(\frac{\lambda_2}{\lambda_1}\right)^5 \frac{e^{hc/\lambda_2 k T_c} - 1}{e^{hc/\lambda_1 k T_c} - 1}.$$

B_λ is the spectral radiance of source at wavelength λ .

Effective temperature

$$L = 4\pi R^2 \sigma T_{\text{eff}}^4,$$

where

L = source power,

R = radius of source,

σ = Stefan-Boltzmann constant.

Blackbody flux densities for stars

The blackbody spectral flux density for a star is given by,

$$F_\lambda = \frac{1.9 \times 10^{-11} \lambda^{-5} p^2 R^2}{e^{14388/\lambda T} - 1} \quad (\text{W cm}^{-2} \mu\text{m}^{-1})$$

where

R = radius of star (in units of the Sun's radius),

p = parallax (arcsec),

T = temperature (K),

λ = wavelength (μm).

(Johnson, H. M. & Wright, C. D., *Ap. J. Suppl.*, **53**, 643, 1983),

Bibliography

The Infrared Handbook, W. I. Wolfe & G. J. Zissis, eds., Office of Naval Research, Department of the Navy, Washington, DC, 1978.
Infrared Astronomy, G. Setti & G. Fazio, eds., D. Reidel Publishing Co., 1977.


Chapter 5

Ultraviolet astronomy


	<i>page</i>
UV stellar spectra	168
Stellar surface fluxes	170
Mass loss rates	170
Stellar EUV sources	171
Background fluxes	172
<i>Cosmic background</i>	172
<i>Geocoronal vacuum ultraviolet emission</i>	172
EUV plasma spectra	173
Spectral features	175
Important strong lines	176
Ultraviolet absorption cross-sections	177
Interstellar extinction	178
<i>Analytical fits to the UV extinction</i>	178
Interstellar EUV attenuation	179
Average interstellar hydrogen densities	180
Neutral hydrogen column density	181
Bibliography	182

UV stellar spectra


Spectral features in stars of different spectral types and luminosities.
(Courtesy of A. K. Dupree, Center for Astrophysics.)



IUE short wavelength spectra of dwarf stars. (Courtesy of A. K. Dupree, Center for Astrophysics.)




UV stellar spectra (cont.) IUE short wavelength spectra of giant and supergiant stars. (Courtesy of A. K. Dupree, Harvard-Smithsonian Center for Astrophysics)




Stellar surface fluxes

The ratio of stellar surface flux to the corresponding solar value for emission lines formed at various temperatures. (Courtesy of A. K. Dupree, Harvard-Smithsonian Center for Astrophysics.)



Mass loss rates

Characteristics of mass loss rates and winds (temperatures and terminal velocities) in stars of various luminosities. (Courtesy of A. K. Dupree, Harvard-Smithsonian Center for Astrophysics.)



Stellar EUV sources ($\leq 5\%$ of sky surveyed)


Source	α 1950	δ 1950	Distance (pc)	Flux ($\text{erg cm}^{-2} \text{s}^{-1}$)			Type
				500-780 Å	170-620 Å	50-150 Å	
HZ 43	$13^{\text{h}} 14^{\text{m}}$	+29 22'	65 ± 15	4×10^{-9}	1.1×10^{-9}	DA White Dwarf	
Feige 24	2 32	+2 12	$80 + 50 - 25$	3×10^{-9}	—	DA White Dwarf	
Proxima Centauri	14 26	-62 28	1.3	—	7×10^{-10}	Flare Star	
SS Cygni	21 40	+43 21	$30 - 50$	—	9×10^{-11}	Dwarf Nova	
HD 192273?	20 14	-69 45	—	7×10^{-10}	—	B Star	
VW Hyi?	4 40	-71	—	—	1.3×10^{-9}	U Gem	
Feige 4?	0 17	+13 36	70	6×10^{-11}	—	DB White Dwarf	

(Adapted from Paresce, F. in *Astrophysics from Spacelab*, P. L. Bernacca & R. Ruffini, D. Reidel Publishing Co., Dordrecht, 1980.)

Background fluxes

Cosmic background

Soft X-ray and EUV background fluxes. (Adapted from Stern, R. & Bowyer, S., *Ap. J.*, **230**, 755, 1979.)





Geocoronal vacuum ultraviolet emission

λ (Å)	Line	Photon radiance (ph cm ⁻² s ⁻¹ sr ⁻¹)	
		Day	Night
304	He II	8.0×10^5	8.0×10^5
584	He I	8.0×10^7	8.0×10^5
834	O II	8.0×10^7	2.0×10^6
1025	H I	1.6×10^8	1.6×10^8
1216	H I	8.0×10^8	8.0×10^8
1304-1356	O I	8.0×10^8	8.0×10^6
1356-1600	N ₂	2.4×10^7	2.4×10^7



$$1 \text{ Rayleigh} = \frac{1}{4\pi} \times 10^6 \text{ ph cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}.$$

EUV plasma spectra

EUV (100–1000 Å) spectrum of an optically thin plasma. Line fluxes are normalized assuming a line width of 1 Å. Logarithmic abundances: H = 12.00, He = 10.93, C = 8.52, N = 7.96, O = 8.82, Ne = 7.92, Mg = 7.42, Si = 7.52, S = 7.20, Fe = 7.60. Dashed curve: free-free continuum; dotted curve: two-photon continuum; solid curve: total continuum including free-bound radiation. (Adapted from Stern, R., Wang, E. & Bowyer, S., *Ap. J. Suppl.*, **37**, 195, 1978.)



EUV plasma spectra (cont.)



An incomplete list of astrophysically important ultraviolet spectral features


Identification	Wavelength (Å)†	Identification	Wavelength (Å)†
[Ne III]	3869	H I, D I	973
[O II]	3727	H I, D I	950
[Ne V]	3426	N II	917
Mg II	2798	H I, D I Ly edge	912
V II	2326	N II	916
C III	1909	O II	834
O III	1663	O III	834
He II	1640	O II	833
C IV	1549	O III	833
O IV	1402	S IV	816
Si IV	1397	O V	760
C II	1335	S IV	745
O I	1302	O II	719
Si II	1264	O II	718
S II	1260	O III	704
N V	1240	S IV	657
H I, D I	1216	O V	630
S III	1207	O III	600
N I	1201	He I	584
N I	1200	O IV	555
Si II	1190	O IV	554
C III	1176	O IV	553
O III	1175	O II	539
Fe II	1145	He I	537
N I	1134	He I	522
N II	1085	O III	508
N II	1084	O III	507
Ar I	1067	He I cont. edge	504
Si IV	1067	Ne VII	465
H ₂	1062	Mg IX	368
H ₂	1050	O III	306
H ₂	1049	He II	304
O VI	1038	Si XI	303
O II	1036	Fe XV	284
O VI	1032	He II	256
H I, D I	1026	Fe XXIV	255
N III	992	He II	243
N III	990	He II cont. edge	227
C III	977	O V	172

† $h\nu$ (eV) = 12 399/ λ (Å).

Brackets denote forbidden transitions.

Important strong lines

Wavelengths of important spectral lines of abundant elements and molecular hydrogen (H_2). Also indicated are the typical element abundances on a logarithmic scale where hydrogen is 12.00, and the temperatures of maximum fractional amount of each ion assuming collisional ionization equilibrium. Regions of continuous absorption by photoionization are indicated for hydrogen and helium. (From *FUSE Science Working Group Report*, NASA, 1983.)




Ultraviolet absorption cross-sections

λ (Å)	Line identification	Absorption cross-sections (10^{-18} cm ²)		
		O	O ₂	N ₂
1215	Ly α	0	0.01	0.000 03
1176	C III	0	1.30	0
1085	N II	0	2.00	0
1038	O VI	0	0.78	0.0007
1032	O VI	0	1.04	0.0007
1026	Ly β	0	1.58	0.0001
977	C III	0	4.0	0.7
897	Ly cont	2.9	13.0	~ 11.0
870	Ly cont	2.9	13.0	~ 10.0
800	Ly cont	2.9	29.0	~ 10.0
791	O IV	3.2	28.0	25.0
703	O III	7.0	25.0	26.0
630	O V	12.0	30.0	24.0
625	Mg X	12.0	25.0	24.0
584	He I	13.0	23.0	23.0
554	O IV	13.0	26.0	25.0
537	He I	12.0	21.0	25.0
504	He I	12.0	25.0	~ 22.0
499	Si XII	12.0	25.0	~ 22.0
465	Ne VII	11.0	23.0	23.0
368	Mg IX	9.0	18.0	16.0
335	Fe XVI	8.8	17.0	14.0
304	He II	9.0	17.0	12.0
284	Fe XV	7.7	15.0	9.8

(Adapted from Sullivan, J. O. & Holland, A. C., NASA CR-371, 1964.)

Interstellar extinction in the UV

The UV extinction $X(x) = A_\lambda/E_{B-V}$ against $x = 1/\lambda$ in microns. A_λ is the extinction in magnitudes; $E_{B-V} = A_B - A_V$ where A_B and A_V are the extinctions at the wavelengths of the B and V filters. The curve is from the analytical fit of the table below. (Figure courtesy of M. J. Seaton, University College London.)



Analytical fit to the UV extinction (see figure above)

Range of $1/\lambda(\mu) = x$	$A_\lambda/E(B-V) = X$
$2.70 \leq 1/\lambda \leq 3.65$	$1.56 + 1.048/\lambda + 1.01/\{(1/\lambda - 4.60)^2 + 0.280\}$
$3.65 \leq 1/\lambda \leq 7.14$	$2.29 + 0.848/\lambda + 1.01/\{(1/\lambda - 4.60)^2 + 0.280\}$
$7.14 \leq 1/\lambda \leq 10.0$	$16.17 - 3.20/\lambda + 0.2975/\lambda^2$

(From Seaton, M. J., *M.N.R.A.S.*, 187, 1979.)

Interstellar EUV attenuation


Distances (parsecs) corresponding to unit optical depth (1/e attenuation) as a function of neutral hydrogen density and wavelength

λ (Å)	$n_{\text{HI}} = 1$ (cm^{-3})	$n_{\text{HI}} = 0.1$ (cm^{-3})	$n_{\text{HI}} = 0.01$ (cm^{-3})	$n_{\text{HI}} = 0.001$ (cm^{-3})
912	0.05	0.5	5	50
500	0.2	2	20	200
200	1.5	15	150	1500
100	10	100	1000	10 000

(From FUSE Science Working Group Report, NASA, 1983.)

Distance at which the attenuation of EUV radiation reaches 90%. An ionized interstellar medium of normal composition is assumed.

(Adapted from Paresce, F. in *Astrophysics from Spacelab*, P. L. Bernacca & R. Ruffini, eds., D. Reidel Pub. Co., 1980.)



Average interstellar hydrogen densities within 100 pc

In the direction of	Distance (pc)	$n_{\text{H I}}$ (cm^{-3})
Sun	—	0.05
α Cen	1.34	0.06–0.30
ε Eri	3.3	0.06–0.20
ε Ind	3.4	~0.1
α CMi	3.5	0.09–0.13
β Gem	10.8	0.02–0.15
α Boo	11.1	0.02–0.15
α Aur	14	0.04–0.05
α Tau	21	0.02–0.15
α Leo	22	0.02
		0.01†
α Eri	28	0.07
α Gru	29	0.09–0.18
		0.18†
HR 1099	33	0.003–0.007
η UMa	42	0.005
G191–B2B	47	>0.03
σ Sgr	57	<0.17
HZ 43	62	<0.013
α Pav	63	<0.1
β Cen	81	0.13
β Lib	83	0.06–0.13
ζ Cen	83	<0.39
α Vir	87	0.037
Feige 24	90	0.02–0.05
λ Sco	100	<0.078


Mean hydrogen column density within 100 pc = $1.8 \times 10^{17} \text{ cm}^{-2} \text{ pc}^{-1}$.

† Multiple entries denote independent measurements.

(Adapted from Cash, W., Bowyer, S. & Lampton, M., *Astron. Astrophys.*, **80**, 67, 1979.)


Neutral hydrogen column density

Neutral hydrogen column density N_{HI} contours projected onto the plane of the galaxy ($b = 0^\circ$). The Sun is at the center of this plot, distances out to 100 pc are indicated, and the direction towards the galactic center ($l = 0^\circ$) is at the bottom. Line *A* is the contour of $N_{\text{HI}} \sim 5 \times 10^{17} \text{ cm}^{-2}$, corresponding to $\tau_{500 \text{ \AA}} = 1$, $\tau_{200 \text{ \AA}} \approx 0.1$, and $\tau_{100 \text{ \AA}} \approx 0.01$. Line *B* is the contour of $N_{\text{HI}} = 25 \times 10^{17} \text{ cm}^{-2}$, corresponding to $\tau_{500 \text{ \AA}} = 5$, $\tau_{200 \text{ \AA}} \approx 0.5$, and $\tau_{100 \text{ \AA}} \approx 0.05$. Line *C* is the contour of $N_{\text{HI}} \sim 50 \times 10^{17} \text{ cm}^{-2}$, corresponding to $\tau_{500 \text{ \AA}} = 10$, $\tau_{200 \text{ \AA}} \approx 1$, and $\tau_{100 \text{ \AA}} \approx 0.1$. All open circles are white dwarfs. Small circles represent stars with $N_{\text{HI}} \leq 5 \times 10^{17} \text{ cm}^{-2}$, medium circles represent stars with $5 \times 10^{17} < N_{\text{HI}} < 25 \times 10^{17} \text{ cm}^{-2}$, the large circles represent stars with $25 \times 10^{17} < N_{\text{HI}} < 50 \times 10^{17} \text{ cm}^{-2}$, and the crosses represent stars with $N_{\text{HI}} > 50 \times 10^{17} \text{ cm}^{-2}$. Stars with measured hydrogen column densities but which are located within 10 pc projected distance are not plotted. (From *FUSE Science Working Group Report*, NASA, 1983.)



Neutral hydrogen column density (*cont.*)

Same as previous diagram but projected onto a plane intercepting the Galactic plane at Galactic longitudes $l = 0^\circ$ and 180° , and passing through the North Galactic Pole (*top*) and South Galactic Pole (*bottom*). The symbols have the same meanings as before, and those stars with projected distances of less than 10 pc are generally not plotted. *W* symbols designate white dwarfs not yet observed. Many of these are within the cavity of low HI absorption and will be observable below 900 \AA . (From *FUSE Science Working Group Report*, NASA, 1983.)



Bibliography

The Universe at Ultraviolet Wavelengths, NASA Conference Publication 2171, 1980.

Proceedings of the Fourth European IUE Conference, ESA SP-218, 1984.


Chapter 6

X-ray astronomy

	<i>page</i>
Uhuru X-ray map	184
The HEAO A1 all sky map	185
Galactic sources: binaries and stars	186
Soft X-ray H–R diagrams	188
Binary X-ray pulsars	189
X-ray emitting supernova remnants	191
Crab Nebula	192
X-ray emission from normal galaxies	193
X-ray emission from clusters of galaxies	193
X-ray emission from active galaxies	193
3C 273	194
Quasar X-ray luminosity	195
X-ray source nomogram	196
Diffuse X-ray background	197
Absorption of X-rays	198
<i>Photoabsorption cross-sections (UV–X-ray)</i>	198
<i>Effective cross-section of the interstellar medium</i>	199
<i>Total photo-ionization cross-section</i>	200
<i>Photoelectric absorption cross-section</i>	201
<i>Attenuation of photons in the atmosphere</i>	203
<i>Photoelectric absorption in the interstellar medium</i>	204
Spectral features	204
Model X-ray spectral distributions	205
Emission lines for low density plasma	206
Plasma emission – power in lines	208
Bibliography	208


Uhuru X-ray map

X-ray sources from the *Fourth Uhuru Catalog* displayed in galactic coordinates. The size of the symbol representing a source is proportional to the logarithm of the peak source intensity. The 339 X-ray sources observed with the UHURU (SAS-A) X-ray observatory are displayed. (Adapted from Forman, W. *et al.*, *Ap. J. Suppl.*, **38**, 357, 1978.)



The HEAO A1 all sky map

Discrete X-ray sources observed by the NRL Sky Survey Experiment on HEAO-1 are displayed in galactic coordinates. The size of the dot representing a source is proportional to the logarithm of the intensity averaged over the time interval when the source was in the field of view. 700 sources are shown. (Courtesy of K. Wood, Naval Research Laboratory.)



Galactic sources: binaries and stars



Source	α 1950 δ 1950	Flux density ^(a) (2-11 keV)		$L(X)$ max (2-11 keV) (erg s^{-1})	V magnitude	$\frac{L(X)}{L(O)}$	Spectral type	Periods	Distance	Remarks
		max (μJy)	min (μJy)							
SMC X-1	01 ^h 15 ^m 45 ^s .6 -73° 42' 22"	57		6×10^{38}	13.3	1.2	B0 I	3.89 ^d 0.71 ^s	65 kpc	Sanduleak 160
β Per	03 04 54.3 40 45 52	9		2×10^{31}	2.2	5×10^{-5}	B8 V/K0 IV/ Am	2.9 ^d	32 pc	Algol
3U 0352+30	03 52 15.1 30 54 01	37		1.2×10^{34}	6.0-6.7	1×10^{-4}	O9.5 (III-V)e	581 ^d (?) 13.9 ^m	350 pc	X Per
A 0620-00	06 20 11.1 -00 19 11	~50 000 $\lesssim 5$		3×10^{38}	10.4 (18.3)	85			1.5-2.5 kpc	X-ray Nova V616 Mon
Vela X-1	09 00 13.2 -40 21 25	280 <28		1.4×10^{36}	6.9	3×10^{-3}	B0.5 Ib	8.97 ^d 283 ^s	1.4 kpc	HD 77581
Cen X-3	11 19 01.9 -60 20 57	224 <21		4×10^{37}	13.4	0.05	O6.5 II-III	2.09 ^d 4.8 ^s	8 kpc	Krzeminski's star
Sco X-1	16 17 04.5 -15 31 15	19 000 6900		2×10^{37}	12.2-13.3	600		0.787 ^d	0.7 kpc	V818 Sco
Her X-1	16 56 01.7 35 25 05	160 11		1.0×10^{37}	13.2	10	A09-F0	34.8 ^d 1.7 ^d 1.2 ^s	5 kpc	HZ Her
3U 1700-37	17 00 32.7 -37 46 29	110 <11		3×10^{36}	6.6	5×10^{-4}	O6.5f	3.4 ^d	1.7 kpc	
4U 1813+50	18 14 58.6 49 50 55	7 2.5		8×10^{33}	13.1-12.3	0.6		0.129 ^d	300 kpc	AM Her

Cyg X-1	19 56 28.9	1320	2×10^{37}	8.9	2×10^{-2}	O9.7 Iab	5.6 ^d	2.5 kpc	Blackhole candidate
	35 03 55.0	260							HDE 226868
Cyg X-3	20 30 37.6	430	1.2×10^{38}				16.8 ^d (?)	10.5 kpc	IR, radio
	40 47 13	90					4.8 ^h		
SS Cygni	21 40 42.6	20	$1-5 \times 10^{32}$	12.1-8.1	1.3		6 ^h 38 ^m	100 pc	Dwarf nova (U Gem)
	43 21 51	5							
Cyg X-2	21 42 36.9	740		15.5	450		11.2 ^d		Sub-dwarf
	38 05 28	220							
X-ray intensity (>0.2 keV) ($\text{erg cm}^{-2} \text{s}^{-1}$)									
UV Cet	1 36 24 -18 13 00	7×10^{-11}		6.8-12.9	2×10^{-3}	dM5.5 e			Flare star
α Aur	5 12 59.5 45 56 58	3×10^{-10}		0.1	1×10^{-5}	G8 III + F	104 ^d	14 pc	Capella
α CMa	6 42 54 -16 39 00	9.5×10^{-12}		-1.5		A IV + D0	44.98 ^y	2.7 pc	Sirius
HZ 43	13 14 00 29 22 00	9×10^{-10}		12.9	6				White dwarf

(^a) Flux density = (integrated 2-11 keV flux)/9 keV; $1 \mu\text{Jy} = 0.242 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1} = 1.51 \times 10^{-3} \text{ keV cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1}$. (Adapted from Bradt, H. V., Doxsey, R. E. & Jernigan, J. G., *COSPAR Symposium on X-ray Astronomy*, Innsbruck, Austria, May 31, 1978).

Soft X-ray H-R diagrams

Soft X-ray (0.15–3 keV) H-R diagram for (a) giant and (b) main sequence stars from the Einstein Observatory stellar survey (K. Topka, thesis, Harvard University, 1980). Thick bars and arrows denote X-ray luminosity medians or their upper limits. The upper limit or lowest detection and the highest detection are also shown.



Binary X-ray pulsars


Source	Optical counter-part	Pulse period (s)	$-\dot{P}/P$ (yr^{-1})	Orbital period (d)	$a_x \sin i$ (lt-s)	$f(M)$ (M_\odot)	Orbital eccentricity	$K_c^{(a)}$ (km s^{-1})	Eclipse half-angle ($^\circ$)	$L_x^{(b)}$ (erg s^{-1})
A 0538-66	Identified	0.069	—	16.66	—	—	>0.4	—	—	1.2×10^{34}
SMC X-1	Sk 160	0.714	7.1×10^{-4}	3.892	53.46(3)	10.8	<0.0007	19(2)	26.5 29	6×10^{38}
Her X-1	HZ Her	1.24	2.9×10^{-6}	1.700	13.1831(3)	0.85	<0.0003	20.2(3.5)	24.4-24.7	7×10^{36}
1E 2259+586	Identified	3.49	—	<0.08	—	—	—	—	—	3×10^{34}
4U 0115+63	Identified	3.61	3.2×10^{-5}	24.31	140.13(10)	5.00	0.3402(2)	—	0	8×10^{36}
Cen X-3	V779 Cen	4.84	2.8×10^{-4}	2.087	39.792(5)	15.5	0.0008(1)	24(6)	35-40	8×10^{37}
4U 1626-67	KZ TrA	7.68	1.9×10^{-4}	0.0288	<0.04	$<8 \times 10^{-5}$	—	280(76)	0	—
2S 1553-54	—	9.26	—	30.7(2.8)	165(30)	5.1(2.9)	<0.07	—	—	—
LMC X-4	Identified	13.5	$<1.2 \times 10^{-3}$	1.408	30(5)	15(8)	<0.2	37.9(2.4)	25.5-33	4×10^{38}
2S 1417-62	—	17.6	9×10^{-3} (?)	>15	>25	—	—	—	—	—
OA0 1653-40	—	38.2	5.4×10^{-3}	—	—	—	—	—	—	—
A 0535+26	HDE	—	—	—	—	—	—	—	—	—
245770	—	104	3.5×10^{-2}	111(?)	—	—	—	≤ 20	—	2×10^{37}
GX 1+4	V2116 Oph	122	2.1×10^{-2}	>15	>60	—	—	—	—	6×10^{37}
GX 304-1	Identified	272	—	132(?)	—	—	—	—	—	3×10^{36}
4U 0900-40	HD 77581	283	1.7×10^{-4}	8.965	113.0(8)	19.3	0.092(5)	21.8(1.2)	31 37	6×10^{36}
4U 1145-61	HEN 715	292	$<10^{-4}$	187(?)	>100	—	—	—	—	6×10^{36}
1E 1145.1-	—	—	—	—	—	—	—	—	—	—
6141	Identified	297	$<10^{-4}$	>12	>50	—	—	—	—	3×10^{36}
A 1118-61	HEN	—	—	—	—	—	—	—	—	—
3-640	—	405	—	—	—	—	—	—	—	5×10^{36}
4U 1538-52	QV Nor	529	$<2 \times 10^{-3}$	3.730	55.2(3.7)	13	—	33(7)	25 33	4×10^{36}
GX 301-2	WRA 977	696	7×10^{-3}	41.4	367(3)	31	0.47(1)	—	0	1×10^{37}
4U 0352+30	X Per	835	1.8×10^{-4}	580(?)	—	—	—	<20	—	1×10^{34}

^(a) K_c = semi-amplitude of the optical Doppler velocity curve.

^(b) L_x = X-ray luminosity (2-11 keV) from Bradt, H. & McClintock. *Ann. Rev. Astron. Astrophys.*, **21**, 13, 1983.

(Adapted from Joss, P. C. and Rappaport, S. A., *Ann. Rev. Astron. Astrophys.*, **22**, 537, 1984.)

Sky map, in galactic coordinates, of 21 binary X-ray pulsars (●) and 27 X-ray burst sources (○). (Reproduced, with permission, from the *Annual Review of Astronomy and Astrophysics*, Vol. 22, (1984, by Annual Reviews Inc. Courtesy of Paul C. Joss, MIT.)



X-ray emitting supernova remnants

Source	α 1950 δ 1950	Distance (kpc)	Diameter (pc)	L_x (0.2-2 keV) (10^{35} erg s $^{-1}$)	Age (yr)	1 GHz flux density (fu)	Angular size (arcmin)
Crab Nebula	05 ^h 31 ^m 21° 59'	2	3	160	900	1000	3.0 × 4.2
Cas A	23 21 58 33	3	3.5	30	300	3000	4.0 × 3.8
Cygnus Loop	20 49 30 30	0.8	40	8	20 000	180	200 × 160
Vela	8 32 -45 00	0.5	44	4	13 000	1800	220 × 180
IC 443	06 14 22 30	1.5	20	0.4	6000	160	40 diameter
Tycho's SNR	00 22 63 52	6	13	40	400	58	6.0 × 7.0
SN 1006	15 00 -41 45	1.2	10	0.2	970	25	30 × 22
GKP SNR	19 31 31 10	1.2	70	0.8	300 000	50	240 × 200
Lupus Loop	15 09 -40	0.5	40	0.1	20 000	340	270 diameter
RCW 86	14 39 -62 15	2.5	28	2	1800	33	40 diameter
RCW 103	16 14 -50 56	3.3	9	1	1000	22	7.0 × 7.9
PKS 1209-52	12 06 -52 10	2	40	0.7	20 000	49	86 × 75
Pup A	08 20 -42 50	1.2	17	6	10 000	145	55 diameter
65.6+6.5	19 31 31 10	1.2	60-80	0.7	<2 × 10 ⁴		180 × 240

Crab Nebula

The observed electromagnetic spectrum of the Crab Nebula and the Crab Pulsar. Dashed lines show corrections made for absorption and scattering of interstellar material. (Adapted from Seward, F. O., *Journal of the British Interplanetary Society*, 31, 83, 1978.)

X-ray spectrum (0.1–100 keV):


$$\frac{dN}{dE} = 10 E^{-2.05} \exp(-\sigma n_H) \text{ photons cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1},$$

$$n_H = 3 \times 10^{21} \text{ cm}^{-2}.$$

X-ray luminosity:

$$L_x = 4.9 \times 10^{37} \text{ (0.1–100 keV)}$$

Distance:	2200 pc
Diameter:	3 pc (5 arcmin)
m_v :	8.6
A_v (absorption):	1.5
Coordinates:	$\alpha = 05^{\text{h}}31^{\text{m}}$, $\delta = +22^{\circ}$
Total radiated energy:	$1.8 \times 10^{38} \text{ erg s}^{-1}$
Age:	900 years (AD 1054)
Pulsar period, P (1969):	0.0331 s
Rate of period increase, \dot{P} :	$422.69 \times 10^{-15} \text{ ss}^{-1}$.



X-ray emission from normal galaxies

Galaxy	Distance (Mpc)	Radius (kpc)	Total mass ($10^{10} M_{\odot}$)	X-ray luminosity ^(a) ($2-10$ keV) (erg s^{-1})	L_x/M ($\text{erg s}^{-1} M_{\odot}^{-1}$)
LMC	0.052	3.5	1.0	4×10^{38}	4×10^{28}
SMC	0.063	1.5	0.15	1×10^{38}	6×10^{28}
M31	0.67	8	30	3×10^{39}	1×10^{28}
Our Galaxy		12	14	5×10^{39}	4×10^{28}

X-ray emission from clusters of galaxies

Cluster	Distance (Mpc)	Radius (Mpc)	Optical luminosity ($10^{11} L_{\odot}$)	Virial mass Luminosity (M_{\odot}/L_{\odot})	X-ray luminosity ^(a) ($2-10$ keV) (erg s^{-1})
Perseus	97	0.44	10.0	461	1×10^{45}
Coma	113	2.63	49.0	1020	5×10^{44}
Centaurus	250				4×10^{43}
Virgo	19	1.07	12.0	668	1.5×10^{43}


X-ray emission from active galaxies

Object	Class of galaxy	X-ray source	$2-10$ keV flux density ($10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$)	$D^{(b)}$ (Mpc)	X-ray lumi- nosity ^(a) ($2-10$ keV erg s^{-1})
NGC 4151	Seyfert	4U 1206+39	10	20	5×10^{42}
Cen A	Giant radio source	4U 1322-42	20	9	2×10^{42}
3C 273	Quasar	4U 1226+02	6	950	7×10^{45}
M87	Radio source in cluster	4U 1228+12	52	25	4×10^{43}
NGC 1275 (Perseus A)	Peculiar Seyfert in cluster	4U 0316+41	29	110	4×10^{44}
3C 120	Radio/optical variable, Seyfert	4U 0432+05	7	190	3×10^{44}
3C 390.3	N galaxy, compact radio source	4U 1847+78	7	336	1×10^{45}
M82	Emission line galaxy	4U 0954+70	6	4	1×10^{41}
Mkn 501	BL Lac object	4U 1651+39	5	180	2×10^{44}

^(a) $L_x = (\text{X-ray flux}) \cdot 4\pi D^2$.


^(b) $D = \frac{zc}{H_0}$ ($H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

The observed electromagnetic spectrum of the quasar 3C273 (3C273 is variable). (From Worrall, D. M. *et al.*, *Ap. J.*, **232**, 683, 1979.)




Quasar X-ray luminosity

Quasar X-ray luminosity (0.5–4.5 keV) versus redshift. (Courtesy H. Tananbaum, Harvard/Smithsonian Center for Astrophysics.)




X-ray source nomogram

Nomogram to compute $\log(f_x/f_v)$ for X-ray sources, where f_x is the X-ray flux density in $\text{erg cm}^{-2} \text{s}^{-1}$ in the 0.3–3.5 keV band and f_v is the flux density in the V band. For each object class indicated (stars: B–F, G, K, M; normal galaxies; active galactic nuclei; BL Lac objects) a continuous horizontal line indicates the range of $\log(f_x/f_v)$ comprising 70% of the known sources in the class and a dashed line indicates the range comprising the highest and lowest 15% of the sources. For example, for an X-ray source with a flux density of $2 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ and a V magnitude of 20, $\log(f_x/f_v)$ is ~ 0.7 and the source is most likely a BL Lac object. (From Maccacaro, T. *et al.*, *Ap. J.*, **326**, 680, 1988).



Diffuse X-ray background


Energy spectrum of diffuse X-ray background. (Courtesy of D. Schwartz, Harvard/Smithsonian Center for Astrophysics.)



Absorption of X-rays


Photoabsorption cross-sections (UV-X-ray) of abundant elements

Photoabsorption cross-sections of the abundant elements in the interstellar medium as a function of wavelength. (Adapted from Cruddace, R., Paresce, F., Bowyer, S. & Lampton, M., *Ap. J.*, **187**, 497, 1974.)




Effective cross-section of the interstellar medium

Effective cross-section (cross-section per hydrogen atom or proton) of the interstellar medium: — gaseous component with normal composition and temperature; - - hydrogen, molecular form; HII region about a B star; HII region about an O star; dust. (Adapted from Cruddace, R., Paresce, F., Bowyer, S. & Lampton, M., *Ap. J.*, 187, 497, 1974.)




Total photoionization cross-section

Total photo-ionization cross-section per hydrogen atom [$\sigma(E/1 \text{ keV})^3$] in units 10^{-22} cm^2 as a function of incident photon energy for a gas having a cosmic elemental abundance. The elements responsible for the discontinuities due to their K edges are shown. (Adapted from Brown, R. L. & Gould, R. J., *Physical Review*, **01**, 2252, No. 8, 1970.)



Photoelectric absorption cross-section

Net photoelectric absorption cross-section per hydrogen atom as a function of energy, scaled by $(E/1 \text{ keV})^3$. The solid line is for relative abundances given in the table of elemental abundances below, with all elements in the gas phase and in neutral atomic form. The dotted line shows the effect of condensing the fraction of each element indicated in the table into $0.3 \mu\text{m}$ grains. The contributions of hydrogen and hydrogen plus helium to the total cross-section are also shown. (From Morrison, R. & McCammon, D., *Ap. J.*, **270**, 119, 1983. Diagram courtesy of D. McCammon.)



Elemental abundances

Element	Abundance ^(a)	Fraction in grains ^(b)
H	12.00	0
He	11.00	0
C	8.65	1
N	7.96	1
O	8.87	0.25
Ne	8.14	0
Na	6.32	1
Mg	7.60	1
Al	6.49	1
Si	7.57	1
S	7.28	1
Cl	5.28	1
Ar	6.58	0
Ca	6.35	1
Cr	5.69	1
Fe	7.52	1
Ni	6.26	1

^(a) \log_{10} abundance relative to hydrogen = 12.00. All values except helium are from Anders and Ebihara, 1982.

^(b) Fraction of atoms of each element assumed depleted from gas phase and condensed into grains of average thickness 2.1×10^{18} atoms cm^{-2} for case shown as dotted line in the diagram.

Coefficients of analytic fit to cross-section

Energy range (keV)	c_0	c_1	c_2
0.030–0.100 ^(a)	17.3	608.1	–2150
0.100–0.284	34.6	267.9	–476.1
0.284–0.400	78.1	18.8	4.3
0.400–0.532	71.4	66.8	–51.4
0.532–0.707	95.5	145.8	–61.1
0.707–0.867	308.9	–380.6	294.0
0.867–1.303	120.6	169.3	–47.7
1.303–1.840	141.3	146.8	–31.5
1.840–2.471	202.7	104.7	–17.0
2.471–3.210	342.7	18.7	0.0
3.210–4.038	352.2	18.7	0.0
4.038–7.111	433.9	–2.4	0.75
7.111–8.331	629.0	30.9	0.0
8.331–10.000	701.2	25.2	0.0


Note: cross-section per hydrogen atom = $(c_0 + c_1 E + c_2 E^2) E^{-3} \times 10^{-24} \text{ cm}^2$ (E in keV).

^(a) Break introduced to allow adequate fit with quadratic: no absorption edge at 0.1 keV.

(From Morrison, R. & McCammon, D., *op. cit.*)


Attenuation of photons in the atmosphere

Attenuation of photons in the 1972 COSPAR International Reference Atmosphere with $1/e$ absorption length plotted as a function of energy and altitude or atmospheric depth.



Photoelectric absorption in the interstellar medium

Photoelectric absorption in the interstellar medium. The vertical axis gives the column density in units of hydrogen atoms cm^{-2} at which the absorption is $1/e$ at the photon energy E_a . (For a hydrogen atom number density of $1/\text{cm}^3$, 1 kpc is equivalent to a column density of 3.1×10^{21} hydrogen atoms cm^{-2} .)



An incomplete list of astrophysically important X-ray spectral features

Identification	Energy (keV)†	Identification	Energy (keV)†
Ne VII	0.127	N VII	0.500
Si XI	0.283	O I K edge	0.532
C I K edge	0.284	O VII	0.569
Si XII	0.303	O VII	0.574
C V	0.308	O VIII	0.654
N I K edge	0.402	O VII	0.666
N VI	0.431	O VII	0.698

continued

Identification	Energy (keV) [†]	Identification	Energy (keV) [†]
Fe I LIII edge	0.707	Si XIII	1.86
Fe I LII edge	0.721	S I K edge	2.472
Fe XVII	0.826	Ar I K edge	3.203
Ne I K edge	0.867	Fe I K α_2	6.391
Ne IX	0.915	Fe I K α_1	6.404
Ne IX	0.922	Fe XXV	6.64
Fe XX	0.996	Fe XXV	6.68
Ne X	1.022	Fe XXV	6.70
Mg I K edge	1.305	Fe XXVI	6.93
Mg XI	1.340	Fe I K β	7.058
Mg XI	1.352	Fe I K edge	7.111
Si K edge	1.839		

[†] λ (Å) = 12.399/ E (keV).

Model X-ray spectral distributions for non-dispersive spectroscopy

$$f(E) = C e^{-\sigma_e(E)N_H} f(S, E) \quad \text{photons cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1},$$

where

C = normalization constant,

N_H = hydrogen column density to source,

$\sigma_e(E)$ = photoelectric cross-section per hydrogen atom for absorption of photons of energy E by interstellar medium. For $E > 3$ keV:

$$\sigma_e(E)N_H \approx (E_0/E)^{8/3}, \quad \text{where } E_0 \text{ is a low energy cutoff parameter.}$$

S is a parameter in the intrinsic spectral shape:

Thermal bremsstrahlung:

$$S = T, \quad f(S, E) = \overline{g(T, E)} e^{-E/kT} / E(kT)^{1/2},$$

where $\overline{g(T, E)}$ is the temperature-averaged Gaunt factor.

Power law:

$$S = n, \quad f(S, E) = E^{-n}.$$

Blackbody:

$$S = T, \quad f(S, E) = E^2 / (e^{-E/kT} - 1).$$

For optically identified sources:



$$N_H \approx (2.22 \pm 0.3) \times 10^{21} A_v \quad \text{cm}^{-2 \dagger}$$



$$A_v = 3.0 E_{B-v} \quad (\text{color excess}).$$

[†] (Gorenstein, P., *Ap. J.*, 198, 40, 1975.)

Emission lines for low density plasma


Power per unit emission integral for some of the strong emission lines for a low density plasma as a function of temperature. Elemental abundances characteristic of the solar corona have been assumed: $A(\log N) = H(12.00)$, He (11.30), C (8.70), O (8.50), Ne (7.60), Mg (7.50), Si (7.70), S (7.30), Ca (6.30), Fe (7.70), and Ni (6.70). (Adapted from Tucker, W. H. and Blumenthal, G. R. in *X-ray Astronomy*, R. Giacconi & H. Gursky, eds., D. Reidel Publishing Co., Dordrecht, 1974.)





Plasma emission – power in lines

The power in lines radiated in various wavelength bands as a function of temperature for a low density plasma. Elemental abundances characteristic of the solar corona have been assumed. The power radiated in Bremsstrahlung (B) and radiative recombination (R) is also shown. (Adapted from Kato, T., *Ap. J. Suppl.*, **30**, 397, 1976.)



Bibliography

- X-ray Astronomy*, R. Giacconi & H. Gursky, eds., D. Reidel Publishing Co., 1974.
- 'Soft X-ray spectrum of a hot plasma', Raymond, J. C. & Smith, B. W., *Ap. J. Suppl.*, **30**, 419 (1977).
- 'Radiation from a hot, thin plasma from 1 to 250 Å', Kato, T., *Ap. J. Suppl.*, **30**, 397 (1976).
- 'Radiation from a high-temperature, low-density plasma: the X-ray spectrum of the solar corona', Tucker, W. H. & Koren, M., *Ap. J.*, **168**, 283 (1971).

'Calculated solar X-radiation from 1 to 60 Å', Mewe, R., *Solar Phys.*, **22**, 459 (1972).

'Calculated solar X-radiation', Mewe, R., *Solar Phys.*, **44**, 383 (1975).


Chapter 7

Gamma-ray astronomy

	<i>page</i>
Gamma-ray source map	212
<i>The Second COS-B Catalog of Gamma-ray Sources</i>	212
Intensities of X- and gamma-ray sources	214
<i>Very high energy gamma-rays</i>	214
Crab Nebula spectra	215
Diffuse gamma-ray background	216
Spectral features	217
Downward gamma-ray flux	218
Bibliography	218

Gamma-ray source map

Region of the sky searched for gamma-ray sources by COS-B (unshaded) and sources detected above 100 MeV. The closed circles denote sources with intensities $\geq 1.3 \times 10^{-6}$ photons $\text{cm}^{-2} \text{s}^{-1}$; open circles denote sources below this level. (Adapted from Scarsi, L. *et al.*, *Proc. 12th ESLAB Symp.*, 1977.)



The Second COS-B Catalog of Gamma-ray Sources

Source name	Galactic coord.		Error radius (°)	Flux ^(a)	Spectral ^(b) hardness	Identification
	<i>l</i> (°)	<i>b</i> (°)				
2CG 006-00	6.7	-0.5	1.0	2.4	0.39 ± 0.08	
2CG 010-31	10.5	-31.5	1.5	1.2	...	
2CG 013+00	13.7	+0.6	1.0	1.0	0.68 ± 0.14	
2CG 036+01	36.5	+1.5	1.0	1.9	0.27 ± 0.07	
2CG 054+01	54.2	+1.7	1.0	1.3	0.20 ± 0.09	
2CG 065+00	65.7	0.0	0.8	1.2	0.24 ± 0.09	
2CG 075+00	75.0	0.0	1.0	1.3	...	
2CG 078+01	78.0	+1.5	1.0	2.5	...	
2CG 095+04	95.5	+4.2	1.5	1.1	...	
2CG 121+04	121.0	+4.0	1.0	1.0	0.43 ± 0.12	
2CG 135+01	135.0	+1.5	1.0	1.0	0.31 ± 0.10	
2CG 184-05	184.5	-5.8	0.4	3.7	0.18 ± 0.04	PSR 0531+21 (Crab)
2CG 195+04	195.1	+4.5	0.4	4.8	0.33 ± 0.04	
2CG 218-00	218.5	-0.5	1.3	1.0	0.20 ± 0.08	
2CG 235-01	235.5	-1.0	1.5	1.0	...	

Source name	Galactic coord.		Error radius (")	Flux ^(a)	Spectral ^(b) hardness	Identification
	<i>l</i> (")	<i>b</i> (")				
2CG 263-02	263.6	-2.5	0.3	13.2	0.36 ± 0.02	PSR 0833-45 (Vela)
2CG 284-00	284.3	-0.5	1.0	2.7	...	
2CG 288-00	288.3	-0.7	1.3	1.6	...	
2CG 289+64	289.3	+64.6	0.8	0.6	0.15 ± 0.07	3C 273
2CG 311-01	311.5	-1.3	1.0	2.1	...	
2CG 333+01	333.5	+1.0	1.0	3.8	...	
2CG 342-02	342.9	-2.5	1.0	2.0	0.36 ± 0.09	
2CG 353+16	353.3	+16.0	1.5	1.1	0.24 ± 0.09	ρ Oph Cloud
2CG 356+00	356.9	+0.3	1.0	2.6	0.46 ± 0.12	
2CG 359-00	259.5	-0.7	1.0	1.8	...	

^(a) Flux: $E > 100$ MeV (10^{-6} photons $\text{cm}^{-2} \text{s}^{-1}$).


^(b) Intensity ($E > 300$ MeV)/Intensity ($E > 100$ MeV), assuming E^{-2} spectra in calculating both intensities.

Notes: error radius defines approximately the 90% confidence error circle; fluxes are approximate, since an E^{-2} spectrum is assumed for simplicity.

(Adapted from Swanenburg, B. N. *et al.*, *Ap. J. (Letters)*, **243**, L69, 1981.)

Intensities of X- and gamma-ray sources

Compilation of the intensities of a variety of X- and gamma-ray sources. The ordinate, the number of photons $\text{cm}^{-2} \text{s}^{-1} \text{MeV}^{-1}$, is multiplied by E^2 . (Adapted from Schönfelder, V. in *Non-Solar Gamma-Rays*, R. Cowsik & R. Wills, eds., Pergamon Press, 1980.)



Very high energy gamma-rays

Cen A: $\bar{F}_{\gamma}(\geq 300 \text{ GeV}) \simeq (4.4 \pm 1) \times 10^{-11} \text{ photons cm}^{-2} \text{ s}^{-1}$


(Grindlay, J. *et al.*, *Ap. J.* (Letters), **197**, L9, 1975)

Crab Pulsar: $(\Delta t)F_{\gamma}(\geq 800 \text{ GeV}) = 4.0 \times 10^{-12} \text{ photons cm}^{-2} \text{ s}^{-2}$


(Grindlay, J. *et al.*, *Ap. J.*, **209**, 592, 1976)

Crab Nebula spectra

Photon number spectrum of the total Crab emission. (Adapted from Schönfelder, V., *op. cit.*, see reference for explanation of symbols.)




Photon number spectrum of the Crab pulsar. (Adapted from Schönfelder, V., *op. cit.*, see reference for explanation of symbols.)



Diffuse gamma-ray background

Spectrum of diffuse gamma-ray background. (Adapted from Fichtel, C. E., Simpson, G. A. & Thompson, D. J., *Ap. J.*, **22**, 833, 1978.)




An incomplete list of astrophysically important gamma-ray spectral features

Identification	Energy (MeV)	Identification	Energy (MeV)
Cf-249	0.34	N-14	2.313
Cf-249	0.39	Ne-20	2.613
Annihil. rad.	0.511	O-16	2.741
Ni-56	0.812	Mg-24	2.754
Fe-56	0.847	Ne-20	3.34
Co-56	0.847	C-12	4.438
Fe-56	1.238	N-14	5.105
Mg-24	1.369	O-16	6.129
Ne-20	1.634	Si-28	6.878
Si-28	1.779	O-16	6.917
Al-26	1.81	O-16	7.117
Neutron capture	2.23		

Downward gamma-ray flux

Measurements of the total downward gamma-ray flux at 5 g cm^{-2} over Palestine, Texas. See original work for references. (From Gehrels, N., Instrumental background in balloon-borne gamma-ray spectrometers and techniques for its reduction, *NASA Technical Memorandum 86162*, 1985.)



Bibliography

Gamma-Ray Astronomy, Chupp, E. L., D. Reidel Publishing Co., 1976.

Chapter 8

Cosmic rays

	<i>page</i>
Chemical composition of primary cosmic rays	220
Relative abundances	221
Abundances in the galactic cosmic rays	222
Relative abundances of nuclei	223
Cosmic ray energy spectra	224
Cutoff rigidity	225
Particle production in the atmosphere	226
Gamma-ray production in the atmosphere	227
Altitude variation of cosmic rays	228
Atmospheric depth	229
Pressure and atmospheric thickness	230
Bibliography	230

Chemical composition of primary cosmic rays

Particle group	Charge	Average atomic weight	Intensity†	Corresponding nuclear intensity
Protons	1	1	1300 ± 100	1300
α -particles (^3He , ^4He)	2	4	94 ± 4	376
Light nuclei (Li, Be, B)	3-5	10	2.0 ± 0.3	20
Middle nuclei (C, N, O, F)	6-9	14	6.7 ± 0.3	94
Heavy nuclei	≥ 10	31	2.0 ± 0.3	62
Very heavy nuclei	≥ 20	51	0.5 ± 0.2	25

† Number of particles with energies greater than 2.5 BeV nucleon⁻¹ in units of m⁻² s⁻¹ sr⁻¹.

Relative abundances


Relative abundances of the chemical elements in the cosmic rays and in the solar system normalized to 100 at carbon.

Element	Z	Cosmic rays	Solar system
H	1	26 000	270 000
He	2	3600	18 728
Li	3	18 ± 2	4.2×10^{-4}
Be	4	10.5 ± 1	6.9×10^{-6}
B	5	28 ± 1	3.0×10^{-3}
C	6	100	100
N	7	25 ± 2	31.7
O	8	91 ± 4	182
F	9	1.7 ± 0.4	2.1×10^{-2}
Ne	10	16 ± 2	29.2
Na	11	2.7 ± 0.4	0.51
Mg	12	19 ± 1	8.99
Al	13	2.8 ± 1	0.72
Si	14	14 ± 2	8.47
P	15	0.6 ± 0.2	8.1×10^{-2}
S	16	3 ± 0.4	4.24
Cl	17	0.5 ± 0.2	4.83×10^{-2}
A	18	1.5 ± 0.3	0.99
K	19	0.8 ± 0.2	3.6×10^{-2}
Ca	20	2.2 ± 0.5	0.611
Sc	21	0.4 ± 0.2	3.0×10^{-4}
Ti	22	1.7 ± 0.3	2.35×10^{-2}
V	23	0.7 ± 0.3	2.22×10^{-3}
Cr	24	1.5 ± 0.4	0.108
Mn	25	0.9 ± 0.2	7.88×10^{-2}
Fe	26	10.8 ± 1.4	7.03
Co	27	<0.2	1.87×10^{-2}
Ni	28	0.4 ± 0.1	0.407
Cu	29		4.58×10^{-3}
Zn	30		1.05×10^{-2}
	31-35	$c. 5 \times 10^{-3}$	2.1×10^{-3}
	36-40	$c. 5 \times 10^{-4}$	9.4×10^{-4}
VVH	41-60	$c. 5 \times 10^{-4}$	3.0×10^{-4}
	61-80	$c. 2 \times 10^{-4}$	4.7×10^{-5}
	>80	$c. 10^{-4}$	4.0×10^{-5}

(From Hillier, R., *Gamma Ray Astronomy*, Clarendon Press, Oxford, 1984.)

Abundances in the galactic cosmic rays

Comparison of the abundances of the elements in the galactic cosmic rays with the solar abundances (normalized to C). (Courtesy of C. Meyer, University of Chicago.)




Relative abundances of nuclei normalized to oxygen

Element	Solar flare cosmic rays	Sun	Galactic cosmic rays
^1H	700	1000	350
^2He	107 ± 14	~ 100	50
^3Li	...	$<< 0.001$	0.3
^4Be ^5B	< 0.02	$<< 0.001$	0.8
^{12}C	0.59 ± 0.07	0.6	1.8
^{14}N	0.19 ± 0.04	0.1	≈ 0.8
^{16}O	1.0	1.0	1.0
^{19}F	< 0.03	$<< 0.001$	≈ 0.1
^{20}Ne	0.13 ± 0.02	?	0.30
^{23}Na	...	0.002	0.19
^{24}Mg	0.043 ± 0.011	0.027	0.32
^{27}Al	...	0.002	0.06
^{28}Si	0.033 ± 0.011	0.035	0.12
^{31}P ^{56}Fe	0.057 ± 0.017	0.032	0.13
^{48}Ti ^{60}Ni	≤ 0.02	0.006	0.28


(Adapted from Johnson, F. S., ed., *Satellite Environment Handbook*, Stanford University Press, 1965.)

Cosmic ray energy spectra

Energy spectrum of the primary cosmic ray electrons.



Integral energy spectrum of cosmic ray particles.



Cutoff rigidity

The Earth's magnetic field affects the penetration of charged particles in the vicinity of the Earth. The minimum rigidity (cutoff rigidity) necessary to reach some geomagnetic latitude L and geocentric radius R is given by:

$$\frac{pc}{ze} = \frac{M}{R^2} \frac{\cos^4 L}{[(1 + \cos \theta \cos^3 L)^{1/2} + 1]^2},$$

where

M is the Earth's dipole moment,


$\left(\frac{pc}{ze}\right)$ is the magnetic rigidity of the particle; for charge $z = 1$ it is numerically equal, when expressed in volts, to the momentum in units of ev/c ,

$\left(\frac{M}{R_0^2}\right) \approx 60 \times 10^9$ volts, where R_0 is the radius of the Earth,

θ is the angle between the direction of arrival of the particle and the tangent to the circle of latitude. ($\theta = 0$ corresponds to arrival from the *west* for *positive* particles; $\theta = 0$ corresponds to arrival from the *east* for *negative* particles.)


Particle production in the atmosphere

Schematic representation of the development of particle production in the atmosphere. (Adapted from Simpson *et al.*, *Phys. Rev.*, **90**, 934, 1953.)




Gamma-ray production in the atmosphere

Schematic diagram of gamma-ray production processes in the atmosphere. Neutrinos are ignored. (From Allkofer, O. C. & Grieder, P. K. F., *Cosmic Rays on Earth*, Physik Daten, ISSN 0344-8401, 1984.)




Altitude variation of cosmic rays

Altitude variation of the main cosmic ray components. (From Allkofer, O. C. & Grieder, P. K. F., *Cosmic Rays on Earth*, Physik Daten, ISSN 0344-8401, 1984.)




Atmospheric depth

Relation between atmospheric depth and altitude for an isothermal atmosphere. (From Allkofer, O. C. & Grieder, P. K. F., *Cosmic Rays on Earth*, Physik Daten, ISSN 0344 8401, 1984.)




Relation between zenith angle and atmospheric depth at sea level in an isothermal atmosphere. (From Allkofer, O. C. & Grieder, P. K. F., *Cosmic Rays on Earth*, Physik Daten, ISSN 0344 8401, 1984.)



Pressure and atmospheric thickness

Relations between altitude and pressure, and altitude and depth in the real atmosphere. (After Cole, A. E. & Kantor, A. J., Air Force Reference Atmosphere, AFGL-TR-78-0051, 1978.)



Bibliography

'Cosmic rays on Earth', Allkofer, O. C. & Grieder, P. K. F. in *Physics Data*, ISSN 0344-8401, 1984, nr. 25-1, Fachinformationszentrum, Karlsruhe.

Chapter 9

Earth's atmosphere and environment

	<i>page</i>
Radiation environment	232
<i>Galactic cosmic radiation</i>	232
<i>Solar high energy particle radiation</i>	232
Trapped radiation	233
Solar irradiance	236
<i>Visible and infrared radiation</i>	236
<i>Ultraviolet and X-ray radiation</i>	236
The solar spectrum	237
The international reference atmosphere	238
Altitude variation of atmospheric constituents	239
Opacity of the atmosphere	240
US standard atmosphere	241
Structure of the upper atmosphere	243
Earth's magnetosphere	244
Bibliography	244

Radiation environment

Galactic cosmic radiation

Flux at sunspot minimum:	$\sim 4 \text{ protons cm}^{-2} \text{ s}^{-1}$ (isotropic)
Integrated yearly rate:	$\sim 1.3 \times 10^8 \text{ protons cm}^{-2}$
Flux at sunspot maximum:	$2.0 \text{ protons cm}^{-2} \text{ s}^{-1}$ (isotropic)
Integrated yearly rate:	$\sim 7 \times 10^7 \text{ protons cm}^{-2}$
Energy range:	40 MeV - 10^{13} MeV; predominantly 10^3 – 10^7 MeV
Integrated dose (without shielding):	~ 4 – 10 rad yr^{-1}

Solar high energy particle radiation

Composition: predominantly protons (H^+) and alpha particles (He^{++}).

Integrated yearly flux at 1 AU:

Energy > 30 MeV, $N \approx 8 \times 10^9 \text{ protons cm}^{-2}$ near solar maximum

$N \approx 5 \times 10^5 \text{ protons cm}^{-2}$ near solar minimum

Energy > 100 MeV, $N \approx 6 \times 10^8 \text{ protons cm}^{-2}$ near solar maximum


$N \approx 5 \times 10^4 \text{ protons cm}^{-2}$ near solar minimum

Maximum dosage with shielding of 5 g cm^{-2} (equivalent thickness):


~ 200 rad per week (3 flares), skin dose at a point detector.

Trapped radiation

Electron distribution in the Earth's field. (Published by Vette in August 1964.)




Proton distribution in the Earth's field. (Published by Vette in September 1963.)




Trapped radiation (cont.)

Omni-directional flux in protons $\text{cm}^{-2} \text{s}^{-1}$. (Adapted from Stassinopoulos, E. G., *World Maps of Constant B, L, and Flux Contours*, NASA SP-3054, 1970.)



Omni-directional flux in electrons $\text{cm}^{-2} \text{s}^{-1}$. (Adapted from Stassinopoulos, E. G., *World Maps of Constant B, L, and Flux Contours*, NASA SP-3054, 1970.)



Solar irradiance (1 AU)***Visible and infrared radiation***

Radiant energy distribution:

approximated by that from a 5800 K blackbody

Fraction of solar radiation:

Above 7000 Å = 53.12 %

Above 4000 Å ~ 91.28 %

3000 Å–30 000 Å = 96.62 %

Ultraviolet and X-ray radiation

Fraction of solar radiation:

Below 4000 Å = 8.72 %

Below 3000 Å = 1.21 %

Below 2000 Å = 0.008 % (variable)

Below 1000 Å = 10^{-4} % (variable)


Principal line emission fluxes at 1.0 AU:

Lyman Alpha H I (1215.67 Å): $51.0 \times 10^{-4} \text{ W m}^{-2}$ He II (303.8 Å): $2.5 \times 10^{-4} \text{ W m}^{-2}$ H I (1025.72 Å): $0.60 \times 10^{-4} \text{ W m}^{-2}$ C III (977 Å): $0.50 \times 10^{-4} \text{ W m}^{-2}$ X-ray flux (W m^{-2}):

	1–8 Å	8–20 Å	20–200 Å
Sunspot min	1×10^{-8}	1×10^{-7}	$\sim 1 \times 10^{-4}$
Sunspot max	3×10^{-6}	2×10^{-5}	$\sim 1 \times 10^{-3}$
Flare activity (large flares)	1×10^{-4}	5×10^{-4}	$\sim 1 \times 10^{-2}$


The solar spectrum

The solar spectral irradiance from radio waves to gamma-rays.
(Courtesy H. Malitson and the National Space Science Data Center.)




International reference atmosphere

(COSPAR International Reference Atmosphere, 1961.)




Altitude variation of atmospheric constituents

Variation with altitude of the various constituents of the atmosphere. The horizontal scale is the logarithm of the particle density n in particles cm^{-3} . (Adapted from Pecker, J., *Space Observatories*, D. Reidel Publishing Company, Dordrecht, 1970.)







Opacity of the atmosphere

Attenuation of photons as a function of wavelength for various constituents of the atmosphere. The ordinate is the mass attenuation coefficient in $\text{cm}^2 \text{g}^{-1}$. (a) X-ray and EUV region; (b) UV, visible, and infrared region; (c) radio region. (Adapted from Pecker, J., *Space Observatories*, D. Reidel Publishing Company, Dordrecht, 1970.)




US standard atmosphere, 1976

(a) Mean free path as a function of geometric altitude. (b) Speed of sound as a function of geometric altitude. (c) Mean molecular weight as a function of geometric altitude. (d) Total pressure and mass density as a function of geometric altitude.




US standard atmosphere, 1976 (cont.)


- (e) Dynamic viscosity as a function of geometric altitude.
- (f) Coefficient of thermal conductivity as a function of geometric altitude.
- (g) Kinetic temperature as a function of geometric altitude.
- (h) Mean air-particle speed as a function of geometric altitude.




(e)



(f)




(g)



(h)


Structure of the upper atmosphere

(Adapted from Harris, M. F. in *American Institute of Physics Handbook*, D. E. Gray, ed., McGraw-Hill Book Company, 1972.)



Earth's magnetosphere

Schematic view of the Earth's magnetosphere.



Bibliography

World Maps of Constant B, L, and Flux Contours,
Stassinopoulos, E. G., NASA SP-3054, 1970.

US Standard Atmosphere, 1976, 1976-0-588-256, US Government
Printing Office.

Satellite Environment Handbook, F. S. Johnson, ed., Stanford University
Press, 1965.

Chapter 10

Relativity

	<i>page</i>
Special relativity	246
Lorentz transformation	246
<i>4-Vector transformation</i>	246
<i>Examples of 4-vectors</i>	247
<i>2nd rank tensor transformation</i>	247
<i>Electromagnetic field strength tensor</i>	247
<i>Covariant formulation of Maxwell's equations</i>	247
<i>Lorentz force</i>	247
Cosmology	248
<i>Robertson–Walker line element</i>	248
<i>Einstein field equations</i>	248
<i>Friedmann universes</i>	248
Bibliography	254

Special relativity

Fundamental kinematical relations for a particle of rest mass m_0 and velocity v :

$$\begin{aligned} p &= m_0 v / (1 - v^2/c^2)^{1/2} && \text{momentum} \\ E &= m_0 c^2 / (1 - v^2/c^2)^{1/2} && \text{total energy} \\ T &= E - m_0 c^2 && \text{kinetic energy} \\ m &= m_0 / (1 - v^2/c^2)^{1/2} && \text{relativistic mass} \\ E_0 &= m_0 c^2 && \text{rest energy} \end{aligned}$$

From the above, the following relations can be derived:

$$\begin{aligned} E &= mc^2 = (m_0^2 c^4 + c^2 p^2)^{1/2} \\ p &= [(E/c)^2 - m_0^2 c^2]^{1/2} \\ v &= c^2 p / E = c [1 - (m_0 c^2 / E)^2]^{1/2} = p / [m_0 c^2 + (p/c)^2]^{1/2} \\ m &= E / c^2 = [m_0^2 + (p/c)^2]^{1/2} \end{aligned}$$

Relativistic Doppler effect:

$$1 + z = \frac{1 + (v/c) \cos \theta}{(1 - v^2/c^2)^{1/2}},$$

where

θ = angle between direction of observation and direction of motion,

$\theta = 0$ for motion directly away from observer,

$$z = (\lambda_{\text{obs}} - \lambda) / \lambda.$$

$$z \approx (v/c) \cos \theta \quad \text{for } v \ll c.$$

Lorentz transformation (Gaussian units)

4-Vector transformation

$$B'_\mu = \sum_{\nu=1}^4 a_{\mu\nu} B_\nu \equiv a_{\mu\nu} B_\nu.$$

For a Lorentz transformation from system k to a system k' moving with a velocity parallel to the z -axis, the transformation coefficients are given by:

$$(a_{\mu\nu}) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \gamma & i\gamma\beta \\ 0 & 0 & -i\gamma\beta & \gamma \end{bmatrix}, \quad \beta = v/c, \gamma = (1 - v^2/c^2)^{-1/2}.$$

Examples of 4-vectors

$$x_\mu = (\mathbf{x}, icT),$$

where $\mathbf{x} = x_1\hat{i} + x_2\hat{j} + x_3\hat{k}$.

$$A_\mu = (\mathbf{A}, i\phi),$$

where \mathbf{A} and ϕ are the electromagnetic vector and scalar potential.

$$J_\mu = (\mathbf{J}, ic\rho),$$

where \mathbf{J} and ρ are the current density and charge density.

$$k_\mu = (\mathbf{k}, i\omega/c),$$

where \mathbf{k} and ω are the wave vector and frequency of a plane electromagnetic wave.

$$p_\mu = (\mathbf{p}, iE/c),$$

where \mathbf{p} and E are the momentum and energy of a particle.

2nd rank tensor transformation

$$S_{\mu\nu} = \sum_{\lambda,\sigma=1}^4 a_{\mu\lambda} a_{\nu\sigma} S_{\lambda\sigma} \equiv a_{\mu\lambda} a_{\nu\sigma} S_{\lambda\sigma}.$$

Electromagnetic field strength tensor

$$F_{\mu\nu} = \frac{\partial A_\nu}{\partial x_\mu} - \frac{\partial A_\mu}{\partial x_\nu} = \begin{bmatrix} 0 & B_3 & -B_2 & -iE_1 \\ -B_3 & 0 & B_1 & -iE_2 \\ B_2 & -B_1 & 0 & -iE_3 \\ iE_1 & iE_2 & iE_3 & 0 \end{bmatrix}.$$

Covariant formulation of Maxwell's equations

$$\frac{\partial F_{\mu\nu}}{\partial x_\nu} = \frac{4\pi}{c} J_\mu; \quad \frac{\partial F_{\mu\nu}}{\partial x_\lambda} + \frac{\partial F_{\lambda\mu}}{\partial x_\nu} + \frac{\partial F_{\nu\lambda}}{\partial x_\mu} = 0,$$

where λ, μ , and ν are any three of the integers 1, 2, 3, 4.

Lorentz force

$$\mathbf{f} = \rho\mathbf{E} + \frac{1}{c}(\mathbf{J} \times \mathbf{B}).$$

$$f_\mu = \frac{1}{c} F_{\mu\nu} J_\nu = \frac{1}{4\pi} F_{\mu\nu} \frac{\partial F_{\nu\lambda}}{\partial x_\lambda} = \frac{\partial T_{\mu\nu}}{\partial x_\nu} = \left(\mathbf{f}, \frac{i}{c} \mathbf{E} \cdot \mathbf{J} \right),$$

where $T_{\mu\nu}$ is the electromagnetic stress-energy-momentum tensor:

$$T_{\mu\nu} = \frac{1}{4\pi} [F_{\mu\lambda} F_{\lambda\nu} + \frac{1}{4} \delta_{\mu\nu} F_{\lambda\sigma} F_{\lambda\sigma}].$$

Cosmology

Robertson-Walker line element (homogeneous and isotropic universe)

$$ds^2 = c^2 dt^2 - R^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2 \theta d\phi^2) \right],$$

where

$R(t)$ = radius of curvature of the universe,

r, θ, ϕ = co-moving spherical coordinates (a co-moving observer is an observer at rest with respect to matter in his vicinity),

k = curvature index = 0, ± 1 ($k = 1$, elliptical closed space; $k = 0$, Euclidean flat space; $k = -1$, hyperbolic open space).

Einstein field equations

$$\frac{3\dot{R}^2}{R^2} + \frac{3kc^2}{R^2} = 8\pi G\rho + \Lambda c^2,$$

$$\frac{2\ddot{R}}{R} + \frac{\dot{R}^2}{R^2} + \frac{kc^2}{R^2} = -\frac{8\pi GP}{c^2} + \Lambda c^2,$$

$$\frac{\ddot{R}}{R} = \frac{\Lambda c^2}{3} - \frac{4\pi G}{3} \left[\rho + \frac{3P}{c^2} \right],$$

where

ρ = mean density of matter and energy,

Λ = cosmological constant,

P = hydrodynamic pressure of matter and radiation,

G = gravitational constant,

$H_0 \equiv \dot{R}_0/R_0$, Hubble constant,

$q_0 \equiv -\ddot{R}_0/R_0H_0^2$, deceleration constant, where the subscript zero denotes the present value.

Friedmann universes ($\Lambda = 0$)

$$\rho_0 + \frac{3P_0}{c^2} = \frac{3H_0q_0}{4\pi G},$$

$$\frac{kc^2}{R_0^2} = \frac{4\pi G}{3q_0} \left[\rho_0(2q_0 - 1) - \frac{3P_0}{c^2} \right].$$

For epochs after the decoupling of matter, P_0 is approximately 0, and we have

Curvature	Space	q_0	$\Omega_0^{(a)}$	Density ρ_0	Expansion
$k = +1$	Closed	$> 1/2$	> 1	$> \frac{3H_0^2}{8\pi G}$	Turns eventually into contraction
$k = 0$	Flat (Euclidean)	$1/2$	1	$\rho_c = \frac{3H_0^2}{8\pi G}^{(b)}$	Stops in infinite future
$k = -1$	Open	$0 \leq q_0 < 1/2$	$0 \leq \Omega_0 < 1$	$< \frac{3H_0^2}{8\pi G}$	Forever

^(a) The 'density parameter' $\Omega \equiv \rho/\rho_c$, where ρ_c is the critical closure density (i.e. for the case $q_0 = 1/2$).

^(b) With $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$, the present critical density becomes $\rho_c = 4.7 \times 10^{-30} \text{ g cm}^{-3}$.

Useful relationships and quantities (Friedmann universe)

Differential volume:

$$dV = \frac{R_H^3}{(1+z)^3} \frac{\{q_0 z + (q_0 - 1)[(1 + 2q_0 z)^{1/2} - 1]\}^2}{q_0^4 (1 + 2q_0 z)^{1/2}} d\Omega dz,$$

where

$$R_H = \frac{c}{H_0}, \quad \text{the Hubble radius.}$$

Time differential:

$$dt = -dz / [(1+z)^2 H_0 (1 + 2q_0 z)^{1/2}].$$

Look-back time:

$$\tau = - \int_0^z dt,$$

$q_0 = 0,$

$$\tau = \frac{1}{H_0} (1 - 1/(1+z)); \quad \lim_{z \rightarrow \infty} \tau = 1/H_0,$$

$q_0 = 1/2,$

$$\tau = \frac{2}{3H_0} (1 - 1/(1+z)^{3/2}); \quad \lim_{z \rightarrow \infty} \tau = 2/3(1/H_0),$$

$q_0 = 1,$

$$\tau = \frac{1}{H_0} [(2z+1)^{1/2}/(z+1) + 2 \tan^{-1}(2z+1)^{1/2} - 1 - \pi/2];$$

$$\lim_{z \rightarrow \infty} \tau = 0.57(1/H_0),$$

where

H_0 = Hubble constant,
 q_0 = deceleration constant,
 z = redshift.

Redshift-magnitude relationship (Friedmann universe):

$$m_{\text{bol}} = 5 \log \left\{ \frac{c}{H_0 q_0^2} [1 - q_0 + q_0 z + (q_0 - 1)(2q_0 z + 1)^{1/2}] \right\} \\ + M_{\text{bol}} + 25 = 5 \log D_L + M_{\text{bol}} + 25.$$

where m_{bol} is the apparent bolometric magnitude of a source with absolute bolometric magnitude M_{bol} and redshift z .

Expanded in powers of z :

$$m_{\text{bol}} = 5 \log \left(\frac{zc}{H_0} \right) + 1.086(1 - q_0)z + \dots + M_{\text{bol}} + 25.$$

where H_0 is in $\text{km s}^{-1} \text{Mpc}^{-1}$; zc is in km s^{-1} , and z is the observed redshift.

$$D_L = \frac{c}{H_0 q_0^2} [1 - q_0 + q_0 z + (q_0 - 1)(2q_0 z + 1)^{1/2}] \\ \approx \frac{cz}{H_0} [1 + 0.5z(1 - q_0)], \quad q_0 z \ll 1.$$

$m_{\text{bol}} - M_{\text{bol}} = m - M - K - A$, $m - M$ = observed distance modulus, for heterochromatic magnitudes, where K = redshift correction, A = interstellar absorption.

$$K = 2.5 \log(1 + z) + 2.5 \log \frac{\int_0^x I(\lambda) S(\lambda) d\lambda}{\int_0^x I\left(\frac{\lambda}{1+z}\right) S(\lambda) d\lambda} \quad \text{[mag]},$$

where $I(\lambda)$ is the incident energy flux per unit wavelength and $S(\lambda)$ is the photometer response function.

Angular diameter-redshift relationship (Friedmann universe)

$$\theta = \frac{l(1+z)^2}{D_L},$$

where

θ = apparent angular diameter of source,
 l = linear diameter of spherical source,
 D_L = luminosity distance.

Observed energy flux density (Friedmann universe)

$$S_0(E_0) = \frac{P_e((1+z)E_0)}{4\pi D_L^2}$$


where P_e is the monochromatic power emitted at the energy $(1+z)E_0$, and S_0 is the observed monochromatic energy flux density at the energy E_0 . The observed energy flux density integrated from E_1 to E_2 is:

$$F_0(E_1, E_2) = \frac{1}{4\pi D_L^2} \int_{(1+z)E_1}^{(1+z)E_2} P_e(E) dE,$$

where $P_e(E)$ is the differential power emitted per unit energy in the emitted rest frame.


Redshift functions

Angular size vs. redshift




Redshift functions (cont.)


Luminosity distance vs. redshift




Angular density of sources per unit z vs. redshift.



Integrated angular density vs. redshift.



Redshift functions (cont.)

**Bibliography**

Gravitational and Cosmology, Weinberg, S., John Wiley and Sons, 1971.
General Relativity and Cosmology, 2nd ed. McVittie, G. C., University of Illinois Press, 1965.

Chapter 11

Atomic physics

	<i>page</i>
Atomic physics	256
<i>Spectroscopic terminology</i>	256
<i>Emission and absorption of radiation</i>	256
X-ray atomic energy levels	258
X-ray energy level diagram	261
X-ray wavelengths	262
<i>Wavelengths of K series lines</i>	262
<i>Wavelengths of the more prominent L group lines</i>	264
Bibliography	265

Atomic physics

Spectroscopic terminology

Orbital angular momentum L or l	0	1	2	3	4	5	6	7	8	
L		S	P	D	F	G	H	I	K	L
l		s	p	d	f	g	h	i	k	l

$L = \sum l$ (individual electrons), orbital angular momentum,

$S = \sum s$ (individual electrons), spin angular momentum,

$J = L + S$ (LS coupling), total angular momentum,

$J = \sum j$; $j = l + s$ (jj coupling),

M = magnetic quantum number; components of J in magnetic field.

n (total quantum number)	1	2	3	4	5	6	7
Shell	K	L	M	N	O	P	Q

The quantities n, l, S, L, J, M define a *Zeeman state*.

The quantities n, l, S, L, J define a *level* which includes $2J + 1$ states, e.g., the atomic level $2p^3\ ^4S_{3,2}^0$.

Interpretation:

2: outer electrons, $n = 2$ (L shell).

p^3 : 3 outer electrons, $l = 1$.

4: multiplicity = 4 ($2S + 1 = 4$, $S = 3/2$, the spin).

S : orbital momentum $L = 0$.


$3/2$: $J = 3/2$.

0: the level has odd parity.

The quantities n, l, S, L define an atomic *term*, the set of $(2S + 1) \times (2L + 1)$ states characterized by given values of L and S .

A transition between two *levels* is called a *spectral line*.

The totality of transitions between two *terms* is a *multiplet*.

Emission and absorption of radiation (cgs units)

where

A_{ki} = Einstein coefficient of spontaneous emission,
 N_k = number of atoms per unit volume in level k ,
 B_{ik} = induced transition probability from level i to level k ,
 I_ν = specific intensity of radiation field at frequency ν ,
 $h\nu = W_k - W_i$, transition energy.

$$g_k B_{ki} = g_i B_{ik},$$

$$\frac{g_k}{g_i} A_{ki} = \frac{2h\nu^3}{c^2} B_{ik} = \frac{8\pi^2 e^2 \nu^2}{mc^3} f_{ik},$$

where f_{ik} = absorption oscillator strength.

$$\sigma_i = \int \sigma_\nu d\nu = \frac{\pi e^2}{mc} f_{ik} N_i, \text{ is the}$$

integrated atomic scattering coefficient for a spectrum line.

σ_ν = atomic scattering coefficient near an absorption line.

Thermal equilibrium

$$\frac{N_k}{N_i} = \frac{g_k}{g_i} e^{-(W_k - W_i/kT)} \quad (\text{Boltzmann's formula}).$$

Doppler shift

$$\Delta\lambda/\lambda \approx v/c \quad (v = \text{velocity of source}).$$

Doppler width of spectral line (FWHM, Maxwellian distribution)

$$\frac{\Delta\lambda_D}{\lambda} = \frac{2[(2 \ln 2)kT/M]^{1/2}}{c} = 7.162 \times 10^{-7} \sqrt{\left[\frac{T(\text{K})}{\text{at. wt.}} \right]}$$

where M is the mass of the radiating atom.

X-ray atomic energy levels (eV)

Z	element	K	L_I	L_{II}	L_{III}	M_I	$M_{II,III}$	$M_{IV,V}$	
1	H	13.598							
2	He	24.587							
3	Li	54.75							
4	Be	111.0							
5	B	188.0		4.7					
6	C	283.8		6.4					
7	N	401.6		9.2					
8	O	532.0	23.7	7.1					
9	F	685.4	31	8.6					
10	Ne	866.9	45	18.3					
11	Na	1072.1	63.3	31.1					
12	Mg	1305.0	89.4	51.4					
13	Al	1559.6	117.7	73.1					
14	Si	1838.9	148.7	99.2					
15	P	2145.5	189.3	132.2					
16	S	2472.0	229.2	164.8					
Permitted transitions									
K series									
L series									
$K\alpha_1 = K - L_{III}$ $K\alpha_2 = K - L_{II}$ $K\beta_1 = K - M_{III}$ $K\beta_3 = K - M_{II}$ $K\beta_2 = K - N_{II,III}$									
$L\alpha_1 = L_{III} - M_V$ $L\alpha_2 = L_{III} - M_{IV}$ $L\beta_1 = L_{II} - M_{IV}$									
Z	element	K	L_I	L_{II}	L_{III}	M_I	$M_{II,III}$	$M_{IV,V}$	
17	Cl	2822.4	270.2	201.6	200.0	17.5	6.8		
18	Ar	3202.9	320.	247.3	245.2	25.3	12.4		
19	K	3607.4	377.1	296.3	293.6	33.9	17.8		
20	Ca	4038.1	437.8	350.0	346.4	43.7	25.4		
21	Sc	4492.8	500.4	406.7	402.2	53.8	32.3	6.6	
22	Ti	4966.4	563.7	461.5	455.5	60.3	34.6	3.7	
23	V	5465.1	628.2	520.5	512.9	66.5	37.8	2.2	
24	Cr	5989.2	694.6	583.7	574.5	74.1	42.5	2.3	
25	Mn	6539.0	769.0	651.4	640.3	83.9	48.6	3.3	
26	Fe	7112.0	846.1	721.1	708.1	92.9	54.0	3.6	

27	Co	7708.9	925.6	793.6	778.6	100.7	59.5	2.9												
28	Ni	8332.8	1008.1	871.9	854.7	111.8	68.1	3.6												
29	Cu	8978.9	1096.6	951.0	931.1	119.8	73.6	1.6												
30	Zn	9658.6	1193.6	1042.8	1019.7	135.9	86.6	8.1												
Z element		K	L _I	L _{II}	L _{III}	M _I	M _{II}	M _{III}	M _{IV}	M _V	M _{VI}	M _{VII}	M _{VIII}	M _{IX}	M _X					
31	Ga	10 367.1	1297.7	1142.3	1115.4	158.1	106.8	102.9												
32	Ge	11 103.1	1414.3	1247.8	1216.7	180.0	127.9	120.8												
Z element		K	L _I	L _{II}	L _{III}	M _I	M _{II}	M _{III}	M _{IV}	M _V	N _I	N _{II}	N _{III}							
47	Ag	25 514.0	3805.8	3523.7	3351.1	717.5	602.4	571.4	372.8	366.7	95.2	62.6	55.9							
53	I	33 169.4	5188.1	4852.1	4557.1	1072.1	930.5	874.6	631.3	619.4	186.4	122.7								
54	Xe	34 561.4	5452.8	5103.7	4782.2		999.0	937.0		672.3		146.7								
55	Ca	35 984.6	5714.3	5359.4	5011.9	1217.1	1065.0	997.6	739.5	725.5	230.8	172.3	161.6							
74	W	69 525.0	12 099.8	11 544.0	10 206.8	2819.6	2574.9	2281.0	1871.6	1809.2	595.0	491.6	425.3							
78	Pt	78 394.8	13 879.9	13 272.6	11 563.7	3296.0	3026.5	2645.4	2201.9	2121.6	722.0	609.2	519.0							
79	Au	80 724.9	14 352.8	13 733.6	11 918.7	3424.9	3147.8	2743.0	2291.1	2205.7	758.8	643.7	545.4							
82	Pb	88 004.5	15 860.8	15 200.0	13 035.2	3850.7	3554.2	3066.4	2585.6	2484.0	893.6	763.9	644.5							
92	U	115 606.1	21 757.4	20 947.6	17 166.3	5548.0	5182.2	4303.4	3727.6	3551.7	1440.8	1272.6	1044.9							
Z element		N _{IV}	N _V	N _{VI}	N _{VII}	O _I	O _{II}	O _{III}	O _{IV}	O _V	P _I	P _{II}	P _{III}							
47	Ag																			
53	I	3.3	49.6			13.6														
54	Xe						3.3													
55	Ca	78.8	76.5			22.7	13.1	11.4												
74	W	258.8	245.4	36.5	33.6	77.1	46.8	35.6	6.1											
78	Pt	330.8	313.3	74.3	71.1	101.7	65.3	51.7	2.2											
79	Au	352.0	333.9	86.4	82.8	107.8	71.7	53.7	2.5											
82	Pb	435.2	412.9	142.9	138.1	147.3	104.8	86.0	21.8	19.2	3.1	0.7								
92	U	780.4	737.7	391.3	380.9	323.7	259.3	195.1	105.0	96.3	70.7	42.3	32.3							

(List from Bearden, J. A. and Burr, A. F., *Rev. Mod. Phys.*, **39**, 125, 1967.)


X-ray atomic energy levels (cont.)

Notation for X-ray lines

Siegbahn notation	Transition	Siegbahn notation	Transition	Siegbahn notation	Transition	Siegbahn notation	Transition
$K\alpha_1$	KL_{II}	$L\gamma_2$	L_1N_{II}	$L\nu$	$L_{II}N_{VI}$	$L\beta_2$	$L_{III}N_V$
$K\alpha_2$	KL_{II}	$L\gamma_3$	L_1N_{III}	$L\gamma_8$	$L_{II}O_I$	Lu	$L_{III}N_{VI,VII}$
$K\beta_3$	KM_{II}	$L\gamma_{11}$	L_1N_V	$L\gamma_6$	$L_{II}O_{IV}$	$L\beta_7$	$L_{III}O_I$
$K\beta_1$	KM_{III}	$L\gamma_4$	L_1O_{III}	Ll	$L_{III}M_I$	$L\beta_3$	$L_{III}O_{IV,V}$
$K\beta_5$	$KM_{IV,V}$	$L\gamma_{13}$	$L_1P_{II,III}$	Ll	$L_{III}M_{II}$	$M\gamma$	$M_{III}N_V$
$K\beta_2$	$KN_{II,III}$	$L\eta$	$L_{II}M_I$	Ls	$L_{III}M_{III}$	$M\beta$	$M_{IV}N_{VI}$
$K\beta_4$	$KN_{IV,V}$	$L\beta_{17}$	$L_{II}M_{III}$	$L\alpha_2$	$L_{III}M_{IV}$	$M\zeta_2$	$M_{IV}N_{II}$
$L\beta_4$	L_1M_{II}	$L\beta_1$	$L_{II}M_{IV}$	$L\alpha_1$	$L_{III}M_V$	$M\zeta_1$	$M_{IV}N_{III}$
$L\beta_3$	L_1M_{III}	$L\gamma_5$	$L_{II}N_I$	$L\beta_6$	$L_{III}N_I$	$M\alpha_2$	M_VN_{VII}
$L\beta_{10}$	L_1M_{IV}	$L\gamma_1$	$L_{II}N_{IV}$	$L\beta_{13}$	$L_{III}N_{IV}$	$M\alpha_1$	M_VN_{VII}
$L\beta_9$	L_1M_V						

X-ray energy level diagram

X-ray energy-level diagram for ^{92}U , showing the transitions permitted by the selection rules $\Delta l = \pm 1$; $\Delta j = \pm 1, 0$. (Adapted from Richtmyer, F. K., Kennard, E. H. & Lauritsen, T., *Introduction to Modern Physics*, McGraw-Hill Book Company, 1955.)



X-ray wavelengths

Wavelengths of K series lines representing transitions in the ordinary X-ray energy-level diagram allowed by the selection principles (Ångstrom)

Siegbahn Sommerfeld transition	$K\alpha_2$ $K\alpha'$ $K-L_{II}$	$K\alpha_1$ $K\alpha$ $K-L_{III}$	$K\beta$ $K\beta_2$ $K-M_{II}$	$K\beta_1$ $K\beta$ $K M_{III}$	$K\beta_2$ $K\gamma$ $K L_{II} N_{III}$
4 Be	115.7				
5 B	67.71				
6 C	44.54				
7 N	31.557				
8 O	23.567				
9 F	18.275				
11 Na	11.885			11.594	
12 Mg	9.869			9.539	
13 Al	8.3205			7.965	
14 Si	7.11106			6.7545	
15 P	6.1425			5.7921	
16 S	5.3637	5.3613		5.0211	
17 Cl	4.7212	4.7182		4.3942	
19 K	3.73707	3.73368		3.4468	
20 Ca	3.35495	3.35169		3.0834	
21 Sc	3.02840	3.02503		2.7739	
22 Ti	2.74681	2.74317		2.5090	
23 V	2.50213	2.49835		2.2797	
24 Cr	2.28891	2.28503		2.0806	
25 Mn	2.10149	2.09751		1.90620	
26 Fe	1.936012	1.932076		1.753013	
27 Co	1.78919	1.78529		1.61744	
28 Ni	1.65835	1.65450		1.47905	1.48561
29 Cu	1.541232	1.537395		1.38935	1.37824
30 Zn	1.43603	1.43217		1.29255	1.28107
31 Ga	1.34087	1.33715		1.20520	1.1938
32 Ge	1.25521	1.25130		1.12671	1.11459
33 As	1.17743	1.17344		1.05510	1.04281
34 Se	1.10652	1.10248		0.99013	0.97791
35 Br	1.04166	1.03759		0.93087	0.91853
36 Kr	0.9821	0.9781		0.8767	0.8643
37 Rb	0.92776	0.92364	0.82749	0.82696	0.81476
38 Sr	0.87761	0.87345	0.78183	0.78130	0.76921
39 Y	0.83132	0.82712	0.73972	0.73919	0.72713
40 Zr	0.78851	0.78430	0.70083	0.70028	0.68850
41 Nb	0.74889	0.74465	0.66496	0.66438	0.65280
42 Mo	0.712105	0.707831	0.631543	0.630978	0.619698
43 Te	0.675	0.672		0.601	
44 Ru	0.64606	0.64174	0.57193	0.57131	0.56051
45 Rh	0.61637	0.61202	0.54509	0.54449	0.53396
46 Pd	0.58863	0.58427	0.52009	0.51947	0.50918
47 Ag	0.56267	0.55828	0.49665	0.49601	0.48603

Wavelengths of K series lines representing transitions in the ordinary X-ray energy-level diagram allowed by the selection principles (cont.)

Siegbahn Sommerfeld transition	$K\alpha_2$ $K\alpha'$ $K-L_{II}$	$K\alpha_1$ $K\alpha$ $K-L_{III}$	$K\beta$ $K\beta_2$ $K-M_{II}$	$K\beta_1$ $K\beta$ $K-M_{III}$	$K\beta_2$ $K\gamma$ $K-L_{II}N_{III}$
48 Cd	0.538 32	0.533 90	0.474 71	0.474 08	0.464 20
49 In	0.515 48	0.511 06	0.454 23	0.453 58	0.444 08
50 Sn	0.494 02	0.489 57	0.434 95	0.434 30	0.424 99
51 Sb	0.473 87	0.469 31		0.416 23	0.407 10
52 Te	0.454 91	0.450 37		0.399 26	0.390 37
53 I	0.437 03	0.432 49	0.382 92	0.383 15	0.374 71
54 Xe		0.417		0.360	
55 Cs	0.404 11	0.399 59	0.354 36	0.353 60	0.345 16
56 Ba	0.388 99	0.384 43	0.340 89	0.340 22	0.332 22
57 La	0.374 66	0.370 04	0.328 09	0.327 26	0.319 66
58 Ce	0.361 10	0.356 47	0.315 72	0.315 01	0.307 70
59 Pr	0.348 05	0.343 40	0.304 39	0.303 60	0.296 25
60 Nd	0.335 95	0.331 25	0.293 51	0.292 75	0.285 73
62 Sm	0.313 02	0.308 33	0.273 25	0.272 50	0.265 75
63 Eu	0.302 65	0.297 90	0.263 86	0.263 07	0.256 45
64 Gd	0.292 61	0.287 82	0.254 71	0.253 94	0.247 62
65 Tb	0.282 86	0.278 20	0.246 29	0.245 51	0.239 12
66 Dy	0.273 75	0.269 03	0.237 87	0.237 10	0.231 28
67 Ho	0.264 99	0.260 30			
68 Er	0.256 64	0.251 97	0.223 00	0.222 15	0.216 71
69 Tm	0.248 61	0.243 87	0.215 58	0.214 87	
70 Yb	0.240 98	0.230 28	0.209 16	0.208 34	0.203 22
71 Lu	0.233 58	0.2282	0.202 52	0.201 71	0.196 49
72 Hf	0.226 53	0.221 73	0.195 83	0.195 15	0.190 42
73 Ta	0.219 73	0.214 88		0.189 91	0.184 52
74 W	0.213 37	0.208 56	0.184 75	0.183 97	0.179 06
76 Os	0.201 31	0.196 45		0.173 61	0.168 75
77 Ir	0.195 50	0.190 65		0.168 50	0.163 76
78 Pt	0.190 04	0.182 23		0.163 70	0.158 87
79 Au	0.184 83	0.179 96		0.159 02	0.154 26
81 Tl	0.174 66	0.169 80		0.150 11	0.145 39
82 Pb	0.170 04	0.165 16		0.146 06	0.141 25
83 Bi	0.165 25	0.160 41		0.142 05	0.136 21
92 U	0.130 95	0.126 40		0.111 87	0.108 42

(From *Smithsonian Physical Tables*.)

Wavelengths of the more prominent L group lines (Ångstrom)

Siegbahn Sommerfeld transition	α_2		α_1		β_1		l		η	
	L_{III}	M_{IV}	L_{III}	M_V	L_{II}	M_V	L_{III}	M_I	L_{II}	M_I
16 S		...							83.75	
20 Ca		36.27					40.90			
21 Sc		31.37					35.71			
22 Ti		27.37					31.33			
23 V		24.31					27.70			
24 Cr		21.53			21.19		23.84		23.28	
25 Mn		19.40			19.04		22.34			
26 Fe		17.57			17.23		20.09		19.76	
27 Co		15.93			15.63		18.25		17.86	
28 Ni		14.53			14.25		16.66		16.28	
29 Cu		13.306			13.027		15.26		14.87	
30 Zn		12.229			11.960		13.97		13.61	
31 Ga		11.27			11.01		12.89		12.56	
32 Ge		10.415			10.153		11.922		11.587	
33 As		9.652			9.395		11.048		10.711	
34 Se		8.972			8.718		10.272		9.939	
35 Br		8.358			8.109		9.564		9.235	
37 Rb		7.3027								
38 Sr		6.8486			6.610		7.822		7.506	
39 Y		6.4357			6.2039		...		7.0310	
							β_2		γ_1	
							γ		δ	
							$L_{III}-N_V$		$L_{II}-N_{IV}$	
40 Zr		6.057			5.8236		5.5742		5.3738	
41 Nb	5.718		5.7120		5.4803		5.2260		5.0248	
42 Mo	5.401		5.3950		5.1665		4.9100			
44 Ru	4.8437		4.8357		4.6110		4.3619		4.1728	
45 Rh	4.5956		4.5878		4.3640		4.1221		3.9357	
46 Pd	4.3666		4.3585		4.1373		3.9007		3.7164	
47 Ag	4.1538		4.1456		3.9266		3.6938		3.5149	
48 Cd	3.9564		3.9478		3.7301		3.5064		3.3280	
49 In	3.7724		3.7637		3.5478		3.3312		3.1553	
50 Sn	3.601 51		3.592 57		3.3779		3.168 61		2.994 94	
51 Sb	3.4408		3.4318		3.2184		3.0166		2.8451	
52 Te	3.2910		3.2820		3.0700		2.8761		2.7065	
53 I	3.1509		3.1417		2.9309		2.7461		2.5775	
55 Cs	2.8956		2.8861		2.6778		2.5064		2.3425	
56 Ba	2.7790		2.7696		2.5622		2.3993		2.2366	
57 La	2.6689		2.6597		2.4533		2.2980		2.1372	
58 Ce	2.5651		2.5560		2.3510		2.2041		2.0443	
59 Pr	2.4676		2.4577		2.2539		2.1148		1.9568	
60 Nd	2.3756		2.3653		2.1622		2.0314		1.8738	
62 Sm	2.2057		2.1950		1.9936		1.8781		1.7231	
63 Eu	2.1273		2.1163		1.9163		1.8082		1.6543	

Wavelengths of the more prominent L group lines (cont.)

Siegbahn Sommerfeld transition	α_2		α_1		β_1		l	η
	L_{III}	M_{IV}	L_{III}	M_V	L_{II}	M_V	$L_{III}-M_I$	L_{II} M_I
64 Gd	2.0526		2.0419		1.8425		1.7419	1.5886
65 Tb	1.9823		1.9715		1.7727		1.6790	1.5266
66 Dy	1.9156		1.9046		1.7066		1.6198	1.4697
67 Ho	1.8521		1.8410		1.6435		1.5637	1.4142
68 Er	1.792 02		1.780 68		1.584 09		1.510 94	1.3611
69 Tm	1.7339		1.7228		1.5268		1.4602	1.3127
70 Yb	1.679 42		1.668 44		1.4725		1.412 61	1.265 12
71 Lu	1.6270		1.616 17		1.420 67		1.367 31	1.219 74
72 Hf	1.577 04		1.566 07		1.3711		1.3235	1.1765
73 Ta	1.529 78		1.518 85		1.324 23		1.281 90	1.135 58
74 W	1.484 38		1.473 36		1.279 17		1.242 03	1.096 30
75 Re	1.4410		1.429 97		1.236 03		1.2041	1.0587
76 Os	1.398 66		1.388 59		1.194 90		1.168 84	1.022 96
77 Ir	1.3598		1.348 47		1.155 40		1.132 97	0.988 76
78 Pt	1.321 55		1.310 33		1.117 58		1.099 74	0.955 99
79 Au	1.285 02		1.273 77		1.081 28		1.068 01	0.924 61
80 Hg	1.249 51		1.238 63		1.046 52		1.037 70	0.8946
81 Tl	1.216 26		1.204 93		1.012 99		1.008 22	0.865 71
82 Pb	1.184 08		1.172 58		0.980 83		0.980 83	0.838 01
83 Bi	1.153 01		1.141 50		0.950 02		0.953 24	0.811 43
90 Th	0.965 85		0.954 05		0.763 56		0.791 92	0.651 76
91 Pa	0.9427		0.9309		0.7407		0.7721	0.6325
92 U	0.920 62		0.908 74		0.718 51		0.753 07	0.613 59

(From *Smithsonian Physical Tables*.)**Bibliography***Atomic Energy Levels*, Moore, C. E., NBS Circular 467, US

Government Printing Office.

An Ultraviolet Multiplet Table, Moore, C. E., NBS Circular 488, US

Government Printing Office.

Chapter 12

Electromagnetic radiation

	<i>page</i>
Blackbody radiation	268
<i>Planck functions</i>	268
<i>Radiation curves</i>	269
Synchrotron radiation	270
<i>Single electron in a magnetic field</i>	270
<i>General distribution of electrons</i>	271
<i>Power law distribution of electrons</i>	271
Compton scattering	272
<i>Compton shift</i>	272
<i>Energy of scattered photon</i>	272
<i>Energy of struck electrons</i>	272
<i>Relation between the scattering angles</i>	272
<i>Klein–Nishina cross-section</i>	272
<i>Klein–Nishina differential cross-sections</i>	273
Inverse Compton scattering	274
<i>Total energy loss rate</i>	274
<i>Spectra</i>	275
Hot plasma emission	276
<i>Bremsstrahlung from a hot plasma</i>	276
<i>Non-thermal bremsstrahlung</i>	276
<i>X-ray line emission from a hot plasma</i>	276
<i>Radiative recombination radiation</i>	277
Maxwell's equations	277
Conversion table	278
Standard definitions in radiative transport theory	279
Maxwell's equations in various systems of units	280
Spectrum nomogram	281
Bibliography	281

Blackbody radiation (cgs units)*Planck functions* (brightness of a blackbody)

$$B_\nu(T) = \frac{2h\nu^3}{c^2} \frac{1}{\left(\exp\left(\frac{h\nu}{kT}\right) - 1\right)} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}$$

$$B_\lambda(T) = \frac{2hc^2}{\lambda^5} \frac{1}{\left(\exp\left(\frac{hc}{\lambda kT}\right) - 1\right)} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ cm}^{-1} \text{ sr}^{-1}$$

$$B_{\tilde{\nu}}(T) = \frac{2hc^2\tilde{\nu}^3}{\left(\exp\left(\frac{hc\tilde{\nu}}{kT}\right) - 1\right)} \text{ erg cm}^{-2} \text{ s}^{-1} (\text{cm}^{-1})^{-1} \text{ sr}^{-1}$$

$$B_\nu(T) d\nu = B_\lambda(T) d\lambda = B_{\tilde{\nu}}(T) d\tilde{\nu}$$

Rayleigh Jeans law

$$h\nu/kT \ll 1$$

$$B_\nu(T) = 2\left(\frac{\nu}{c}\right)^2 kT$$

Wien's law

$$h\nu/kT \gg 1$$

$$B_\nu(T) = \frac{2h\nu^3}{c^2} \exp\left(-\frac{h\nu}{kT}\right)$$

Stefan Boltzmann law

$$\text{total emittance} = \pi \int_0^\infty B_\nu(T) d\nu = \sigma T^4 \text{ erg cm}^{-2} \text{ s}^{-1}$$

$$\text{where } \sigma = \frac{2\pi^5 k^4}{15c^2 h^3} = 5.67 \times 10^{-5} \text{ erg cm}^{-2} \text{ deg}^{-4} \text{ s}^{-1}$$

*Wien displacement law**Maximizing B_ν :*

$$\nu_m = 5.9 \times 10^{10} T \text{ Hz}$$

$$\lambda_m = 0.51 T^{-1} \text{ cm}$$

Maximizing B_λ :

$$\nu_m = 10.3 \times 10^{10} T \text{ Hz}$$

$$\lambda_m = 0.29 T^{-1} \text{ cm}$$


Mean photon energy

$$\langle hv \rangle = \frac{\int_0^\infty B_\nu(T) d\nu}{\int_0^\infty (B_\nu(T)/hv) d\nu} = \left(\frac{\zeta(4)}{\zeta(3)} \right) \left(\frac{\Gamma(4)}{\Gamma(3)} \right) kT = 2.7012kT.$$

where $\zeta(n)$ = Riemann zeta function; $\Gamma(n)$ = gamma function.

Radiation curves

Planck-law radiation curves. (Adapted from Kraus, J. D., *Radio Astronomy*, McGraw-Hill Book Company, 1966.)



Synchrotron radiation (cgs units)

Single electron in a magnetic field

Total radiated power:

$$P = 1.6 \times 10^{-15} \gamma^2 B^2 \beta^2 \sin^2 \alpha \quad \text{erg s}^{-1},$$

where

$$\gamma = (1 - \beta^2)^{-1/2} = E/mc^2,$$

E = total energy of particle,

$$\beta = v/c,$$

B = magnetic induction in Gauss,

α = pitch angle, angle between B and velocity vector.

Synchrotron lifetime:

$$t_s = \frac{3 \times 10^8}{\gamma B^2 \beta^2 \sin^2 \alpha} \quad \text{s.}$$

Spectrum:


$$P(\nu) = 2.3 \times 10^{-22} B \sin \alpha F(\nu/\nu_c) \quad \text{erg s}^{-1} \text{ Hz}^{-1} \quad (\alpha \gg 1/\gamma).$$

$$\nu_c = 4.3 \times 10^6 B \gamma^2 \sin \alpha \quad \text{Hz} \quad (\text{critical frequency}).$$

$$F(x) = x \int_x^\infty d\xi K_{5/3}(\xi), \quad x = \nu/\nu_c.$$

$K_{5/3}(\xi)$ is the modified Bessel function of fractional order 5/3.

A plot of the function $F(x)$ or, equivalently, the dimensionless synchrotron spectrum. $F(\nu/\nu_c)$ reaches its maximum value of 0.918 at $\nu_m = 0.29\nu_c$. (Adapted from Tucker, W. H. & Blumenthal, G. R. in *X-ray Astronomy*, R. Giacconi & H. Gursky, eds., D. Reidel Publishing Company, Dordrecht, 1974.)



General distribution of electrons

$$\frac{dP}{dV dv} = \int \int d\gamma d\Omega_x n(\gamma, \alpha) P(\nu), \text{ the spectral emission per unit volume,}$$


where $n(\gamma, \alpha) d\gamma d\Omega_x$ = density of electrons with Lorentz factor between γ and $\gamma + d\gamma$ and pitch angle between α and $\alpha + d\alpha$; $d\Omega_x = 2\pi \sin \alpha d\alpha$; $P(\nu)$ = single electron spectrum.

Power law distribution of electrons

$$n(\gamma, \alpha) = N\gamma^{-s} g(\alpha) / 4\pi,$$


and for local isotropy $g(\alpha) = 1$.

$$\frac{dP}{dV dv} = 1.7 \times 10^{-21} N a(s) B (4.3 \times 10^6 B/\nu)^{(s-1)/2} \text{ erg s}^{-1} \text{ cm}^{-3} \text{ Hz}^{-1}.$$



(Tucker, W. H. & Blumenthal, G. R., *op. cit.*)

Compton scattering (cgs units)

**Compton shift**

$$\frac{c}{\nu'} - \frac{c}{\nu_0} = \lambda' - \lambda_0 = \frac{h}{m_e c} (1 - \cos \theta).$$

Energy of scattered photon

$$h\nu' = \frac{m_e c^2}{1 - \cos \theta + (1/\alpha)}, \quad \alpha = h\nu_0/m_e c^2.$$

Energy of the struck electron

$$T = h\nu' - h\nu_0,$$

$$T = h\nu_0 \frac{\alpha(1 - \cos \theta)}{1 + \alpha(1 - \cos \theta)},$$

$$T_{\max} = \frac{h\nu_0}{1 + (1/2\alpha)}.$$

Relation between the scattering angles, ϕ and θ

$$\cot \phi = (1 + \alpha) \frac{1 - \cos \theta}{\sin \theta} = (1 + \alpha) \tan \frac{\theta}{2}.$$

Klein-Nishina cross-section for unpolarized incident radiation


$$\frac{d\sigma}{d\Omega} = r_0^2 \left[\frac{1}{1 + \alpha(1 - \cos \theta)} \right]^3 \left(\frac{1 + \cos^2 \theta}{2} \right) \left[1 + \frac{\alpha^2(1 - \cos \theta)^2}{(1 + \cos^2 \theta)[1 + \alpha(1 - \cos \theta)]} \right] \text{ cm}^2 \text{ electron}^{-1} \text{ sr}^{-1}.$$

where $r_0 = \frac{e^2}{m_e c^2}$, classical electron radius = 2.82×10^{-13} cm.


$$\int \frac{d\sigma}{d\Omega} d\Omega = \frac{8\pi}{3} r_0^2 (1 - 2\alpha + 5.2\alpha^2 - 13.3\alpha^3 + \dots) \text{ cm}^2 \text{ electron}^{-1}.$$

Klein Nishina differential cross-sections

Differential cross-sections, $d\sigma(\theta)/d\Omega$, for the production of secondary photons from Compton scattering. Curves are shown for six different values of primary photon energy. (From Davisson & Evans, *Rev. Mod. Phys.*, **34**, 79, 1953.)




Differential cross-sections, $d\sigma(\theta)/d\theta$, for the production of secondary photons from Compton scattering. (From Davisson & Evans, *Rev. Mod. Phys.*, **34**, 79, 1953.)



Klein-Nishina differential cross-sections (cont.)

Differential cross-sections, $d\sigma_e(\phi)/d\phi$, for the production of secondary electrons from Compton scattering. (From Davisson & Evans, *Rev. Mod. Phys.*, **34**, 79, 1953.)

**Inverse Compton scattering (cgs units)**

Compton collisions between relativistic electrons and low frequency photons.

Thomson limit

$$\gamma h\nu_0 \ll m_e c^2,$$

where γ is the Lorentz factor for the relativistic electrons.

Total energy loss rate (Thomson limit)

$$-\frac{dE}{dt} = \frac{4}{3}\sigma_T c \gamma^2 u = 2.6 \times 10^{-14} \gamma^2 u \quad \text{erg s}^{-1} \text{ electron}^{-1},$$

where

u = radiation energy density,

$$\sigma_T = \frac{8\pi}{3} r_0^2, \text{ the Thomson cross-section,}$$

$$r_0 = \frac{e^2}{m_e c^2}.$$


Spectra

Thomson limit power law electron distribution function and blackbody radiation: when the initial radiation field is given by a blackbody distribution and the density of electrons is given by a power law, viz., $N(\gamma) d\gamma = N\gamma^{-s} d\gamma$ (the density of electrons with a Lorentz factor between γ and $\gamma + d\gamma$), the spectral power density is:

$$\frac{dP}{dV d\nu} = 4.2 \times 10^{-40} N b(s) T^3 (2.1 \times 10^{10} T/\nu)^{(s-1)/2}$$

erg cm⁻³ s⁻¹ Hz⁻¹,

with T in degrees Kelvin.



(Tucker, W. H. & Blumenthal, G. R., *op. cit.*)

Hot plasma emission (cgs units)***Bremsstrahlung from a hot plasma***

For a Maxwellian distribution of electron velocities, the spectral emission per unit volume:

$$\frac{dP_B(T)}{dV dv} = 6.8 \times 10^{-38} T^{-1/2} e^{-E/kT} N_e N_Z Z^2 \overline{g_B(T, E)} \quad \text{erg cm}^{-3} \text{ s}^{-1} \text{ Hz}^{-1},$$

where

N_e = electron density,

N_Z = ion density (charge z),

$E = h\nu$ = photon energy, and

g_B is the Gaunt factor.

$$\begin{aligned} \overline{g_B(T, E)} &\approx \frac{\sqrt{3}}{\pi} \ln\left(\frac{4kT}{\Gamma E}\right), \quad \text{for } E \ll kT \quad (\ln \Gamma = 0.577) \\ &\approx (E/kT)^{-0.4}, \quad \text{for } E \sim kT. \end{aligned}$$

The total bremsstrahlung emission:

$$\frac{dP_B}{dV} = 1.4 \times 10^{-27} T^{1/2} N_e N_Z Z^2 g_B(T) \quad \text{erg cm}^{-3} \text{ s}^{-1},$$

where $g_B(T) \approx 1.2$.

$$\frac{dP_B}{dV} = 2.4 \times 10^{-27} T^{1/2} N_e^2 \quad \text{erg cm}^{-3} \text{ s}^{-1}$$

for a plasma with cosmic abundances, since the contribution from all ions

$$\sum N_e N_Z Z^2 \approx 1.4 N_e^2.$$

Non-thermal bremsstrahlung

For a flux density $J(E) = J_0 E^{-s} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ erg}^{-1}$ of non-thermal electrons, the spectral emission per unit volume:

$$\frac{dP_B}{dV dv} = 1.2 \times 10^{-6} Z^2 N_Z J_0 \frac{E^{-s-1}}{s} \quad \text{erg cm}^{-3} \text{ s}^{-1} \text{ Hz}^{-1}.$$

X-ray line emission from a hot plasma (electron collisional excitation)

For a Maxwellian distribution of electron velocities, the power emitted per unit volume due to excitations of level n' of ion Z in the ground state n is:

$$\frac{dP_L}{dV} = 2.7 \times 10^{-15} T^{-1/2} e^{-E_{nn'}/kT} \int_{nn'} \overline{g_{nn'}} N_e N_Z \quad \text{erg cm}^{-3} \text{ s}^{-1},$$

where

$E_{nn'}$ = energy of excitation,

f_{nm} = oscillator strength for the transition,

g_{nm} = mean Gaunt factor ≈ 0.2 for $kT \cdot E_{nm} < 1$.

In some cases, a line is produced by a transition to a state other than the ground state. In this case, the emitted power above is not equal to the power in the line and the branching ratio to the state of interest must be taken into account. For $1s-2p$ transitions in hydrogen and helium-like ions, this branching is not significant.

Radiative recombination radiation

$$\frac{dP_{RR,Z,n}}{dV dE} = \frac{2.8 \times 10^{-6} T^{-3/2} e^{(I_{Z-1,n} - E)/kT}}{n^3} N_e N_Z Z^4 \quad \text{erg cm}^{-3} \text{s}^{-1} \text{erg}^{-1},$$

where

$$E = W_f + I_{Z-1,n},$$

E = energy of emitted photon,

W_f = energy of free electron,

$I_{Z,n}$ = ionization energy of level n for ion Z .

(Adapted from Tucker, W. H. & Blumenthal, G. R. in *X-ray Astronomy*, R. Giacconi & H. Gursky, eds., D. Reidel Publishing Company, Dordrecht, 1974.)

Maxwell's equations (Gaussian units)

$$\nabla \cdot \mathbf{D} = 4\pi\rho, \quad \nabla \times \mathbf{H} = \frac{4\pi\mathbf{J}}{c} + \frac{1}{c} \frac{\partial \mathbf{D}}{\partial t},$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} = 0.$$

Constitutive relations for an isotropic, permeable, conducting dielectric:

$$\mathbf{D} = \epsilon\mathbf{E}, \quad \mathbf{J} = \sigma\mathbf{E}, \quad \mathbf{B} = \mu\mathbf{H}.$$

Macroscopic media:

$$\text{polarization:} \quad \mathbf{P} = \frac{1}{4\pi} (\mathbf{D} - \mathbf{E}),$$

$$\text{magnetization:} \quad \mathbf{M} = \frac{1}{4\pi} (\mathbf{B} - \mathbf{H}).$$

Vector and scalar potentials:

$$\mathbf{B} = \nabla \times \mathbf{A}, \quad \mathbf{E} = -\nabla\phi - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t}.$$

For homogeneous, isotropic media, Maxwell's equations become:

$$\nabla^2 \phi = \frac{\epsilon\mu}{c^2} \frac{\partial^2 \phi}{\partial t^2} - \frac{1}{c} \frac{\partial}{\partial t} \left(\nabla \cdot \mathbf{A} + \frac{\epsilon\mu}{c} \frac{\partial \phi}{\partial t} \right) - \frac{4\pi\rho}{\epsilon}$$

$$\nabla^2 \mathbf{A} = \frac{\mu\epsilon}{c^2} \frac{\partial^2 \mathbf{A}}{\partial t^2} + \nabla \left(\nabla \cdot \mathbf{A} + \frac{\epsilon\mu}{c} \frac{\partial \phi}{\partial t} \right) - \frac{4\pi\mu}{c} \mathbf{J}$$

Lorentz gauge:

$$\left(\nabla \cdot \mathbf{A} + \frac{\epsilon\mu}{c} \frac{\partial \phi}{\partial t} \right) = 0.$$

Coulomb gauge:

$$\nabla \cdot \mathbf{A} = 0.$$

Conversion table for given amount of physical quantity

Physical quantity	Symbol	Rationalized MKS	Gaussian
Charge	q	1 coulomb	3×10^9 statcoulombs
Charge density	ρ	1 coul m ⁻³	3×10^3 statcoulombs cm ⁻³
Current	I	1 ampere	3×10^9 statamperes
Current density	J	1 amp m ⁻²	3×10^5 statamperes cm ⁻²
Electric field	E	1 volt m ⁻¹	$1/3 \times 10^{-4}$ statvolt cm ⁻¹
Potential	ϕ, V	1 volt	1/300 statvolt
Polarization	P	1 coul m ⁻²	3×10^5 statcoulombs cm ⁻²
Displacement	D	1 coul m ⁻²	$12\pi \times 10^5$ statvolt cm ⁻¹
Conductivity	σ	1 mho m ⁻¹	9×10^9 s ⁻¹
Magnetic induction	B	1 weber m ⁻²	10 ⁴ gauss
Magnetic field	H	1 ampere-turn m ⁻¹	$4\pi \times 10^{-3}$ oersted
Magnetization	M	1 weber m ⁻²	$1/4\pi \times 10^4$ gauss

(Adapted from *Classical Electrodynamics*, Jackson, J. D., John Wiley and Sons, 1962.)

Standard definitions in radiative transport theory

Quantity	Symbol	Units (cgs)
Specific intensity or radiance	I_ν	$\text{erg cm}^{-2} \text{s}^{-1} \text{Hz}^{-1} \text{sr}^{-1}$
Brightness	$B_\nu = -I_\nu$	$\text{erg cm}^{-2} \text{s}^{-1} \text{Hz}^{-1} \text{sr}^{-1}$
Flux density	$F_\nu = \int I_\nu \cos \theta \, d\Omega$	$\text{erg cm}^{-2} \text{s}^{-1} \text{Hz}^{-1}$
Mean intensity	$J_\nu = \frac{1}{4\pi} \int I_\nu \, d\Omega$	$\text{erg cm}^{-2} \text{s}^{-1} \text{Hz}^{-1}$
Radiation density	$u_\nu = \frac{1}{c} \int I_\nu \, d\Omega = \frac{4\pi}{c} J_\nu$	$\text{erg cm}^{-3} \text{Hz}^{-1}$
Emission coefficient	j_ν	$\text{erg cm}^{-3} \text{s}^{-1} \text{Hz}^{-1} \text{sr}^{-1}$
Emissivity	$\epsilon_\nu = \frac{4\pi}{\rho} j_\nu$ (isotropic emission, $\rho = \text{density}$)	$\text{erg gm}^{-1} \text{s}^{-1} \text{Hz}^{-1}$
Linear absorption coefficient	$\alpha_\nu = n\sigma_\nu$ ($n = \text{number density}$, $\sigma_\nu = \text{cross-section}$)	cm^{-1}
Mean free path	$l_\nu = \frac{1}{\alpha_\nu}$	cm
Optical depth	$\tau_\nu = \int \alpha_\nu \, ds$	dimensionless


Maxwell's equations in various systems of units ($\epsilon_0, \mu_0, \mathbf{D}, \mathbf{H}$, macroscopic Maxwell's equations, and Lorentz force equation in various systems of units)

System	ϵ_0	μ_0	\mathbf{D}, \mathbf{H}	Macroscopic Maxwell's equations	Lorentz force per unit charge
Electrostatic (esu)	1	c^{-2}	$\mathbf{D} = \mathbf{E} + 4\pi\mathbf{P}$ $\mathbf{H} = c^2\mathbf{B} - 4\pi\mathbf{M}$	$\nabla \cdot \mathbf{D} = 4\pi\rho$ $\nabla \times \mathbf{H} = 4\pi\mathbf{J} + \frac{\partial \mathbf{D}}{\partial t}$ $\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0$ $\nabla \cdot \mathbf{B} = 0$	$\mathbf{E} + \mathbf{r} \times \mathbf{B}$
Electro-magnetic (emu)	c^{-2}	1	$\mathbf{D} = (1/c^2)\mathbf{E} + 4\pi\mathbf{P}$ $\mathbf{H} = \mathbf{B} - 4\pi\mathbf{M}$	$\nabla \cdot \mathbf{D} = 4\pi\rho$ $\nabla \times \mathbf{H} = 4\pi\mathbf{J} + \frac{\partial \mathbf{D}}{\partial t}$ $\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0$ $\nabla \cdot \mathbf{B} = 0$	$\mathbf{E} + \mathbf{r} \times \mathbf{B}$
Gaussian	1	1	$\mathbf{D} = \mathbf{E} + 4\pi\mathbf{P}$ $\mathbf{H} = \mathbf{B} - 4\pi\mathbf{M}$	$\nabla \cdot \mathbf{D} = 4\pi\rho$ $\nabla \times \mathbf{H} = \frac{4\pi}{c}\mathbf{J} + \frac{1}{c}\frac{\partial \mathbf{D}}{\partial t}$ $\nabla \times \mathbf{E} + \frac{1}{c}\frac{\partial \mathbf{B}}{\partial t} = 0$ $\nabla \cdot \mathbf{B} = 0$	$\mathbf{E} + \frac{\mathbf{r}}{c} \times \mathbf{B}$
Heaviside-Lorentz	1	1	$\mathbf{D} = \mathbf{E} + \mathbf{P}$ $\mathbf{H} = \mathbf{B} - \mathbf{M}$	$\nabla \cdot \mathbf{D} = \rho$ $\nabla \times \mathbf{H} = \left(\mathbf{J} + \frac{\partial \mathbf{D}}{\partial t}\right)$ $\nabla \times \mathbf{E} + \frac{1}{c}\frac{\partial \mathbf{B}}{\partial t} = 0$ $\nabla \cdot \mathbf{B} = 0$	$\mathbf{E} + \frac{\mathbf{r}}{c} \times \mathbf{B}$
Rationalized mks	10^7 $4\pi c^2$	$4\pi \times 10^{-7}$	$\mathbf{D} = \epsilon_0\mathbf{E} + \mathbf{P}$ $\mathbf{H} = (1/\mu_0)\mathbf{B} - \mathbf{M}$	$\nabla \cdot \mathbf{D} = \rho$ $\nabla \times \mathbf{H} = \mathbf{J} + \frac{\partial \mathbf{D}}{\partial t}$ $\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0$ $\nabla \cdot \mathbf{B} = 0$	$\mathbf{E} + \mathbf{r} \times \mathbf{B}$

(Adapted from Jackson, J. D., *Classical Electrodynamics*, John Wiley and Sons, 1962.)

Spectrum nomogram

Electromagnetic spectrum nomogram.



Bibliography

- Radiation Processes in Astrophysics*, Tucker, W., The MIT Press, 1975.
Radiative Processes in Astrophysics, Rybicki, G. B. & Lightman, A. P., John Wiley and Sons, 1979.
Classical Electrodynamics, Jackson, J. D., John Wiley and Sons, 1962.
Astrophysical Formulae, Lang, K. R., Springer-Verlag, 1980.

Chapter 13

Plasma physics

	<i>page</i>
Plasma physics parameters	284
<i>Frequencies</i>	284
<i>Velocities</i>	284
<i>Lengths</i>	284
<i>Miscellaneous</i>	285
Approximate parameters	286
Bibliography	287

Plasma physics parameters (Gaussian units)

Z_i = charge on i th ion (units of electronic charge),
 N_e, N_i = electron, ion particle densities, respectively (cm^{-3}),
 T = temperature (K),
 B = magnetic flux density (gauss),
 ρ = density (g cm^{-3}),
 A = mass in amu,
 g = gravitational acceleration (cm s^{-2}),
 $\omega = 2\pi\nu$ (rad s^{-1}).

Frequencies (Hz)

plasma oscillation frequency:

$$\nu_{pe} = (N_e e^2 / \pi m_e)^{1/2} = 8.978 \times 10^3 N_e^{1/2} \text{ Hz},$$

electron gyrofrequency:

$$\nu_{ce} = (e/2\pi m_e c) B = 2.7994 \times 10^6 B,$$

ion gyrofrequency:

$$\nu_{ci} = (Ze/2\pi m_i c) B = 1.535 \times 10^3 Z_i B/A.$$

Velocities (cm s^{-1})

electron rms thermal velocity:

$$v_{Te} = (3kT/m_e)^{1/2} = 6.745 \times 10^5 T^{1/2} \text{ cm s}^{-1},$$

atoms, ions rms thermal velocity:

$$v_{Ti} = (3kT/m)^{1/2} = 1.580 \times 10^4 (T/A)^{1/2},$$

ion sound velocity:

$$c_s = (kT/m_i)^{1/2} = 9.12 \times 10^3 (T/A)^{1/2},$$

Alfvén speed:

$$v_A = B/(4\pi\rho)^{1/2} = 0.282 B/\rho^{1/2},$$

electron drift velocity in crossed magnetic and electric field

$$= 3 \times 10^{10} E_{\perp} (\text{statvolts cm}^{-1})/B(\text{gauss}),$$

electron drift velocity in magnetic and gravitational field

$$= m_e g c / e B = 5.686 \times 10^{-8} g/B.$$

Lengths (cm)

Debye length:

$$\lambda_D = (kT/4\pi e^2 N_e)^{1/2} = 6.92 (T/N_e)^{1/2} \text{ cm}.$$

electron gyroradius:

$$\begin{aligned} a_e &= m_e v_{\perp} / c e B = 5.69 \times 10^{-8} v_{\perp} / B \\ &\simeq 2.21 \times 10^{-2} T^{1/2} B, \end{aligned}$$

ion gyroradius:

$$\begin{aligned} a_i &= m_i v_{\perp} / c Z_i e B = 1.036 \times 10^{-4} v_{\perp} A / Z_i B \\ &\simeq 0.945 T^{1/2} A^{1/2} / Z_i B. \end{aligned}$$

Miscellaneous

electrical resistivity:

$$\eta = 9 \times 10^{-9} (\ln \Lambda) T^{-3/2} \text{ s},$$

Coulomb logarithm:

$$\ln \Lambda \simeq 9.00 + 3.45 \log T - 1.15 \log N_e,$$

thermal conductivity

$$= 1.0 \times 10^{-6} T^{5/2} \text{ erg cm}^{-1} \text{ s}^{-1} \text{ K}^{-1},$$

life of magnetic field in a plasma:

$$\begin{aligned} \tau &= 4\pi L^2 / \eta c^2 \\ &= 1.5 \times 10^{-12} L^2 (\ln \Lambda)^{-1} T^{3/2} \text{ s} \\ & (L \text{ is the characteristic scale of the field}). \end{aligned}$$

Maxwellian velocity distribution:

$$f(v) dv = 4\pi \left(\frac{m}{2\pi kT} \right)^{3/2} e^{-mv^2/(2kT)} v^2 dv,$$

$$\int_0^{\infty} f(v) dv = 1,$$

$$\bar{v} = (8kT/\pi m)^{1/2} \text{ cm s}^{-1},$$

$$\begin{aligned} v_{\text{rms}} &= (3kT/m)^{1/2} \\ &= 6.7 \times 10^5 T^{1/2} \text{ for electrons} \\ &= 1.57 \times 10^4 T^{1/2} \text{ for protons,} \end{aligned}$$

$$\frac{1}{2} m_e v_{\text{rms}}^2 = 3kT/2 = 2.1 \times 10^{-16} T \text{ erg.}$$

Classical skin depth:

$$d \equiv c/2\pi\nu_{pe} = 5.32 \times 10^5 N_e^{-1/2} \text{ cm.}$$

Electron-ion collision frequency:

$$\nu_{ei} \simeq 28 N_i T_e^{-3/2} \frac{\ln \Lambda}{20} \text{ s}^{-1}.$$

Approximate parameters for some astrophysical plasmas

Plasma	N_e (cm^{-3})	T_e (K)	B (gauss)	ν_{pe} (Hz)	λ_D (cm)
Ionosphere	$10^3 - 10^6$	$10^2 - 10^3$	10^{-1}	$3 \times 10^5 - 10^7$	$7 \times 10^{-2} - 7$
Interplanetary space	$1 - 10^4$	$10^2 - 10^3$	$10^{-6} - 10^{-5}$	$10^4 - 10^6$	$7 \times 10^{-1} - 2 \times 10^2$
Solar corona	$10^8 - 10^{12}$ (flare)	$10^6 - 10^7$ (flare)	$10^{-5} - 1$	$10^8 - 10^{10}$	$10^{-2} - 2$
Solar chromosphere	10^{12}	$2 - 3 \times 10^4$	10^3	10^{10}	10^{-3}
Stellar interiors	10^{27}	$10^{7.5}$	—	3×10^{17}	10^{-9}
Planetary nebulae	$10^3 - 10^5$	$10^3 - 10^4$	$10^{-4} - 10^{-3}$	$3 \times 10^5 - 3 \times 10^6$	$7 \times 10^{-1} - 2 \times 10^1$
H II regions	$10^2 - 10^3$	$10^3 - 10^4$	10^{-6}	$10^5 - 3 \times 10^5$	$7 - 7 \times 10^1$
H I regions	10^{-3}	10^2	—	3×10^2	2×10^3
White dwarfs	10^{32}	10^7	10^6 (surface)	10^{20}	2×10^{-12}
Pulsars	10^{42} (center)	—	$10^{1.2}$ (surface)	$10^{2.5}$	—
	10^{12} (surface)	—	—	10^{10}	—
Interstellar space	$10^{-3} - 10$	10^2	10^{-6}	$3 \times 10^2 - 3 \times 10^4$	$2 \times 10^1 - 2 \times 10^3$
Intergalactic space	$\lesssim 10^{-5}$	$10^5 - 10^6$	$\lesssim 10^{-8}$	< 30	2×10^6

N_e = electron density; T_e = electron temperature; B = magnetic field; ν_{pe} = plasma frequency; λ_D = Debye length.

Bibliography

Revised and Enlarged Collection of Plasma Physics Formulas and Data,
Book, D. L., NRL Mem. Report 3332, Naval Research Laboratory,
Washington, DC, 1977.

Chapter 14

Experimental astronomy and astrophysics

	<i>page</i>
Attenuation of electromagnetic radiation	291
<i>Extrapolated absorption and absorption-jump ratios</i>	292
<i>Mass attenuation coefficients, ranges, cross-sections</i>	292
Fluorescence yields	304
Passage of charged particles through matter	304
<i>Total ionization loss by electrons or positrons</i>	304
<i>Radiative loss for non-relativistic electrons</i>	305
<i>Ionization and radiative loss for highly relativistic electrons</i>	306
<i>Total ionization loss by heavy charged particles</i>	307
<i>Approximate range energy relationships</i>	307
<i>Charged particles in silicon</i>	309
Radioactive sources	311
<i>X-ray sources</i>	311
<i>Alpha sources</i>	312
<i>Gamma-ray energy standards</i>	312
<i>Electron energy standards</i>	313
Characteristics of synchrotron radiation	314
<i>Practical formulae for electron storage rings</i>	316
X-ray spectroscopy	318
<i>Crystal spectroscopy</i>	318
<i>Crystal properties</i>	319
<i>Useful characteristic lines for X-ray spectroscopy</i>	320
<i>Grating spectroscopy</i>	320
Reflection of X-rays	322
<i>Optical constants at soft X-ray wavelengths</i>	325
<i>Reflectivity vs. wavelength (energy)</i>	326
Wolter type I mirror system	332

Vacuum technology	333
Equivalents for various cryogenic fluids	334
Optical point spread function	334
<i>The line spread function</i>	335
<i>The edge trace</i>	335
<i>The modulation transfer function</i>	335
<i>The full-width half maximum</i>	336
<i>The root mean square radius</i>	336
<i>The encircled energy function</i>	336
<i>The half-power radius</i>	336
<i>Point spread function for a circular aperture</i>	337
Optical telescopes	339
Photometry	341
<i>Spectral luminous efficiency</i>	341
<i>Summary of typical sources/parameters</i>	343
<i>Conversion table for various photometric units</i>	343
<i>Luminance values for various sources</i>	344
<i>Typical values of natural scene illuminance</i>	344
<i>Radiant responsivity</i>	345
<i>Standard units, symbols, and defining equations</i>	346
Detectors	348
<i>Visible and ultraviolet light detectors</i>	348
<i>X-ray and gamma-ray detectors</i>	358
Microcomputer topics	370
<i>ASCII character code</i>	370
<i>Boolean algebra</i>	373
<i>Logic gates</i>	374
<i>The RS-232-C standard</i>	375
<i>Parallel interface</i>	376
<i>IEEE 488 interface</i>	377
<i>Data transmission</i>	378
Bibliography	378

Attenuation of electromagnetic radiation

$$I = I_0 e^{-(\mu/\rho)t}$$

I_0 = initial intensity of a collimated photon beam,

I = intensity of beam after traversing a thickness t of material density ρ ,

(μ/ρ) = total mass attenuation coefficient

$$= \sigma/\rho + \tau/\rho + k/\rho,$$

σ/ρ = total Compton mass attenuation coefficient,

τ/ρ = photoelectric mass absorption coefficient

$$\left(\sim \frac{Z^4}{A} E^{-8.3} \text{ between absorption edges} \right).$$

k/ρ = pair production mass attenuation coefficient.

Mixtures of materials:

$$\mu/\rho = \sum_i (\mu_i/\rho_i) \omega_i,$$


where

μ_i/ρ_i = mass attenuation coefficient of element i ,

ω_i = fraction by weight of element i .

Photon mean free path = $[(\mu/\rho)\rho]^{-1}$.

Relative importance of the three major types of electromagnetic interactions. The lines show the values of Z and $h\nu$ for which the two neighboring effects are just equal. (Evans, R. D., *The Atomic Nucleus*, McGraw-Hill, 1955, with permission.)



Extrapolated absorption and absorption-jump ratios (r) at the K-edge

Z	element	Atomic weight	Density (g cm ⁻³)	λ_k (Å)	E_k (keV)	Fluor. yield ω_k	(μ, ρ) (cm ² g ⁻¹)	$(\mu/\rho)^+$	r
4	Be	9.01	1.85	112	0.111		179 000	(5000)	(35)
5	B	10.81	2.34	66.0	0.188		88 400	3130	28.3
6	C	12.01	2.26	43.7	0.284		53 900	2230	24.2
7	N	14.01	0.001 25	30.9	0.402		32 800	1540	21.4
8	O	16.00	0.001 43	23.3	0.532		22 400	1160	19.3
9	F	19.00	0.001 70	18.1	0.685		14 800	846	17.5
10	Ne	20.18	0.000 90	14.3	0.867		10 950	687	15.94
11	Na	22.99	0.97	11.56	1.072		7760	525	14.78
12	Mg	24.31	1.74	9.50	1.305		6000	440	13.63
13	Al	26.98	2.70	7.95	1.560	0.036	4500	355	12.68
14	Si	28.09	2.33	6.74	1.839	0.047	3640	307	11.89
15	P	30.97	1.82	5.77	2.146	0.060	2800	251	11.18
16	S	32.06	2.07	5.01	2.472	0.076	2340	222	10.52
17	Cl	35.45	0.003 17	4.38	2.822	0.094	1840	185	9.92
18	Ar	39.95	0.001 78	3.87	3.203	0.115	1440	154	9.34
19	K	39.10	0.86	3.44	3.607	0.138	1300	148	8.79
20	Ca	40.08	1.55	3.07	4.038	0.163	1120	135	8.28

(After Henke, B. L. & Elgin, R. L. in *Advances in X-ray Analysis*, Vol. 13, Plenum Press, New York, 1970.)

Mass attenuation coefficients (cm² g⁻¹)

Wave-length (Å)	Poly-propylene (CH ₂)	Mylar (C ₁₀ H ₈ O ₄)	Teflon (CF ₂)	Air		Methane (CH ₄)	Energy (eV)
				O = 21% N = 78% Ar = 1%	P 10 CH ₄ = 10% Ar = 90%		
2.0	8	14	28	21	230	7	6199.0
4.0	69	116	220	148	162	60	3099.5
6.0	234	384	700	481	467	205	2066.3
8.0	550	870	1540	1090	1020	479	1549.8
10.0	1040	1630	2800	2020	1850	910	1239.8
12.0	1740	2680	4250	3310	3010	1520	1033.2
14.0	2660	4040	6700	4980	4500	2330	885.6
16.0	3830	5800	9400	7100	6400	3350	774.9
18.0	5200	7800	12 600	9500	8400	4570	688.8
20.0	6900	10 200	2780	12 400	10 900	6100	619.9
22.0	8800	12 900	3540	15 700	13 500	7700	563.5
24.0	11 000	8500	4430	14 100	16 400	9700	516.6
26.0	13 500	10 400	5400	17 100	19 600	11 800	476.8
28.0	16 200	12 400	6500	20 400	22 800	14 100	442.8
30.0	19 200	14 700	7800	24 000	26 300	16 800	413.3
32.0	22 400	17 200	9100	2290	29 700	19 600	387.4
34.0	25 900	19 900	10 600	2650	33 300	22 700	364.6


Mass attenuation coefficients (cont.)

Wave-length (Å)	Poly- propylene (CH ₂)	Mylar (C ₁₀ H ₈ O ₄)	Teflon (CF ₂)	Air O = 21% N = 78% Ar = 1%	P 10 CH ₄ = 10% Ar = 90%	Methane (CH ₄)	Energy (eV)
36.0	29 600	22 800	12 100	3040	36 900	25 900	344.4
38.0	33 600	25 800	13 800	3460	40 500	29 300	326.3
40.0	37 800	29 100	15 600	3810	37 600	33 000	309.9
42.0	42 200	32 500	17 500	4270	40 900	36 900	295.2
44.0	1940	3350	6900	4780	42 600	1700	281.8
46.0	2180	3760	7800	5300	45 600	1910	269.5
48.0	2430	4170	8600	5900	48 900	2130	258.3
50.0	2690	4590	9500	6400	52 000	2350	248.0
52.0	2960	5100	10 500	6300		2590	238.4
54.0	3240	5600	11 500	7000		2840	229.6
56.0	3540	6100	12 500	7600		3100	221.4
58.0	3860	6600	13 600	8200		3380	213.8
60.0	4190	7200	14 800	8900		3660	206.6
62.0	4540	7800	16 000	9700		3970	200.0
64.0	4880	8400	17 300	10 400		4270	193.7
66.0	5200	9000	18 600	11 200		4570	187.8
68.0	5700	9700	19 900	12 100		4940	182.3
70.0	6000	10 300	21 300	12 900		5200	177.1
72.0	6400	11 100	22 700	13 800		5600	172.2
74.0	6800	11 800	24 200	14 700		6000	167.5
76.0	7300	12 500	25 700	15 600		6400	163.1
78.0	7700	13 300	27 200	16 600		6700	158.9
80.0	8100	14 100	28 800	17 600		7100	155.0
82.0	8600	14 900	30 500	18 600		7600	151.2
84.0	9100	15 800	32 100	19 700		7900	147.6
86.0	9600	16 600	33 800	20 800		8400	144.2
88.0	10 100	17 500	35 500	21 900		8800	140.9
90.0	10 600	18 400	37 300	23 100		9300	137.8
92.0	11 100	19 400	39 100	24 200		9700	134.8
94.0	11 700	20 300	40 900	25 400		10 300	131.9
96.0	12 200	21 300	43 000	26 700		10 700	129.1
98.0	12 800	22 300	44 600	28 000		11 200	126.5
100.0	13 400	23 300	46 300	29 200		11 800	124.0
105.0	15 000	26 000	51 000	32 700		13 100	118.1
110.0	16 600	28 800	56 000	36 200		14 500	112.7
115.0	18 200	31 600	61 000	39 900		15 900	107.8
120.0	20 000	34 600	67 000	43 800		17 500	103.3
125.0	21 900	38 000	72 000	48 000		19 200	99.2
130.0	23 900	41 100	78 000	52 000		20 900	95.4
135.0	25 900	44 600	83 000	57 000		22 700	91.8
140.0	28 100	48 200	89 000	61 000		24 600	88.6
145.0	30 300	52 000	95 000	65 000		26 500	85.5
150.0	32 600	55 000	101 000	71 000		28 500	82.7


(Adapted from *The Handbook of Chemistry and Physics*, CRC Press, Cleveland, 1976.)

Mass attenuation coefficients (cont.)

Total mass-absorption coefficients for gamma-rays in all elements from Be to U. Energy range 1.8 keV- 10 MeV. (From *Nucleonics*, 19, 62, 1961.)




Mass attenuation coefficients (cont.)



RANGE-ENERGY CURVES FOR PENETRATING RADIATION


(Adapted from Filius, R., *Satellite Instruments Using Solid State Detectors*, Ph.D. thesis, State University of Iowa, Iowa City, 1963.)



(Adapted from Peterson, L. in *Annual Review of Astronomy and Astrophysics*, Annual Review Inc., Palo Alto, California, 1975.)



Mass attenuation coefficients (cont.)

Photoelectric mass absorption coefficients for various window or filter materials.




Mass attenuation coefficients (cont.)

Photoelectric mass absorption coefficients for various gases.




Mass attenuation coefficients (cont.)


Photoelectric cross-section of xenon and argon as a function of photon energy. (Adapted from Anderson, F. D., *A High Resolution Large Area Gas Scintillation Proportional Counter for Use in X-ray Astronomy*, Ph.D. thesis, Columbia University, 1978.)




Photon mean free path versus energy for various gases at STP.



Mass attenuation coefficients for photons in air. The curve marked 'total absorption' is $(\sigma_a/\rho) = (\sigma_{\kappa}/\rho) + (\tau/\rho) + (\kappa/\rho)$, where σ_a , τ , and κ are the corresponding linear coefficients for Compton absorption, photoelectric absorption, and pair production. When the Compton scattering coefficient σ_s is added to μ_a , we obtain the curve marked 'total attenuation' which is $(\mu_0/\rho) = (\mu_a/\rho) + (\sigma_s/\rho)$. The total Rayleigh scattering coefficient is elastic and is confined to small angles, it has not been included in μ_0/ρ . In computing these curves, the composition of 'air' was taken as 78.04 volume per cent nitrogen, 21.02 volume per cent oxygen, and 0.94 volume per cent argon. At 0°C and 760 mm Hg pressure, the density of air is $\rho = 0.001293 \text{ g cm}^{-3}$. (Adapted from Evans, R. D., *The Atomic Nucleus*, McGraw-Hill, 1955, with permission.)




Mass attenuation coefficients for photons in water. (From Evans, R. D., op. cit.)



Mass attenuation coefficients (cont.)


Mass attenuation coefficients for photons in lead.

(From Evans, R. D., *op. cit.*)




Mass attenuation coefficients for photons in aluminum.


(From Evans, R. D., *op. cit.*)



Mass attenuation coefficients for sodium iodide. The 'Compton total' attenuation coefficient $\sigma_t/\rho = \sigma_a/\rho + \sigma_s/\rho$ is shown explicitly. (From Evans, R. D., *op. cit.*)




Mass attenuation coefficients for photons in xenon. (Adapted from Chupp, E. L., *Gamma-Ray Astronomy*, D. Reidel Publishing Co., Dordrecht, 1976, with permission.)




Mass attenuation coefficients (cont.)


Mass attenuation coefficient for photons in hydrogen. (From Chupp, E. L., *op. cit.*)




Mass attenuation coefficient for photons in polystyrene. (From Chupp, E. L., *op. cit.*)



Mass attenuation coefficient for photons in cesium iodide. (From Chupp, E. L., *op. cit.*)




Mass attenuation coefficient for photons in germanium. (From Chupp, E. L., *op. cit.*)



Fluorescence yields

Fluorescence yields for K and L shells for $5 \leq Z \leq 110$. The plotted curve for the L shell represents an average of L_1 , L_2 and L_3 effective yields. (From Kortright, J. B. in *X-ray Data Booklet*, Lawrence Berkeley Laboratory, University of California, 1986. Data from Krause, M. O., *J. Phys. Chem. Ref. Data*, **8**, 307, 1979.)



Passage of charged particles through matter

Total ionization loss by electrons or positrons (valid for all cases except extremely relativistic electrons)

$$-\left(\frac{dT}{ds}\right)_{\text{ion}} = \frac{2\pi e^4}{m_0 v^2} NZ \left\{ \ln \left[\frac{m_0 c^2 T}{I^2 (1 - \beta^2)} \right] - \beta^2 \right\} \text{ erg cm}^{-1}$$

(all quantities in cgs units).

For numerical calculations:

$$-\left(\frac{dT}{ds}\right)_{\text{ion}} = 4\pi r_0^2 \frac{m_0 c^2}{\beta^2} NZ \left[\ln \left[\beta \left(\frac{T + m_0 c^2}{I} \right) \left(\frac{T}{m_0 c^2} \right)^{1/2} \right] - \frac{1}{2} \beta^2 \right] \text{ MeV cm}^{-1},$$

where

$$\beta^2 = (v/c)^2 = 1 - [(T/m_0 c^2) + 1]^{-2},$$

v = velocity of particle,

$$m_0 c^2 = 0.51 \text{ MeV},$$

$$4\pi r_0^2 = 1.00 \times 10^{-24} \text{ cm}^2,$$

$r_0 = e^2/m_0 c^2$ = classical electron radius,

NZ = electrons cm^{-3} ,

Z = atomic number.

I = mean excitation potential (MeV).

$I \simeq (13 \times 10^{-6})Z$ (MeV).

T = kinetic energy (MeV).

N = atoms cm^{-3}

(e.g., 0.1 MeV electron: 4.7 keV per cm of air).

	Mean excitation potential I		I/Z
	Z	I (eV)	
H ₂	1	19	19
He	2	44	22
Be	4	64	16
Air	7.2	94	13.1
Al	13	166	12.7
Ar	18	230	12.8
Cu	29	371	12.8
Ag	47	586	12.5
Xe	54	660	12.2
Au	79	1017	12.8
Pb	82	1070	13.1

(List from Sternheimer, R. M., *Methods of Experimental Physics*, L. Marton, ed., Vol. 5, part A, Academic Press, New York, 1961.)

For $I \ll T \ll m_0c^2$:

$$-\left(\frac{dT}{ds}\right)_{\text{ion}} = \frac{2\pi e^4}{T} NZ \ln\left(\frac{T\sqrt{2}}{I}\right) \text{ erg cm}^{-1}.$$

In the range 0–10 MeV:

$$-\left(\frac{dT}{ds}\right)_{\text{ion}} \approx \frac{45}{\beta^2} \text{ ion pairs (air-cm)}^{-1}.$$

Radiative loss for non-relativistic electrons

$$-\left(\frac{dT}{ds}\right)_{\text{rad}} = 3.09 \times 10^{-27} NZ^2(T + m_0c^2) \text{ MeV cm}^{-1},$$

where

$T + m_0c^2$ = total energy of electron in MeV,

N = atoms $\text{cm}^{-3} = (\rho/A) \times 6.022 \times 10^{23}$,

Z = atomic number of absorber.

A = atomic weight of absorber,

ρ = density of absorber.

Ionization and radiative loss for highly relativistic electrons ($T \gg m_0c^2$)

$$-\left(\frac{dT}{ds}\right)_{\text{ion}} = \frac{2\pi e^4 NZ}{m_0 v^2} \left[\ln \frac{m_0 v^2 T}{2I^2(1-\beta^2)} - (2\sqrt{(1-\beta^2)} - 1 + \beta^2) \ln 2 + 1 - \beta^2 + \frac{1}{8}(1 - \sqrt{(1-\beta^2)})^2 \right] \text{ erg cm}^{-1}.$$

(Bethe, H. A., *Handbuch der Physik*, Vol. 24, p. 273, Julius Springer, Berlin, 1933.)

$$-\left(\frac{dE}{ds}\right)_{\text{rad}} = N \frac{Z^2}{137} r_0^2 E \left[4 \log \left(\frac{2E}{mc^2} \right) - \frac{4}{3} \right] \text{ erg cm}^{-1}$$

(for $mc^2 \ll E \ll 137mc^2 Z^{-1/3}$).

$$-\left(\frac{dE}{ds}\right)_{\text{rad}} = N \frac{Z^2 r_0^2}{137} E \left[4 \log(183Z^{-1/3}) + \frac{2}{9} \right] \text{ erg cm}^{-1}$$

(for $E \gg 137mc^2 Z^{-1/3}$).

$$E = T + m_0c^2$$

(Bethe, H. A. & Heitler, W., *Proc. Roy. Soc. (London)*, **A146**, 83, 1934.)

$$-\left(\frac{dT}{d\xi}\right)_{\text{rad}} = \frac{T}{X_0}, \quad T = T_0 e^{-\xi/X_0},$$

where

ξ is the distance travelled measured in g cm^{-2} , T_0 , the initial energy,

X_0 , the radiation length = $\frac{716M_A}{Z(Z+1.3)[\ln(183Z^{-1/3}) + \frac{1}{8}]} \text{ g cm}^{-2}$

(M_A = atomic weight).

Radiation lengths X_0 and critical energy T_c for various substances

Absorber	Z	M_A	X_0 (gm cm^{-2})	T_c (MeV)
Hydrogen	1	1	58	340
Helium	2	4	85	220
Carbon	6	12	42.5	103
Nitrogen	7	14	38	87
Oxygen	8	16	34.2	77
Aluminium	13	27	23.9	47
Argon	18	39.9	19.4	34.5
Iron	26	55.8	13.8	24
Copper	29	63.6	12.8	21.5
Lead	82	207.2	5.8	6.9
Air			36.5	83
Water			35.9	93

(List from Bethe, H. A. & Ashkin, J., *Experimental Nuclear Physics*, E. Segrè, ed., Vol. I, John Wiley and Sons, New York, 1953.)

$T_i \approx \frac{1600m_0c^2}{Z}$, the energy at which ionization losses equal radiation losses.

$$\frac{(dT/dx)_{\text{rad}}}{(dT/dx)_{\text{ion}}} \approx \frac{TZ}{1600mc^2}$$

Total ionization loss by heavy charged particles (kinetic energy \leq rest mass)

$$-\left(\frac{dT}{ds}\right)_{\text{ion}} = \frac{4\pi z^2 e^4}{m_0 V^2} NZ \left[\ln \frac{2m_0 V^2}{I} - \ln(1 - \beta^2) - \beta^2 \right] \text{ erg cm}^{-1}$$

(all quantities in cgs units).

For numerical calculations:

$$-\left(\frac{dT}{ds}\right)_{\text{ion}} = 4.58 \times 10^{-4} \frac{z^2 NZ}{V^2} \left[\ln \left(\frac{1.14 \times 10^{-15} V^2}{I} \right) - \ln(1 - \beta^2) - \beta^2 \right],$$

where

z = particle charge,

Z = absorber atomic number,

V = velocity of particle in $\text{cm s}^{-1} = 3 \times 10^{10} \beta$,

N = absorber atomic density = $\rho/A \times 6.022 \times 10^{23}$,

I = mean excitation potential (in eV),

$I \approx 13Z$ (eV),

$(1 - \beta^2) = (1 + T/M)^{-2}$,

T = kinetic energy in MeV,

M = mass of particle in MeV

(e.g., 2 MeV α particle in Si; $0.27 \text{ MeV } \mu\text{m}^{-1}$).

Approximate range-energy relationships

Range-energy for monoenergetic electrons

$20 \text{ eV} \leq E \leq 10 \text{ keV}$:

$$\ln((Z/A)R_{\text{ex}}) = -4.5467 + 0.31104 \ln E + 0.07773(\ln E)^2$$

where

R_{ex} = extrapolated range in $\mu\text{g cm}^{-2}$ (25% precision),

Z/A = charge to mass ratio for absorbing medium,

E = energy in eV.

(Iskef, H. *et al.*, *Phys. Med. Biol.*, **28**, 535, 1983.)

Approximate range-energy relationships (cont.)

10 keV $\lesssim E \lesssim$ 3 MeV:

$$R_{\text{ex}} (\text{mg cm}^{-2}) = 412E (\text{MeV})^n, \text{ where}$$

$$n = 1.265 - 0.0954 \ln E (\text{MeV}).$$

1 MeV $\lesssim E \lesssim$ 20 MeV:

$$R_{\text{ex}} (\text{mg cm}^{-2}) = 530E (\text{MeV}) - 106.$$

Continuous β -ray spectra

β -ray spectra are exponentially attenuated with a mass-absorption coefficient nearly independent of the absorbing material:

$$\mu/\rho (\text{cm}^2 \text{gm}^{-1}) = 17E_m^{-1.14},$$

where E_m (in MeV) is the maximum energy of the β -ray spectrum.

The thickness of absorber required to reduce the β -ray intensity to one-half its original value:

$$t_{1/2} (\text{mg cm}^{-2}) = 0.693/(\mu/\rho) = 41E_m^{1.14}.$$

Range-energy relationships for heavy particles

Alpha particles in air at 15 °C, 760 mm, 4–15 MeV,

$$R (\text{cm}) = (0.005E (\text{MeV}) + 0.285)E (\text{MeV})^{1.5}.$$

Protons in air at 15 °C, 760 mm, 10–200 MeV,

$$R (\text{cm}) = 100 \left(\frac{E (\text{MeV})}{9.3} \right)^{1.8}.$$

Range of heavy particles in other materials

Bragg-Kleeman rule:

$$\frac{R_1}{R_0} = \frac{\rho_0}{\rho_1} \sqrt{\left(\frac{A_1}{A_0} \right)} \quad (\pm 15\%),$$

where

ρ = density

A = atomic weight.

For mixtures:

$$\sqrt{A} = \sum_i n_i \sqrt{A_i},$$


where n_i = atomic fraction of element i .

For air, $\sqrt{A_0} = 3.81$, $\rho_0 = 1.226 \times 10^{-3} \text{ gm cm}^{-3}$ at 15 °C, 760 mm, and therefore:


$$R_1 = 3.2 \times 10^{-4} \frac{\sqrt{A_1}}{\rho_1} R_{\text{air}}.$$

Charged particles in silicon

Range energy curves for charged particles in silicon. Channeling of ions between crystal planes can result in significant variations from the data shown here. (Adapted from *ORTEC Manual on Surface Barrier Detectors*.)




Specific energy loss for charged particles in silicon. Channeling of ions between crystal planes can result in significant variations from the data shown here. (Adapted from *ORTEC Manual on Surface Barrier Detectors*.)




Charged particles in silicon (cont.)

Specific energy loss for electrons in silicon. Channeling of ions between crystal planes can result in significant variations from the data shown here. (Adapted from *ORTEC Manual on Surface Barrier Detectors*.)




Specific energy loss for protons in silicon. Channeling of ions between crystal planes can result in significant variations from the data shown here. (Adapted from *ORTEC Manual on Surface Barrier Detectors*.)



Charged particles in silicon (cont.)

Beta-ray range energy curve in silicon. Channeling of ions between crystal planes can result in significant variations from the data shown here. (Adapted from *ORTEC Manual on Surface Barrier Detectors*.)



Radioactive sources (Adapted from Lederer, C. M. *et al.*, *Table of Isotopes*, John Wiley & Sons, 1968.)

X-ray sources

Source	X-ray energy (keV)	Half-life
⁴⁹ V	Ti X-rays: 4.5, 4.9	330 d
⁵⁵ Fe	Mn X-rays: 5.9, 6.5	2.6 yr
⁵⁹ Ni	Co X-rays: 6.9, 7.7	8 × 10 ⁴ yr
¹⁰⁹ Cd	Ag X-rays: 22, 25	453 d
²⁰⁷ Pb	Pb X-rays: 73, 75, 85, 87	30 yr

Alpha sources

Source	Alpha energies (MeV)	Half-life
²⁴¹ Am	5.443, 5.486	458 yr
²⁵² Cf	6.117	2.65 yr
²⁴² Cm	6.017, 6.115	163 d
²⁴⁴ Cm	5.81, 5.77	17.6 yr
¹⁴⁸ Gd	3.18	84 yr
²³⁷ Np	4.78, 4.65	2.14 × 10 ⁶ yr
²⁰⁸ Po	5.11	2.93 yr
²¹⁰ Po	5.305	138 d
²³⁹ Pu	5.105, 5.143, 5.156	2.4 × 10 ⁴ yr
²²⁶ Ra	4.782, 5.490, 6.002, 7.687	1600 yr
²²⁸ Th	5.427 through 8.785	1.91 yr
²³⁰ Th	4.617, 4.684	8 × 10 ⁴ yr
²³⁵ U	4.336 through 4.597	7.1 × 10 ⁸ yr
²³⁸ U	4.200	4.5 × 10 ⁹ yr

Gamma-ray energy standards

Source	γ -ray energy (keV)	Half-life	Source	γ -ray energy (keV)	Half-life
⁵⁷ Co	14.359	268 d	⁵⁴ Mn	834.861	314 d
²⁴¹ Am	26.350	458 yr	⁴⁶ Sc	899.25	84.2 d
²⁴¹ Am	59.554	458 yr	⁸⁸ Y	898.033	108 d
²⁰³ Hg	70.830	47 d	²⁰⁷ Bi	1063.578	30 yr
²⁰³ Hg	72.871	47 d	⁶⁵ Zn	1115.522	246 d
¹³¹ I	80.164	8.05 d	⁴⁶ Sc	1120.50	84.2 d
²⁰³ Hg	82.572	47 d	⁶⁰ Co	1173.231	5.26 yr
²⁰³ Hg	84.916	47 d	²² Na	1274.552	2.58 yr
⁵⁷ Co	121.969	268 d	⁴¹ Ar	1293.641	1.85 hr
⁵⁷ Co	136.328	268 d	⁶⁰ Co	1332.518	5.26 yr
¹⁴¹ Ce	145.433	32.5 d	²⁴ Na	1368.526	15.0 hr
¹³⁹ Ce	165.85	140 d	⁵² V	1434.19	3.77 m:n
²⁰³ Hg	279.150	47 d	¹²⁴ Sb	1691.24	60.9 d
¹³¹ I	284.307	8.05 d	²⁸ Al	1778.77	2.31 m:n
⁵¹ Cr	320.102	27.8 d	⁸⁸ Y	1836.111	108 d
¹³¹ I	364.493	8.05 d	Th C''	2614.47	1.91 yr
¹⁹⁸ Au	411.795	2.70 d	²⁴ Na	2753.92	15.0 hr
⁷ Be	477.556	53 d	¹² B(β^-) ¹² C	4438.41	—
m_0c^2	511.003	—	¹⁴ C(d, p, β^+) ¹⁵ N	5298.53	—
⁸⁵ Sr	513.95	64 d	¹⁶ O(³ He, α) ¹⁵ O	5240.03	—
²⁰⁷ Bi	569.62	30 yr	¹⁴ N(d, p) ¹⁵ N	5270.10	—
Th C'	583.139	1.91 yr	¹⁶ O*	6127.8	—
¹³⁷ Cs	661.632	30 yr	¹³ C(p, γ) ¹⁴ N	9169.0	—
⁹⁵ Nb	765.83	35 d			

Electron energy standards

Source	Conversion-electron energy (keV)	Half-life
Cd ¹⁰⁹	62.19	453 d
Cd ¹⁰⁹	84.2	453 d
Ce ¹⁴¹	103.44	33 d
Ce ¹³⁹	126.91	140 d
Ce ¹⁴¹	138.63	33 d
Ce ¹³⁹	159.61	140 d
Hg ²⁰³	193.64	46.9 d
Hg ²⁰³	264.49	46.9 d
Au ¹⁹⁸	328.69	2.698 d
Sn ¹¹³	363.8	115 d
Sn ¹¹³	387.6	115 d
Au ¹⁹⁸	397.68	2.698 d
Bi ²⁰⁷	481.61	30 yr
Bi ²⁰⁷	554.37	30 yr
Cs ¹³⁷	624.15	30.0 yr
Cs ¹³⁷	655.88	30.0 yr
Co ⁵⁸	803.35	71.3 d
Co ⁵⁸	809.62	71.3 d
Mn ⁵⁴	828.86	303 d
Mn ⁵⁴	834.17	303 d
Y ⁸⁸	881.86	108 d
Y ⁸⁸	895.76	108 d
Bi ²⁰⁷	975.57	30 yr
Bi ²⁰⁷	1048.1	30 yr
Zn ⁶⁵	1106.46	345 d
Zn ⁶⁵	1114.35	245 d

Characteristics of synchrotron radiation

The angular and spectral distribution of radiation emitted by a relativistic electron in instantaneously circular motion is, according to Schwinger (*Phys. Rev.*, **75**, 1912, 1949), given by (cgs units):

$$\frac{\partial^2 P}{\partial \lambda \partial \psi} = \frac{27}{32\pi^3} \frac{e^2 c}{R^3} \left(\frac{\lambda_c}{\lambda}\right)^4 \gamma^8 (1 + (\gamma\psi)^2)^2 \left[K_{2,3}^2(\xi) + \frac{(\gamma\psi)^2}{1 + (\gamma\psi)^2} K_{1,3}^2(\xi) \right] \text{ erg s}^{-1} \text{ rad}^{-1} \text{ cm}^{-1},$$

where

e = electron charge,

c = velocity of light,

R = radius of curvature,

λ = wavelength,


$\lambda_c = \frac{4}{3}\pi R\gamma^{-3}$, the 'critical wavelength',

$\gamma = E/m_0c^2$, where E = electron energy and m_0 = electron mass,

ψ = vertical angle measured from the plane of the orbit,

$K_{1,2}(\xi)$, $K_{2,3}(\xi)$ are modified Bessel functions of the second kind, where $\xi \equiv (\lambda_c/2\lambda)(1 + (\gamma\psi)^2)^{3/2}$.

The first term within the brackets corresponds to radiation polarized in the plane of the orbit, the second to radiation polarized perpendicular to the orbital plane.



The universal synchrotron radiation function $G(y)$ for monoenergetic electrons as a function of $1/y$.

The spectral distribution of power is obtained by integrating the above equation over the angle ψ :

$$\frac{dP}{d\lambda} = \frac{3^5 2}{16\pi^2} \frac{e^2 c}{R^3} \left(\frac{E}{mc^2} \right)^7 G(y),$$

where

$$G(y) = y^3 \int_y^\infty K_{5/3}(\eta) d\eta, \quad y = \lambda_c / \lambda.$$

The total power radiated,

$$P = \int \frac{dP}{d\lambda} d\lambda = \frac{2}{3} \frac{e^2 c}{R^2} \left(\frac{E}{mc^2} \right)^4.$$


The photon flux distribution is obtained by dividing the power distribution function by the photon energy:

$$\frac{\partial^3 N}{\partial \lambda \partial \psi \partial t} = \frac{\lambda}{hc} \frac{\partial^2 P}{\partial \lambda \partial \psi} \quad \text{photons s}^{-1} \text{ rad}^{-1} \text{ cm}^{-1}.$$


The radiation emitted by the electron in the plane of its orbit is 100% polarized. Above and below this plane, the radiation is elliptically polarized.

Polarization:

$$(I_{\parallel} - I_{\perp}) / (I_{\parallel} + I_{\perp}) = \frac{K_{2/3}^2(\xi) - [(\gamma\psi)^2 / (1 + (\gamma\psi)^2)] K_{1/3}^2(\xi)}{K_{2/3}^2(\xi) + [(\gamma\psi)^2 / (1 + (\gamma\psi)^2)] K_{1/3}^2(\xi)}$$



Fraction of radiation, integrated over vertical angle ψ , that is parallel polarized (From Krinsky, S. *et al.* in *Handbook of Synchrotron Radiation*, E. Koch, ed., North-Holland Publishing Co., 1983, with permission.)



Dependence on the vertical angle ψ of the intensities of the parallel (solid line) and perpendicular (dashed line) polarization components of the photon flux. The individual curves, plotted for $\lambda/\lambda_c = \frac{1}{3}$, 1, 10, and 100, are individually normalized to the intensity in the orbital plane ($\psi = 0$) at the respective λ/λ_c value. Note that the abscissa, ψ multiplied by the electron energy γ , makes these curves universal. (From Krinsky, S. *et al.* in *Handbook of Synchrotron Radiation*, E. Koch, ed., North-Holland Publishing Co., 1983, with permission.)

Practical formulae for electron storage rings

E : electron energy R : radius of ring J : electron current.

Energy loss per turn, per electron:

$$\delta E \text{ (keV)} = 88.5 \frac{E^4 \text{ (GeV)}}{R \text{ (m)}}$$

Critical wavelength:

$$\lambda_c \text{ (\AA)} = 5.59 \frac{R \text{ (m)}}{E^3 \text{ (GeV)}}$$

Characteristic energy:

$$\epsilon_c \text{ (eV)} = 2218 \frac{E^3 \text{ (GeV)}}{R \text{ (m)}} = 2.96 \times 10^{-7} \frac{\gamma^3}{R \text{ (m)}}$$

Emission angle:

$$\approx \frac{1}{\gamma} = \frac{m_0 c^2}{E}$$

with

$$\gamma = \frac{E}{m_0 c^2} = 1957 E \text{ (GeV)}$$

Energy of a photon:

$$E(\text{KeV}) = \frac{12.40}{\lambda(\text{\AA})}$$

Photon flux angular distribution:

$$\frac{\partial^4 N}{\partial \lambda \partial \psi \partial \theta \partial t} = 8.267 \times 10^{-5} \left(\frac{\lambda_c}{\lambda} \right)^{3\gamma^2} \frac{1}{R} F \left(\frac{\lambda_c}{\lambda}, \gamma \psi \right) J$$

photons s⁻¹ mrad⁻¹ mrad⁻¹ Å⁻¹,

where

$$F \left(\frac{\lambda_c}{\lambda}, \gamma \psi \right) = (1 + \gamma^2 \psi^2)^2 \left[K_{2,3}^2(\xi) + \frac{\gamma^2 \psi^2}{1 + \gamma^2 \psi^2} K_{1,3}^2(\xi) \right],$$

$$\xi = \frac{1}{2} \frac{\lambda_c}{\lambda} (1 + \gamma^2 \psi^2)^{3/2},$$

where

θ = horizontal angle,

ψ = vertical angle,

J = current (mA),

R = radius of ring (m).

Photon flux integrated over all vertical angles

$$\frac{\partial^3 N}{\partial \lambda \partial \theta \partial t} = 7.9 \times 10^{11} G(\gamma) J(\text{mA}) \frac{[E(\text{GeV})]^7}{[R(\text{m})]^2} \lambda(\text{\AA})$$

photons s⁻¹ mrad⁻¹ Å⁻¹,

$$\frac{\partial^3 N}{\partial E \partial \theta \partial t} = 5.56 \times 10^7 G(\gamma) J(\text{mA}) \frac{[E(\text{GeV})]^7}{[R(\text{m})]^2} \lambda^3(\text{\AA})$$

photons s⁻¹ mrad⁻¹ eV⁻¹,

for $\lambda \gg \lambda_c$


$$\frac{\partial^3 N}{\partial \lambda \partial \theta \partial t} = 9.35 \times 10^{13} J(\text{mA}) \frac{[R(\text{m})]^{1/3}}{[\lambda(\text{\AA})]^{4/3}} \text{photons s}^{-1} \text{mrad}^{-1} \text{\AA}^{-1}.$$

V. Kostroun (*Nuc. Inst. Meth.*, **172**, 371, 1980) provides series expressions for the modified Bessel functions of fractional order which are suitable for evaluation with programmable calculators or desktop computers.


X-ray spectroscopy

Crystal spectroscopy

Collimated single crystal Bragg spectrometer. Bragg condition: $n\lambda = 2d \sin \theta$, where d is the effective spacing of the crystal planes that participate in the reflection. (Adapted from Burek, A., *Space Sci. Inst.*, 2, 53, 1976.)



Single crystal rocking curve C_c , $\theta_B + \Delta = \sin^{-1}(n\lambda/2d)$, where Δ = refraction correction, d = physical spacing of reflecting planes, and θ_B = Bragg angle ignoring refraction. (Adapted from Burek, A., *Space Sci. Inst.*, 2, 53, 1976.)



Crystal properties

Crystal	Density (g cm ⁻³)	Plane	2d (Å)	Integrated reflectivity (θ = 60°)
Quartz	2.66	1010	8.350	6.25 × 10 ⁻⁵ (D)
		1011	6.592	1.23 × 10 ⁻⁴ (D)
		202 $\bar{3}$	2.750	≈ 1.5 × 10 ⁻⁵ (D)
		2243	2.028	≈ 6 × 10 ⁻⁶ (D)
Topaz	3.49-3.57	303	2.712	(40) 6 × 10 ⁻⁵ (D)
		040	4.40	
		400	2.3246	(43) 1 × 10 ⁻⁵ (D)
		200	4.64	
Calcite	2.710	211	6.083	1.62 × 10 ⁻⁴ (D)
Silicon	2.33	111	6.284	1.2 × 10 ⁻⁴ (D)
		220	3.840	8 × 10 ⁻⁵ (D)
		200	5.44169	
Germanium	5.33	111	6.545	
		220	4.000	2.3 × 10 ⁻⁴ (D)
		200	5.66897	
Beryl (golden)	2.66	1010	15.9549	≈ 6 × 10 ⁻⁵ (D)
Sylvite	1.99	200	6.292	
Halite	2.164	200	5.641	
KBr	2.756	200	6.584	
Fluorite	3.18	111	6.306	
		200	5.4744	
Aluminum	2.699	200	4.057	
		111	4.676	
LiF	2.64	420	1.80	
		200	4.027	10 ⁻⁴ - 3 × 10 ⁻⁴ (D)
		220	2.848	
Graphite	2.21	002	6.708	1.52 × 10 ⁻³ (S)
Mica	2.77-2.88	002	19.84	≈ 2 × 10 ⁻⁵ (S)
Clinochlore	2.6-3.3	001	28.392	
ADP	1.803	101	10.648	9.0 × 10 ⁻⁵ (S)
		220	5.305	1.4 × 10 ⁻⁵ (S)
		200	7.50	
EDDT	1.538	020	8.808	1.15 × 10 ⁻⁴ (S)
PET	1.39	002	8.742	2.2 × 10 ⁻⁴ (S)
SHA	1.3	110	13.98	
KAP	1.636	001	26.5790	5 × 10 ⁻⁵ (S)
RAP	1.94	001	26.121	1.5 × 10 ⁻⁴ (S)
TIAP	2.7	001	25.7567	7.0 × 10 ⁻⁴ (S)
CsAP	2.178	001	25.68	
NH ₄ AP	1.415	002	26.14	1.5 × 10 ⁻⁴ (S)
NaAP	1.504	002	26.42	

D = double crystal; S = single crystal.

(Adapted from Burek, A., *Space Sci. Inst.*, 2, 53, 1976.)

Useful characteristic lines for X-ray spectroscopy

Wavelength (Å)	Energy (keV)	Element	Designation
1.54	8.04	Cu	$K\alpha_{1,2}$
1.66	7.47	Ni	$K\alpha_{1,2}$
1.94	6.40	Fe	$K\alpha_{1,2}$
2.29	5.41	Cr	$K\alpha_{1,2}$
2.75	4.51	Ti	$K\alpha_{1,2}$
3.60	3.44	Sn	$L\alpha_{1,2}$
4.15	2.98	Ag	$L\alpha_{1,2}$
5.41	2.29	Mo	$L\alpha_{1,2}$
6.86	1.80	Sr	$L\alpha_{1,2}$
7.13	1.74	Si	$K\alpha_{1,2}$
8.34	1.49	Al	$K\alpha_{1,2}$
8.99	1.38	Se	$L\alpha_{1,2}$
9.89	1.25	Mg	$K\alpha_{1,2}$
10.44	1.19	Ge	$L\alpha_{1,2}$
12.25	1.01	Zn	$L\alpha_{1,2}$
13.34	0.930	Cu	$L\alpha_{1,2}$
14.56	0.852	Ni	$L\alpha_{1,2}$
15.97	0.776	Co	$L\alpha_{1,2}$
17.59	0.705	Fe	$L\alpha_{1,2}$
18.32	0.677	F	$K\alpha$
19.45	0.637	Mn	$L\alpha_{1,2}$
21.64	0.573	Cr	$L\alpha_{1,2}$
23.62	0.525	O	$K\alpha$
27.42	0.452	Ti	$L\alpha_{1,2}$
31.36	0.395	Ti	$L\beta$
31.60	0.392	N	$K\alpha$
44.7	0.277	C	$K\alpha$
58.4	0.212	W	$N_V N_{VII}$
64.38	0.193	Mo	$M\zeta$
67.6	0.183	B	$K\alpha$
82.1	0.151	Zr	$M\zeta$
114	0.109	Be	$K\alpha$

Grating spectroscopy

Concave grating equation:

$$\pm m\lambda = d(\sin \alpha + \sin \beta),$$

where m is the spectral order, d is the groove separation, α is the angle of incidence, and β is the angle of diffraction. The negative sign applies when the spectrum lies between the central image ($\alpha = \beta$) and the tangent to the grating (sometimes referred to as the 'outside order'). When the spectrum lies between the incident beam and the central image, the positive sign must be used, and the spectrum is referred to as the 'inside order'. The signs of α and β are opposite when they lie on different sides of the grating normal.


Angular dispersion (α fixed):

$$\frac{d\beta}{d\lambda} = \frac{m}{d \cos \beta}$$


Plate factor:

$$\frac{d\lambda}{dl} = \frac{d \cos \beta}{mR}, \quad \frac{d\lambda}{dl} = \frac{\cos \beta}{mR(1/d)} \times 10^4 \text{ \AA mm}^{-1},$$

where R is in meters, $1/d$ is the number of lines mm^{-1} , and l is the distance along the Rowland circle.




Optical layout of basic spectrograph.



Grazing incidence spectrograph.

(Adapted from Samson, J., *Techniques of Vacuum Ultraviolet Spectroscopy*, John Wiley and Sons, 1967.)

Reflection of X-rays



In the X-ray band the complex refractive index n^* is usually expressed as:

$$n^* = (1 - \delta) - i\beta,$$

with

$$\delta = \frac{1}{2\pi} \left(\frac{e^2}{m_e c^2} \right) \left(\frac{N_0 \rho}{A} \right) f_1 \lambda^2$$

and

$$\beta = \mu_a \lambda = \frac{1}{2\pi} \left(\frac{e^2}{m_e c^2} \right) \left(\frac{N_0 \rho}{A} \right) f_2 \lambda^2,$$

where

$$\frac{e^2}{m_e c^2} = r_e = \text{the classical electron radius,}$$

N_0 = Avogadro's number,

ρ = density,

A = atomic weight,

λ = wavelength of the incident radiation,

f_1 = real part of the atomic scattering factor, and

f_2 = imaginary part of the atomic scattering factor.

$$f_1 = (\pi r_e h c)^{-1} \int_0^x \frac{E'^2 \mu_a(E') dE'}{E^2 - E'^2} + Z,$$

$$f_2 = \frac{\pi}{2} (\pi r_e h c)^{-1} E \mu_a,$$

where

Z = atomic number,

h = Planck's constant,

c = velocity of light,

E = incident photon energy, and

μ_a = atomic photoabsorption cross-section.

The atomic photoabsorption cross-section μ_a is related to the mass absorption coefficient (μ/ρ) or to the linear absorption coefficient μ by:

$$\mu_a = \left(\frac{A}{N_0} \right) \left(\frac{\mu}{\rho} \right) = \left(\frac{A}{N_0 \rho} \right) \mu.$$

Numerically,

$$\delta = 2.701 \times 10^{-6} \frac{\rho(\text{g cm}^{-3})}{A} \lambda^2(\text{\AA}) f_1,$$

$$\beta = 2.701 \times 10^{-6} \frac{\rho(\text{g cm}^{-3})}{A} \lambda^2(\text{\AA}) f_2 = \frac{\lambda \mu}{4\pi} \left(\frac{\mu}{\rho} \right).$$

The reflection of X-rays by a perfectly smooth surface for an angle of incidence θ_i is given by the Fresnel equations:

$$R_s = \frac{a^2 + b^2 - 2a \cos \theta_i + \cos^2 \theta_i}{a^2 + b^2 + 2a \cos \theta_i + \cos^2 \theta_i}$$

for perpendicular polarization, and

$$R_p = R_s \frac{a^2 + b^2 - 2a \sin \theta_i \tan \theta_i + \sin^2 \theta_i \tan^2 \theta_i}{a^2 + b^2 + 2a \sin \theta_i \tan \theta_i + \sin^2 \theta_i \tan^2 \theta_i}$$

for parallel polarization, where

$$2a^2 = [(n^2 - \beta^2 - \sin^2 \theta_i)^2 + 4n^2 \beta^2]^{1/2} + (n^2 - \beta^2 - \sin^2 \theta_i),$$

and

$$2b^2 = [(n^2 - \beta^2 - \sin^2 \theta_i)^2 + 4n^2 \beta^2]^{1/2} - (n^2 - \beta^2 - \sin^2 \theta_i),$$

with $n = 1 - \delta$.

Since the real part of the index of refraction is less than 1, near total external reflection occurs at a grazing angle ϕ_c given by Snell's law:

$$\cos \phi_c = 1 - \delta,$$


$$\phi_c \approx \sqrt{2\delta} \quad \text{for } \delta \ll 1.$$

Since $\beta \neq 0$, reflection is not total for $\phi < \phi_c$ but is less than 1. Away from an absorption edge,

$$\delta \approx 2.70 \times 10^{-6} \left(\frac{Z_c}{A} \right) \rho \lambda^2,$$

where Z_c is the number of electrons associated with wavelengths greater than λ . $Z_c = Z$ for $\lambda < \lambda_k$ (K edge).

Calculated specular reflectivity of an ideal surface as a function of normalized grazing angle ϕ/ϕ_c for various values of β/δ . (After Hendrick, *JOSA*, 47, 165, 1957.)



Optical constants at soft X-ray wavelengths

λ (Å)	$\delta \times 10^4$	$\beta \times 10^4$
NICKEL		
1.25†	0.17	0.02
2.0†	0.42	0.01
5.0†	2.85	0.4
8.34	5.91 ± 0.17	1.9 ± 0.2
9.89	8.00 ± 0.20	3.2 ± 0.3
10.44	8.70 ± 0.20	4.4 ± 0.4
12.25	8.85 ± 0.21	5.3 ± 0.5
13.34	10.20 ± 0.22	7.2 ± 0.7
14.56	1.37 ± 0.08	9.6 ± 1.0
15.97	12.30 ± 0.20	2.1 ± 0.2
17.59	17.50 ± 0.30	2.8 ± 0.3
19.4	21.70 ± 0.30	3.7 ± 0.4
21.64	30.50 ± 0.40	6.1 ± 0.6
23.62	33.30 ± 0.40	7.3 ± 0.7
24.78	39.40 ± 0.44	9.1 ± 0.9
27.42	47.30 ± 0.48	10.0 ± 1.0
31.36	60.00 ± 0.50	15.0 ± 1.5
41.3†	120.0	51.3
47.7†	148.6	76.5
GOLD		
1.25†	0.28	0.02
2.0†	0.77	0.1
5.0†	2.94	2.4
8.34	9.15 ± 0.21	3.1 ± 0.3
9.89	11.50 ± 0.24	4.4 ± 0.4
12.254	18.00 ± 0.30	9.4 ± 0.9
13.34	21.20 ± 0.30	12.3 ± 1.2
14.56	24.10 ± 0.30	15.7 ± 1.6
15.97	26.60 ± 0.40	18.6 ± 1.9
17.59	29.20 ± 0.40	23.4 ± 2.3
19.45	33.50 ± 0.40	26.8 ± 2.7
21.64	34.80 ± 0.40	28.5 ± 2.9
23.62	37.60 ± 0.40	37.6 ± 3.8
24.78	40.00 ± 0.40	56.0 ± 5.6
27.42	40.80 ± 0.50	61.0 ± 6.0
31.36	47.50 ± 0.50	61.5 ± 6.2
44.7	58.00 ± 0.30	58.0 ± 5.9
67.6	136.00 ± 1.0	102.0 ± 10.5



† δ and β are calculated for these wavelengths.



$n^* = (1 - \delta) - i\beta$, complex refractive index.



(Adapted from Ershov, O. A., *Optical Spectroscopy*, **22**, 66, 1967; Lukirskii, A. P. et al., *Optical Spectroscopy*, **16**, 168, 1964.)



Reflectivity vs. wavelength (energy) for various grazing angles and materials



(Courtesy of M. Hettrick, Lawrence Berkeley Laboratory, Berkeley, CA.)
Whenever possible, direct measurements should be made of grazing incidence reflectivity in the X-ray region because of uncertainties in the optical constants.













Reflectivity versus wavelength for various materials and grazing angles. (Adapted from Giacconi, R. *et al.*, *Space Science Review*, 9, 3, 1969.)



Wolter type I mirror system



The equations for a paraboloid and hyperboloid which are concentric and confocal can be written as:

$$r_p^2 = P^2 + 2PZ + [4e^2Pd/(e^2 - 1)] \quad (\text{paraboloid}),$$

$$r_h^2 = e^2(d + Z)^2 - Z^2 \quad (\text{hyperboloid}).$$

The origin is at the focus for axial rays, Z is the coordinate along the axis of symmetry, and r is the radius of the surface at Z .

RMS blur circle radius:

$$\sigma = \frac{(\xi + 1) \tan^2 \theta}{10 \tan \alpha} \left(\frac{L_p}{Z_0} \right) + 4 \tan \theta \tan^2 \alpha \quad \text{radians}$$

and

$$\xi = \alpha_p^* / \alpha_h^*$$

(α_p^* and α_h^* are the grazing angles between the two surfaces and the path of an axial ray that strikes at an infinitesimal distance from the intersection).

For most telescope designs: $\xi = 1$.

$$\alpha = \frac{1}{4} \tan^{-1}(r_0/Z_0) = \frac{1}{2}(\alpha_p^* + \alpha_h^*),$$

θ = angle between incident rays and optical axis.

Geometrical collecting area:

$$A \approx 2\pi r_0 L_p \tan \alpha.$$

Effective collecting area:


$$A_e(\alpha, E) \approx AR^2(\alpha, E) \approx 8\pi Z_0 L_p R^2(\alpha, E) \alpha^2,$$

where R is the Fresnel reflectivity at energy E and mean grazing angle α .

(Adapted from Van Speybroeck, L. & Chase, R., *Ap. Opt.* **11**, 440, 1972.)

Vacuum technology

Vacuum nomograph. (Adapted from Roth, A., *Vacuum Technology*, North-Holland Pub. Co., 1976.)



Pumping speed of an aperture of area A :

$$\frac{dV}{dt} = A(\text{cm}^2) \sqrt{[1.32 \times 10^7 T(\text{K})/\text{mol. wt.}] \text{ cm}^3 \text{ s}^{-1}}.$$

Kinetic theory of gases

Mean free path, $\lambda = 1/\sqrt{2} \pi n \sigma^2$

viscosity, $\eta = \rho \bar{v} \lambda / 3$

heat conductivity, $K = \eta c_v$,

mean speed, $\bar{v} = \sqrt{[2.1 \times 10^8 T(\text{K})/\text{mol. wt.}] \text{ cm s}^{-1}}$,

where

n = number of molecules cm^{-3} ,

ρ = gas density in g cm^{-3} ,

σ = mol. diameter,

c_v = specific heat capacity at constant volume,

$\varepsilon = 2.5$ and 1.9 for monatomic and diatomic gas, respectively.

Equivalents for various cryogenic fluids

Fluid	Boiling point at 1 atm	Weight in pounds	Ft ³ at 70 F and 1 atm	Liquid liters at b.p.	Liquid gallons at b.p.	Heat of vapor. (Btu)
Nitrogen	-320.4 F	1	13.81	0.5618	0.1484	85.2
	-195.8 C	0.0724	1	0.0407	0.0108	6.168
	77.3 K	1.780	24.58	1	0.2642	151.7
		6.738	92.94	3.785	1	574.1
Helium	-452.1 F	1	96.8	3.628	0.9585	8.8
	-268.9 C	0.0103	1	0.375	0.0099	0.0906
	4.2 K	0.2756	26.68	1	0.2642	2.425
		1.043	101.0	3.785	1	9.178
Oxygen	-297.4 F	1	12.09	0.3973	0.1050	91.7
	-183.0 C	0.827	1	0.0329	0.0087	7.584
	90.1 K	2.517	30.43	1	0.2642	230.8
		9.527	115.2	3.785	1	873.6
Hydrogen	-423.2 F	1	192.3	6.481	1.712	193
	-252.9 C	0.0052	1	0.0337	0.0089	1.004
	20.2 K	0.1543	29.67	1	0.2642	29.78
		0.5841	112.3	3.785	1	112.7
Argon	-302.6 F	1	9.680	0.3235	0.0855	70.2
	-185.9 C	0.1033	1	0.0334	0.0088	7.251
	87.2 K	3.091	29.92	1	0.2642	217.0
		11.70	113.3	3.785	1	821.3

Optical point spread function

The irradiance distribution of the monochromatic image of a point object, $h_\lambda(x, y; \alpha, \beta)$, is called the point spread function (PSF) of an optical system. x and y are the coordinates of the image points and α and β are the coordinates of the ideal image of the object (a point). If $f_\lambda(x, y)$ is the ideal image of an extended monochromatic object, the image produced by the optical system is given by:

$$g_\lambda(x, y) = \iint_{-\infty}^{\infty} h_\lambda(x, y; \alpha, \beta) f_\lambda(\alpha, \beta) d\alpha d\beta.$$

In some cases, the optical system is *shift-invariant* (at least, over a restricted field):

$$h_\lambda(x, y; \alpha, \beta) = K_\lambda(x - \alpha, y - \beta).$$

and the above integral can be written as a convolution:

$$g_{\lambda}(x, y) = \iint_{-r}^{+r} K_{\lambda}(x_{\lambda} - \alpha, y - \beta) f_{\lambda}(x, \beta) d\alpha d\beta.$$

In general, for optical systems, the PSF is wavelength dependent, *shift-varying*, and asymmetric. Over a restricted field, say within a few arc minutes of the optical axis, it is approximately shift-invariant and symmetric and it is possible to simplify the deconvolution of an image and use Fourier transforms.

In the following discussion we will consider the PSF to be shift invariant and symmetric and will drop the λ subscript.

The PSF is the function that completely characterizes the imaging properties of an optical system. Several useful functions and quantities can be derived from it:

The line spread function

$$A(x) = \int_{-r}^x K(x, y) dy \quad (K(x, y) \text{ is the PSF})$$

represents the intensity distribution for a line object.

The edge trace

$$I(x_0) = \int_{-r}^{x_0} A(x) dx$$

represents the intensity distribution for a knife edge object.

The modulation transfer function (MTF)

This represents the response of an optical system to an object with a sinusoidally varying radiance G of spatial frequency ν :

$$\text{MTF}(\nu) = \frac{M_i(\nu)}{M_o(\nu)},$$

where M_i and M_o are the modulation of the image and object, respectively.

The modulation is given by:

$$M_{i \text{ or } o}(\nu) = \left(\frac{\max - \min}{\max + \min} \right) \text{ image or object.}$$

Since the radiance of the object (image) varies sinusoidally, we can write:

$$G_o(x) = a_o + b_o \sin 2\pi\nu x,$$

$$G_i(x) = a_i + b_i \sin 2\pi\nu x.$$

Then

$$M_o = \frac{(a_o + b_o) - (a_o - b_o)}{(a_o + b_o) + (a_o - b_o)} = \frac{b_o}{a_o},$$

$$M_1 = \frac{(a_1 + b_1) - (a_1 - b_1)}{(a_1 + b_1) + (a_1 - b_1)} = \frac{b_1}{a_1},$$

and the modulation transfer function,

$$\text{MTF} = \frac{(b_1/a_1)}{(b_0/a_0)}.$$

The MTF is also given by the absolute value of the Fourier transform of the line spread function:

$$\text{MTF}(v) = \int_{-x}^x A(x) e^{-2\pi i v x} dx.$$

The **full-width-half-maximum (FWHM)** of a rotationally symmetric PSF is the width of the function at half its peak value. If $K(x, y) = K(r)$, where r is the radial coordinate in the image plane, and the radius r_0 is such that:

$$K(r_0) = \frac{1}{2}K(0) \quad (\text{half the peak value of the PSF}),$$

then

$$\text{FWHM} = 2r_0; \quad r_0 = \text{half-width},$$

or half-width-half-maximum (HWHM).

The **root mean square (rms) radius** $\langle r \rangle$ is defined by:

$$\langle r \rangle^2 = \frac{\int_0^x r^2 K(r) r dr}{\int_0^x K(r) r dr}.$$

The **encircled energy function** $E(r)$, the fraction of the total imaged photons that are within a circle of radius r , is given by:

$$E(r) = \frac{\int_0^r K(r) r dr}{\int_0^\infty K(r) r dr}.$$

The radius of the circle which contains 50% of the imaged photons, the **half-power radius**, $r_{1/2}$ is defined by:

$$E(r_{1/2}) \equiv 0.50 = \frac{\int_0^{r_{1/2}} K(r) r dr}{\int_0^\infty K(r) r dr}.$$

In order to complete the discussion of the point spread function, we give here the various functions and parameters derived from a Gaussian point spread since this can be a useful description of the inner core of the PSF.

The form of the radial symmetric Gaussian is given as:

$$\text{PSF} = C e^{-r^2/2\sigma^2},$$

where C and σ are arbitrary constants. We have derived the following functions and quantities:

the line spread function,

$$A(x) = C\sigma\sqrt{2\pi}e^{-x^2/2\sigma^2},$$

the edge trace,

$$I(x_0) = C\sigma\sqrt{\frac{2\pi}{2}}(1 + \text{erf}(x_0/\sigma\sqrt{2}))$$

where $\text{erf}(z)$ is the error function.

the modulation transfer function,

$$M(v) = C e^{-2(\pi v \sigma)^2}, \text{ and}$$

the encircled energy,

$$E(r) = 1 - e^{-r^2/2\sigma^2}.$$

The table below gives the relations between the FWHM, the rms radius, and the half-power radius for the Gaussian spread function.

PARAMETERS FOR THE GAUSSIAN SPREAD FUNCTION

Spread function	rms radius $\langle r \rangle$	Full width half maximum (FWHM)	Half power radius $r_{1/2}$
$C e^{-r^2/2\sigma^2}$	$\sigma\sqrt{2}$	2.36σ	1.18σ

Point spread function for a circular aperture (diffraction by a circular aperture)

If the transmission of the system is uniform over the (circular) aperture and the system is aberration-free, the illuminance distribution in the image becomes:

$$P(y, z) = \pi \left(\frac{\text{NA}}{\lambda} \right)^2 P_t \left[\frac{2J_1(m)}{m} \right]^2,$$

where NA is the numerical aperture of the system, J_1 is the first-order Bessel function:

$$J_1(x) = \frac{x}{2} - \frac{(x/2)^3}{1^2 \cdot 2} + \frac{(x/2)^5}{1^2 \cdot 2^2 \cdot 3} - \dots,$$

P_t is the total power in the point image, and m is the normalized radial

Point spread function for a circular aperture (diffraction by a circular aperture) (cont.)



coordinate:

$$m = \frac{2\pi}{\lambda} \text{NA}(y^2 + z^2)^{1/2} = \frac{2\pi}{\lambda} \text{NA} \cdot s.$$

The fraction of the total power falling within a radial distance s_0 of the center of the pattern is given by $1 - J_0^2(m_0) - J_1^2(m_0)$, where J_0 is the zero-order Bessel function:


$$J_0(x) = 1 - \left(\frac{x}{2}\right)^2 + \frac{(x/2)^4}{1^2 2^2} - \frac{(x/2)^6}{1^2 2^2 3^2} + \dots$$

Fraunhofer diffraction at a rectangular aperture (a) and at a circular aperture (b). (Adapted from Born, M. & Wolf, E., *Principles of Optics*, Pergamon Press, 1984.)







Optical telescopes

Configurations of optical telescopes. (Adapted from Sinnott, R. W., *Sky and Telescope*, July 1980.)



Optical telescopes (cont.)

Focus configurations for optical telescopes. (Adapted from *Survey of Catadioptric Optical Systems*, J. B. Galligan, ed., Itek Corporation, 1966.)



Photometry

Table 1.10.10.10.1

Spectral luminous efficiency. Relative luminosity values for photopic and scotopic vision

Wavelength (nm)	Photopic $V(\lambda)$ ($B > 3 \text{ cd m}^{-2}$)	Scotopic $V(\lambda)$ ($B < 3 \times 10^{-5} \text{ cd m}^{-2}$)
350		0.0003
360		0.0008
370		0.0022
380	0.000 04	0.0055
390	0.000 12	0.0127
400	0.0004	0.0270
410	0.0012	0.0530
420	0.0040	0.0950
430	0.0116	0.157
440	0.023	0.239
450	0.038	0.339
460	0.060	0.456
470	0.091	0.576
480	0.139	0.713
490	0.208	0.842
500	0.323	0.948
510	0.503	0.999
520	0.710	0.953
530	0.862	0.849
540	0.954	0.697
550	0.995	0.531
560	0.995	0.365
570	0.952	0.243
580	0.870	0.155
590	0.757	0.0942
600	0.631	0.0561
610	0.503	0.0324
620	0.381	0.0188
630	0.265	0.0105
640	0.175	0.0058
650	0.107	0.0032
660	0.061	0.0017
670	0.032	0.0009
680	0.017	0.0005
690	0.0082	0.0002
700	0.0041	0.0001
710	0.0021	—
720	0.001 05	—
730	0.000 52	—
740	0.000 25	—
750	0.000 12	—
760	0.000 06	—
770	0.000 03	—


$K(\lambda)$, spectral luminous efficacy for
scotopic and photopic vision.

Scotopic:

$$\max K = K(511 \text{ nm}) = 1746 \text{ lm W}^{-1}.$$

Photopic:

$$\max K = K(555 \text{ nm}) = 680 \text{ lm W}^{-1}.$$



Summary of typical sources/parameters for the most commonly used radiant energy sources

Lamp type	DC input power (watts)	Arc dimensions (mm)	Luminous flux (lm)†	Luminous efficiency (lm W ⁻¹)	Average luminance (cd mm ⁻²)
Mercury short arc (high pressure)	200	2.5 × 1.8	9500	47.5	250
Xenon short arc	150	1.3 × 1.0	3200	21	300
Xenon short arc	20 000	12.5 × 6	1 150 000	57	3000 (in 3 mm × 6 mm)
Zirconium arc	100	1.5 (diam.)	250	2.5	100
Vortex-stabilized argon arc	24 800	3 × 10	422 000	17	1400
Tungsten light bulbs	{ 10 100 1000	{ — — —	{ 79 1630 21 500	{ 7.9 16.3 21.5	{ 10 to 25
Fluorescent lamp standard warm white	40	—	2560	64	—
Carbon arc, non-rotating	2000	≈ 5 × 5	36 800	18.4	175 to
rotating	15 800	≈ 8 × 8	350 000	22.2	800
Deuterium lamp	40	1.0 (diam.)	(Nominal irradiance at 250 nm at 30 cm = 0.2 μW cm ⁻² nm ⁻¹)		

† Luminous flux Φ in lumens from a source of total radiant power $W(\lambda)$ watts per unit wavelength:

$$\Phi = 680 \int_0^x W(\lambda) V(\lambda) d\lambda,$$

where $V(\lambda)$ represents the spectral luminous efficiency.

Conversion table for various photometric units

Luminous intensity (I)

1 candela (cd) = 1 lumen/steradian (lm sr⁻¹)

Luminous flux (Φ) [lumen (lm)]

4π lumens = total flux from uniform point source of 1 candela

Illuminance (E)

1 footcandle (fc) = 1 lumen foot⁻²

1 lux (lx) = 1 lumen m⁻² = 0.0929 footcandle


*Luminance (L)*1 footlambert (fL) = $1/\pi$ candela foot⁻²,1 nit (nt) = 1 candela m⁻² = 0.2919 footlambert*Luminance values for various sources*

Source	Luminance (fL)	Luminance (cd m ⁻²)
Sun, as observed from Earth's surface at meridian	4.7×10^8	1.6×10^9
Moon, bright spot, as observed from Earth's surface	730	2500
Clear blue sky	2300	7900
Lightning flash	2×10^{10}	7×10^{10}
Atomic fission bomb, 0.1 ms after firing, 90-ft diameter ball	6×10^{11}	2×10^{12}
Tungsten filament lamp, gas-filled, 16lm W ⁻¹	2.6×10^6	9×10^6
Plain carbon arc, positive crater	4.7×10^6	1.6×10^7
Fluorescent lamp, T-12 bulb, cool white, 430 mA, medium loading	2000	7000
Color television screen, average brightness	50	170

Typical values of natural scene illuminance

Sky condition	Approximate levels of illuminance (lux)
Direct sunlight	$1-1.3 \times 10^5$
Full daylight (not direct sunlight)	$1-2 \times 10^4$
Overcast day	10^3
Very dark day	10^2
Twilight	10
Deep twilight	1
Full moon	10^{-1}
Quarter moon	10^{-2}
Moonless, clear night sky	10^{-3}
Moonless, overcast night sky	10^{-4}

Natural illuminance on the Earth for the hours immediately before and after sunset with a clear sky and no moon



Radiant responsivity

Calculation of radiant responsivity from luminous responsivity for photocathodes:

The response of a photocathode (in amperes) to the total radiation $W(\lambda)$ watts per unit wavelength is:

$$\int \sigma R(\lambda) W(\lambda) d\lambda,$$

where the relative spectral response of the photocathode is $R(\lambda)$ ($R_{\max} = 1$) and σ is the absolute radiant response at the peak of the response curve (amperes per watt). The light flux (in lumens) is given by:

$$680 \int V(\lambda) W(\lambda) d\lambda,$$

where $V(\lambda)$ is the spectral luminous efficiency. The luminous responsivity of the photocathode in amperes per lumen is then given by:

$$S = \frac{\sigma \int R(\lambda) W(\lambda) d\lambda}{680 \int V(\lambda) W(\lambda) d\lambda}$$

and, therefore,

$$\sigma = \frac{680S \int V(\lambda) W(\lambda) d\lambda}{\int R(\lambda) W(\lambda) d\lambda}.$$

(The material in the preceding sections was adapted from Engstrom, R. W., *Photomultiplier Handbook*, RCA Corporation, 1980.)

Quantity ^(a)	Symbol ^(a)	Defining equation ^(b)	Commonly used units ^(c)	Symbol
RADIOMETRIC				
Radiant energy	Q (Q_e)		erg joule kilowatt-hour joule per cubic meter erg per cubic centimeter erg per second twatt	J kWh J m ³ erg cm ⁻³ erg s ⁻¹ W
Radiant density	w (w_e)	$w = dQ/dV$		
Radiant flux	Φ (Φ_e)	$\Phi = dQ/dt$		
Radiant flux density at a surface				
Radiant exitance (radiant emittance) ^(d)	M (M_e)	$M = d\Phi/dA$	watt per square centimeter	W cm ⁻²
Irradiance	E (E_e)	$E = d\Phi/dA$	twatt per square meter, etc.	W m ⁻²
Radiant intensity	I (I_e)	$I = d\Phi/d\omega$ ^(h)	twatt per steradian	W sr ⁻¹
Radiance	L (L_e)	$L = d^2\Phi/d\omega(dA \cos \theta)$ $= dI/(dA \cos \theta)$ ^(e)	watt per steradian and square centimeter twatt per steradian and square meter	W sr ⁻¹ cm ⁻² W sr ⁻¹ m ⁻²
Emissivity	ϵ	$\epsilon = M/M_{\text{blackbody}}$ ^(f)	one (numeric)	—
PHOTOMETRIC				
Absorptance	α (α_v, α_e)	$\alpha = \Phi_a/\Phi_i$ ^(g)	one (numeric)	
Reflectance	ρ (ρ_v, ρ_e)	$\rho = \Phi_r/\Phi_i$ ^(g)	one (numeric)	
Transmittance	τ (τ_v, τ_e)	$\tau = \Phi_t/\Phi_i$ ^(g)	one (numeric)	
Luminous energy (quantity of light)	Q (Q_v)	$Q_v = \int_{380}^{760} K(\lambda)Q_e \lambda d\lambda$	lumen-hour †lumen-second (talbot)	lm-h lm-s
Luminous density	w (w_v)	$w = dQ/dV$	†lumen-second per cubic meter	lm-s m ⁻³
Luminous flux	Φ (Φ_v)	$\Phi = dQ/dt$	†lumen	lm

Standard units, symbols, and defining equations for fundamental photometric and radiometric quantities (cont.)

Quantity ^(a)	Symbol ^(a)	Defining equation ^(b)	Commonly used units ^(c)	Symbol
Luminous flux density at a surface			lumen per square foot	lm ft ⁻²
Luminous exitance (luminous emittance) ^(d)	M (M_v)	$M = d\Phi/dA$	footcandle (lumen per square foot)	fc
Illumination (illuminance)	E (E_v)	$E = d\Phi/dA$	flux (lm m ⁻²)	lx
Luminous intensity (candlepower)	I (I_v)	$I = d\Phi/d\omega$ ^(e)	phot (lm cm ⁻²)	ph
Luminance (photometric brightness)	L (L_v)	$L = d^2\Phi/d\omega dA \cos \theta$ $= dI/(dA \cos \theta)$ ^(e)	teandela (lumen per steradian)	cd
			candela per unit area	cd m ⁻² , etc.
			stilb (cd cm ⁻²)	sb
			nit (tcd m ⁻²)	nt
			footlambert (cd per π ft ²)	fl.
			lambert (cd per π cm ²)	L
Luminous efficacy	K	$K = \Phi_e/\Phi_v$ ^(h)	apostilb (cd per π m ²)	asb
Luminous efficiency	V	$V = K/K_{\max}$ ^(h)	flumen watt ⁻¹	lm W ⁻¹
			one (numeric)	

^(a) The symbols for photometric quantities are the same as those for the corresponding radiometric quantities. When it is necessary to differentiate them, the subscripts v and e , respectively, should be used, e.g., Q_v and Q_e . Quantities may be restricted to a narrow wavelength band by adding the word spectral and indicating the wavelength. The corresponding symbols are changed by adding a subscript λ , e.g., Q_λ for a spectral concentration or a λ in parentheses, e.g., $K(\lambda)$, for a function of wavelength.

^(b) The equations in this column are given merely for identification.

^(c) International System (SI) unit indicated by dagger (†).

^(d) To be deprecated.

^(e) θ is the angle between line of sight and normal to surface considered.

^(f) M and $M_{\text{blackbody}}$ are, respectively, radiant exitance of a measured specimen and of a blackbody at the same temperature as the specimen.

^(g) Φ_i is incident flux, Φ_a is absorbed flux, Φ_r is reflected flux, Φ_t is transmitted flux.

^(h) K_{\max} is the maximum value of the $K(\lambda)$ function.

⁽ⁱ⁾ ω is the solid angle through which flux from point source is radiated.


(*Photoelectronic Imaging Devices*, Vol. 1, L. M. Biberman & S. Nudelman, eds., Plenum Press, 1971, with permission.)

Detectors


Visible and ultraviolet light detectors

Photodiode

Schematics of photodiodes (a) sealed with semi-transparent photocathode, (b) open (or sealed) with opaque photocathode. (From Timothy, J. G. & Madden, R. P. in *Handbook on Synchrotron Radiation*, E. Koch, ed., North-Holland Publishing Co., 1983, with permission.)



Quantum efficiencies of opaque Cs_2Te and CsI photocathodes. (From Timothy, J. G. & Madden, R. P., *op. cit.*)



Quantum efficiencies of transfer standard detectors available from NBS. (From Timothy, J. G. & Madden, R. P., *op. cit.*)






Image intensifiers

Generation I electrostatically focused image intensifier. (Reproduced with permission of the publisher, Howard W. Sams & Co., Indianapolis, *Image Tubes*, by Illes P. Csorba, © 1985.)




The electrostatic image-inverting generation II image intensifier employs a microchannel plate (MCP). (From Csorba, I. P., *op. cit.*)




Photomultipliers

Schematic of typical photomultiplier showing some electron trajectories (RCA).




Typical anode sensitivity and amplification characteristics of a 9-dynode photomultiplier tube as a function of applied voltage (RCA).



- 12 DYNODES 14, FOCUSING ELECTRODES
13, ANODE 15 PHOTOCATHODE

Typical voltage-divider arrangement for fast pulse response and high peak current systems.



Source and detector matching

The average power radiating from a light source is given by:

$$P = P_0 \int_0^{\infty} W(\lambda) d\lambda,$$

where P_0 is the incident power in watts per unit wavelength at the peak of the relative spectral radiation characteristic, $W(\lambda)$, which is normalized to unity.

The resulting photocathode current I_k , when the light is incident on the detector, is given by:

$$I_k = \sigma P_0 \int_0^{\infty} W(\lambda) R(\lambda) d\lambda,$$

where σ is the radiant sensitivity of the photocathode in amperes per watt at the peak of the relative curve, and $R(\lambda)$ represents the relative photocathode spectral response as a function of wavelength normalized to unity at the peak.

$$I_k = \sigma P \frac{\int_0^{\infty} W(\lambda) R(\lambda) d\lambda}{\int_0^{\infty} W(\lambda) d\lambda}$$

The ratio of the dimensionless integrals can be defined as the *matching factor*, M .

$$M = \frac{\int_0^{\infty} W(\lambda) R(\lambda) d\lambda}{\int_0^{\infty} W(\lambda) d\lambda}$$

Spectral matching factors

Source	Photocathodes										Photopic eye	Scotopic eye
	S1	S4	S10	S11	S17	S20	S25					
PHOSPHORS												
P1	0.278	0.498	0.807	0.687	0.892	0.700	0.853	0.768	0.743			
P4	0.310	0.549	0.767	0.661	0.734	0.724	0.861	0.402	0.452			
P7	0.312	0.611	0.805	0.709	0.773	0.771	0.882	0.411	0.388			
P11	0.217	0.816	0.949	0.914	0.954	0.877	0.953	0.201	0.601			
P15	0.385	0.701	0.855	0.787	0.871	0.802	0.904	0.376	0.495			
P16	0.830	0.970	0.853	0.880	0.855	0.902	0.922	0.003	0.042			
P20	0.395	0.284	0.612	0.427	0.563	0.583	0.782	0.707	0.354			
P22B	0.217	0.893	0.974	0.960	0.948	0.927	0.979	0.808	0.477			
P22G	0.278	0.495	0.807	0.686	0.896	0.699	0.855	0.784	0.747			
P22R	0.632	0.036	0.264	0.055	0.077	0.368	0.623	0.225	0.008			
P24	0.279	0.545	0.806	0.696	0.827	0.725	0.869	0.540	0.621			
P31	0.276	0.533	0.811	0.698	0.853	0.722	0.868	0.626	0.651			
NaI	0.534	0.923	0.885	0.889	0.889	0.900	0.933	0.046	0.224			

Spectral matching factors (cont'd)

Source	Photocathodes										Photopic eye	Scotopic eye
	S1	S4	S10	S11	S17	S20	S25					
LAMPS												
2870/2856 std	0.516†	0.046†	0.095†	0.060†	0.072†	0.112†	0.227†	0.071†	0.040†			
Fluorescent	0.395	0.390	0.650	0.496	0.575	0.635	0.805	0.502	0.314			
SUN												
In space	0.535†	0.308†	0.388†	0.328†	0.380†	0.406†	0.547†	0.179†	0.172†			
+ 2 air masses	0.536†	0.236†	0.348†	0.277†	0.315†	0.360†	0.513†	0.197†	0.175†			
Day sky	0.537†	0.520†	0.556†	0.508†	0.589†	0.581†	0.700†	0.170†	0.218†			
BLACKBODIES												
6000 K	0.533†	0.308†	0.376†	0.320†	0.375†	0.397†	0.521†	0.167†	0.159†			
3000 K	0.512†	0.053†	0.102†	0.067†	0.080†	0.120†	0.232†	0.075†	0.044†			
2870 K	0.504†	0.044†	0.090†	0.057†	0.069†	0.106†	0.216†	0.067†	0.038†			
2856 K	0.500†	0.042†	0.088†	0.055†	0.068†	0.103†	0.211†	0.065†	0.037†			
2810 K	0.493†	0.039†	0.081†	0.051†	0.062†	0.097†	0.150†	0.061†	0.034†			
2042 K	0.401†	0.008†	0.023†	0.011†	0.014†	0.033†	0.090†	0.018†	0.007†			

† Entry valid only for 300–1200 nm wavelength interval.

Nominal composition and characteristics of various photocathodes


Nominal composition	JETEC response designation	Conversion factor ^(a) (lumen watt ⁻¹ at λ_{max})	Luminous responsibility ($\mu\text{A lumen}^{-1}$)	Wavelength of maximum response, λ_{max} (nm)	Responsivity at λ_{max} (mA watt ⁻¹)	Quantum efficiency at λ_{max} (%)	Dark emission at 25°C (fA cm ⁻²)
Ag-O-Cs	S-1	92.7	25	800	2.3	0.36	900
Ag-O-Rb	S-3	285	6.5	420	1.8	0.55	
Cs ₃ Sb	S-19	1603	40	330	64	24	0.3
Cs ₃ Sb	S-4	1044	40	400	42	13	0.2
Cs ₃ Sb	S-5	1262	40	340	50	18	0.3
Cs ₃ Bi	S-8	757	3	365	2.3	0.77	0.13
Ag-Bi-O-Cs	S-10	509	40	450	20	5.6	70
Cs ₃ Sb	S-13	799	60	440	48	14	4
Cs ₃ Sb	S-9	683	30	480	20	5.3	
Cs ₃ Sb	S-11	808	60	440	48	14	3
Cs ₃ Sb	S-21	783	30	440	23	6.7	
Cs ₃ Sb	S-17	667	125	490	83	21	1.2
Na ₂ KSb	S-24	758	85	420	64	19	0.0003

Nominal composition and characteristics of various photocathodes (cont'd)

Nominal composition	JETEC response designation	Conversion factor ^(a) (lumen watt ⁻¹ at λ_{max})	Luminous responsibility ($\mu\text{A lumen}^{-1}$)	Wavelength of maximum response, λ_{max} (nm)	Responsivity at λ_{max} (mA watt ⁻¹)	Quantum efficiency at λ_{max} (%)	Dark emission at 25°C (fA cm ⁻²)
K ₂ CsSb		1117	85	400	95	29	0.02
Rb-Cs-Sb		767	120	450	92	25	1
Na ₂ KSb:Cs		429	150	420	64	19	0.4
Na ₂ KSb:Cs	S-20	428	150	420	64	19	0.3
Na ₂ KSb:Cs	S-25	276	160	420	44	13	—
Na ₂ KSb:Cs	ERMA II	220	200	530	44	10.3	2.1
Na ₂ KSb:Cs	ERMA III	160	230	575	37	8	0.2
GaAs:Cs-O		116	1025	850	119	17	92
GaAsP:Cs-O		310	200	450	61	17	0.01
In _{0.06} Ga _{0.94} As:Cs-O		200	250	400	50	15.5	220
In _{0.12} Ga _{0.88} As:Cs-O		255	270	400	69	21	40
In _{0.18} Ga _{0.82} As:Cs-O		280	150	400	42	13	75
Cs ₂ Te				250	25	12.4	0.0006
Cs I				120	24	20	Very low
Cu I				150	13	10.7	Very low
K Cs-Rb-Sb		672	125	440	84	24	Very low

^(a) These conversion factors are the ratio of the radiant responsibility at the peak of the spectral response characteristic in amperes per watt to the luminous responsibility in amperes per lumen for a tungsten lamp operated at a color temperature of 2856 K.

Typical photocathode spectral response characteristics (from RCA Corp.)



Short wavelength transmission limits of some UV window materials

Material	Approximate limit (10 ^u , 2 mm thick)
LiF	1040 Å
MgF ₂	1120
CaF ₂	1220
SrF ₂	1280
BaF ₂	1340
Al ₂ O ₃ (sapphire)	1410
SiO ₂ (fused quartz)	1600

UV fluorescent converters (wavelength shifters)

Sodium salicylate	Diphenylstilbene
Tetraphenyl butadiene	<i>p</i> -Terphenyl
Coronene	Dimethyl POPOP
<i>p</i> -Quaterphenyl	POPOP

X-ray and gamma-ray detectors*Detection principles - quantum efficiency*

In general, the quantum efficiency, $\varepsilon(E)$, for an incident photon of energy E is determined by the transmission of the detector window or any 'dead layer' and by the absorption of the detector medium:

$$\varepsilon(E) = e^{-(\mu/\rho)_w \rho_w t_w} (1 - e^{-(\mu/\rho)_d \rho_d t_d}),$$

where $(\mu/\rho)_w$ and $(\mu/\rho)_d$ are the mass absorption coefficients of the detector window (or 'dead layer') and detector medium, respectively, ρ_w and ρ_d are the densities of the detector window (or 'dead layer') and detector medium, respectively, and t_w and t_d are the thicknesses of the detector window (or 'dead layer') and detector medium, respectively.

Detection principles - point source detection with X-ray telescopes

The fluctuation δN_S in the number of counts from a point source of flux density F photons $\text{cm}^{-2} \text{s}^{-1} \text{keV}^{-1}$ is given by:

$$\delta N_S = (A_{\text{eff}} \Delta E F t + f^2 \omega B_i \Delta E t + A_{\text{eff}} \omega j_D \Delta E t)^{1/2},$$

where

A_{eff} = effective area (cm^2) of telescope including detector,

ΔE = energy interval (keV),

t = observing time (s),

f = focal length (cm) of telescope,

ω = solid angle (sr) of picture element,

B_i = internal background ($\text{ct cm}^{-2} \text{s}^{-1} \text{keV}^{-1}$) of detector,

j_D = diffuse X-ray background ($\text{photons cm}^{-2} \text{s}^{-1} \text{keV}^{-1} \text{sr}^{-1}$).




The background, both internal and from diffuse X-rays, is assumed to be steady and well known. For a strong source, the signal-to-noise ratio $N_S/\delta N_S = (A_{\text{eff}} \Delta E F t)^{1/2}$ is given by the fluctuations in the source only.

For a weak source, fluctuations in the background determine the signal-to-noise ratio:

$$\frac{N_S}{\delta N_S} = \frac{(A_{\text{eff}} \Delta E F t)^{1/2}}{\omega^{1/2} (f^2 B_i / A_{\text{eff}} + j_D)^{1/2}}.$$


Scintillation detector

Illustration representing a NaI scintillation detector showing sequence of events producing output from electron multiplier and various processes which contribute to response of detector to a gamma-ray source. (Adapted from Heath, R. L., *Scintillation Spectrometry*, USAEC Report, IDO-16880, 1964.)



A typical pulse-height spectrum obtained with a NaI(Tl) spectrometer, illustrating the energy response of inorganic scintillators. The scale of the abscissa is 1 keV per channel.

Gas proportional counter



Since a proportional counter has internal gain, the system noise can be neglected and the energy resolution is:

$$(\Delta E)_{\text{FWHM}} = 2.35[(F + f)WE]^{1/2} \text{ eV},$$

where

E = energy deposited in counter (eV),

F = Fano factor,

f = a factor to account for variance in the gas gain,

W = mean energy to form an ion pair (eV).

As an example, for methane gas:

$$F = 0.26$$


$$f = 0.75$$

$$W = 27 \text{ eV},$$

so that for a proportional counter:


$$\frac{E}{(\Delta E)_{\text{FWHM}}} = 2.6E^{1/2} \quad (\text{with } E \text{ in keV}).$$

Total number of ion pairs collected in a gas-filled chamber as a function of the voltage across electrodes of the chamber.



Position sensitive gas proportional detector

Readout system of detector. Incident photon is absorbed at point a ; electrons drift toward anode-cathode planes. An avalanche at the anode (A) gives rise to pulse distributions at the cathodes (K_{\parallel} and K_{\perp}). The position (X, Y) is obtained by analog summation and division. (Adapted from Bade, E. *et al.*, *Nucl. Inst. and Meth.*, **201**, 193, 1982.)



Typical performance

Spatial resolution:

0.25 mm (FWHM) at 1 keV.

Energy resolution:

$$\frac{E}{(\Delta E)_{\text{WHM}}} = 2.2E^{1.2} \quad (\text{with } E \text{ in keV}).$$

Format:

10 cm \times 10 cm.

The solid-state detector

$$(\Delta E)_{\text{FWHM}} = 2.35[(\eta\sigma)^2 + (F\eta E)]^{1/2} \quad \text{eV},$$

where


η = conversion factor (Si: 3.6 eV per electron-hole pair; Ge: 2.9 eV per electron-hole pair),

σ = detector rms noise (electrons),

F = Fano factor (Si: 0.14; Ge: 0.13),


E = photon or particle energy (eV).

Schematic diagram of a solid state detector. (Adapted from Enge, H., *Introduction to Nuclear Physics*, Addison-Wesley, 1966.)



Charge-coupled device (CCD)

(Front-side illuminated)



↑
X-rays

(Back-side illuminated)

$$(\Delta E)_{\text{FWHM}} = 2.35 [(\eta\sigma)^2 + (\eta i_d t)^2 + \eta FE]^{1/2} \text{ keV},$$

where

i_d = dark current (electrons s^{-1}),

σ = rms readout noise (electrons),


η = mean energy required to produce one electron hole pair
(0.0036 keV for silicon),

t = integration time (s),

F = Fano factor (~ 0.15),

E = energy of incident photon (keV).

Expected quantum efficiency (defined as the probability that an incident X-ray photon is detected as an 'event') vs. energy. The calculations consider only the interactions of X-rays in Si. for two hypothetical CCD's whose dead-layer and substrate thicknesses are separately within the range spanned by real devices. There will be a low energy cutoff (not shown) depending on the minimum signal which can be discriminated against the system noise.



Microchannel plate detector

Typical performance

Spatial resolution:

20–30 μm (FWHM).


Quantum efficiency:

25% at 1.5 keV (CsI photocathode).

Format:

25–100 mm in diameter.

Schematic diagram of a microchannel plate detector.
(Adapted from Behr, A. in Landolt-Bornstein, subvol. 2a, Springer-Verlag, 1981.)



Properties of common X-ray detectors

Detector	Energy range (keV)	$\Delta E/E^{(a)}$ at 5.9 keV (%)	Dead time/event (μs)	Maximum count rate (s^{-1})
Geiger counter	3-50	none	200	10^4
Gas ionization in current mode	0.2-50	n/a	n/a	10^{11} ^(b)
Gas proportional	0.2-50	15	0.2	10^5
Multiwire proportional chamber	3-50	20	0.2	10^5 /anode wire
Scintillation [NaI(Tl)]	3-10 000	40	0.25	10^6
Semiconductor [Si(Li)]	1-60	3.0	4-30	5×10^4
Semiconductor (Ge)	1-10 000	3.0	4-40	5×10^4

^(a) FWHM.

^(b) Maximum count rate density is limited by space-charge effects to around 10^{11} photons $\text{s}^{-1} \text{cm}^{-3}$.

(From Thompson, A. C. in *X-ray Data Booklet*, Lawrence Berkeley Laboratory, University of California, 1986.)

Ionization and excitation data for a number of gases

Gas	Atomic number	First ionization potential (eV)	Second ionization potential (eV)	First excited state (eV)	Principal emission wavelengths (Å)
He	2	24.48	54.40	20.9 19.8 meta	584 3888 5875
Ne	10	21.56	41.07	16.68 16.53 meta 16.62 meta	734 743 5400 5832 5852 6402
Ar	18	15.76	27.62	11.56 11.49 meta 11.66 meta	1048 1066 6965 7067 7503 8115
Kr	36	14.00	24.56	9.98 9.86 meta 10.51 meta	1236 5570 5870
Xe	54	12.13	21.2	8.39 8.28 meta 9.4 meta	1296 1470 4501 4624 4671
H	1	13.60		10.2	1215 4861 6562
N	7	14.53	29.59	6.3	1200 4110
O	8	13.61	35.11	9.1	1302 7771
H ₂		15.4		11.2	
N ₂		15.8		6.1	
O ₂		12.5			
I ₂		9.0		1.9	1782 2062

(Adapted from Rice-Evans, P., *Spark, Streamer, Proportional and Drift Chambers*, The Richelieu Press, London, 1974.)

Properties of scintillation and solid-state detector materials

Material	Density (g cm^{-3})	Band gap (eV)	λ of max. emission (\AA)	Decay time ^(a) (μs)	Index of refrac- tion ^(m)	Energy ^(c) (eV)	K-edge (keV)	Scintillation conversion ^(d) efficiency (%)	Notes
SCINTILLATORS									
NaI(Tl)	3.67	5.38	4100	0.23	1.85	—	1.07, 33.2	100	Hygroscopic
CaF ₂ (Eu)	3.18	—	4350	0.94	1.47	—	0.68, 4.04	50	Non-hygroscopic
CsI(Na)	4.51	5.67	4200	0.63	1.84	—	33.2, 36.0	80	Hygroscopic
CsI(Tl)	4.51	5.67	5650	1.0	1.80	—	33.2, 36.0	45	Non-hygroscopic
Plastics	1.06	—	3500-4500	0.002-0.020	Varies	—	0.284	20-30	Non-hygroscopic
Liquids	0.86	—	3500-4500	0.002-0.008	Varies	—	0.284	20-30	Non-hygroscopic
SOLID-STATE									
Si(Li)	2.35	1.21	—	—	—	3.6	1.84	—	LN ₂ required during operation
Ge(Li)	5.36	0.785	—	—	—	2.9	11.1	—	LN ₂ required during operation

^(a) Room temperature, exponential decay constant.

^(b) At emission maximum.

^(c) Per electron-hole pair.

^(d) Referred to NaI(Tl) with S-11 photocathode.

(Adapted from *Harshaw Scintillation Phosphors*, The Harshaw Chemical Company.)

Properties of materials used in X-ray detector systems

	Atomic number, Z	Density (STP) (g cm ⁻³)	Shell energy ^(a) (keV)	X-ray lines ^(b) (keV)	Fluorescence yield ^(c)	Energy at which photoelectric equals Compton cross-section (keV)
PROPORTIONAL COUNTER GASES						
Methane (CH ₄)	6	0.713 × 10 ⁻³	0.284	0.277		20
Ne	10	0.901 × 10 ⁻³	0.867	0.849, 0.858	0.01	38
Ar	18	1.78 × 10 ⁻³	3.203 (0.285, 0.246, 0.244)	2.96	0.105	72
Kr	36	3.74 × 10 ⁻³	14.32 (1.92, 1.73, 1.67)	12.64, 14.12	0.625 (0.04)	170
Xe	54	5.85 × 10 ⁻³	34.56 (5.45, 5.10, 4.78)	29.67, 33.78 (4.10, 4.49, 5.30)	0.875 (0.14)	300
CRYSTALS						
NaI	53	3.61	33.16 (5.19, 4.86, 4.56)	28.47, 32.30 (3.93, 4.22, 4.80)	0.865 (0.13)	260
CsI	55	4.54	35.97	30.81, 34.99 (4.28, 4.62, 5.28)	0.885 (0.15)	300
Si	14	2.33	1.84	1.74, 1.83	0.04	53
Ge	32	5.36	11.10 (1.42, 1.41, 1.21)	9.88, 10.98 (1.19, 1.22)	0.49 (0.01)	145

Properties of materials used in X-ray detector systems (cont'd)

	Atomic number, Z	Density (STP) (g cm ⁻³)	Shell energy ^(a) (keV)	X-ray lines ^(b) (keV)	Fluorescence yield ^(c)	Energy at which photoelectric equals Compton cross-section (keV)
WINDOWS, FILTERS						
Be	4	1.82	0.111	0.109		14
Mylar	8	1.4	0.532	0.525		
(C ₁₀ H ₈ O ₄) _n	6		0.284	0.277		
Formvar	8	1.2 ^(d)	0.532	0.525		
(C ₅ H ₇ O ₂) _n	6		0.284	0.277		
Polypropylene						
(CH ₂) _n	6	0.95 ^(d)	0.284	0.277		22
Air (1.3% Ar, 23.2% O, 75.5% N)	18	1.29 × 10 ⁻³	3.20	2.96	0.105	27
	8		0.532	0.525		
	7		0.400	0.392		
Mg	12	1.74	1.30	1.25, 1.30	0.02	46
Al	13	2.7	1.56	1.49, 1.55	0.03	50
Fe	26	7.87	7.11	6.40, 7.06	0.31	135
			(0.849, 0.722, 0.709)			
Ni	28	8.9	8.33	7.47, 8.27	0.38	140
			(1.01, 0.877, 0.858)			
Cu	29	8.96	8.99	8.04, 8.91	0.40	145
			(1.10, 0.954, 0.935)			

^(a) The first line is the K-shell energy. The three L-shell energies are listed in the parentheses.


^(b) The first line for each material gives K α and K β energies. L α , L β , and L γ are listed in the parentheses.

^(c) The first line is the K fluorescence yield. The L fluorescence yield is listed in the parentheses. If the fluorescence yield is less than 1%, the space is left blank.

^(d) Variable, depending on the detailed preparation of a batch of film.


Compton telescope for high energy gamma-rays

a) Configuration of a Compton telescope which relies on the detection of scattered photons: S, scatterer; C, collector; A_1 , A_2 , anticoincidence detectors. b) Basic configuration of a Compton telescope which relies on the detection of secondary electrons. (Adapted from Hillier, R., *Gamma Ray Astronomy*, Clarendon Press, 1984.)



Spark chamber telescope for high energy gamma rays

Diagram showing the basic design of a spark chamber with plates (S), anticoincidence shield (A), and triggering detectors (C_1 and C_2). (Adapted from Hillier, R., *Gamma Ray Astronomy*, Clarendon Press, 1984.)



Microcomputer topics***ASCII character code***

This chart shows the ASCII character set and corresponding code numbers, in decimal, hexadecimal and binary form.

Decimal	Binary	Hexadecimal	ASCII character
0	00000000	00	(NUL)
1	00000001	01	(SOH)
2	00000010	02	(STX)
3	00000011	03	(ETX)
4	00000100	04	(EOT)
5	00000101	05	(ENQ)
6	00000110	06	(ACK)
7	00000111	07	(BEL)
8	00001000	08	(BS)
9	00001001	09	(HT)
10	00001010	0A	(LF)
11	00001011	0B	(VT)
12	00001100	0C	(FF)
13	00001101	0D	(CR)
14	00001110	0E	(SO)
15	00001111	0F	(SI)
16	00010000	10	(DLE)
17	00010001	11	(DC1)
18	00010010	12	(DC2)
19	00010011	13	(DC3)
20	00010100	14	(DC4)
21	00010101	15	(NAK)
22	00010110	16	(SYN)
23	00010111	17	(ETB)
24	00011000	18	(CAN)
25	00011001	19	(EM)
26	00011010	1A	(SUB)
27	00011011	1B	(ESC)
28	00011100	1C	(FS)
29	00011101	1D	(GS)
30	00011110	1E	(RS)
31	00011111	1F	(US)
32	00100000	20	(SP)
33	00100001	21	!
34	00100010	22	"
35	00100011	23	#
36	00100100	24	\$
37	00100101	25	%
38	00100110	26	&
39	00100111	27	'
40	00101000	28	(
41	00101001	29)
42	00101010	2A	*

ASCII character code (cont.)

Decimal	Binary	Hexadecimal	ASCII character
43	00101011	2B	+
44	00101100	2C	,
45	00101101	2D	-
46	00101110	2E	.
47	00101111	2F	
48	00110000	30	0
49	00110001	31	1
50	00110010	32	2
51	00110011	33	3
52	00110100	34	4
53	00110101	35	5
54	00110110	36	6
55	00110111	37	7
56	00111000	38	8
57	00111001	39	9
58	00111010	3A	:
59	00111011	3B	;
60	00111100	3C	<
61	00111101	3D	=
62	00111110	3E	>
63	00111111	3F	?
64	01000000	40	@
65	01000001	41	A
66	01000010	42	B
67	01000011	43	C
68	01000100	44	D
69	01000101	45	E
70	01000110	46	F
71	01000111	47	G
72	01001000	48	H
73	01001001	49	I
74	01001010	4A	J
75	01001011	4B	K
76	01001100	4C	L
77	01001101	4D	M
78	01001110	4E	N
79	01001111	4F	O
80	01010000	50	P
81	01010001	51	Q
82	01010010	52	R
83	01010011	53	S
84	01010100	54	T
85	01010101	55	U
86	01010110	56	V
87	01010111	57	W
88	01011000	58	X
89	01011001	59	Y
90	01011010	5A	Z

ASCII character code (cont.)

Decimal	Binary	Hexadecimal	ASCII character
91	01011011	5B	[
92	01011100	5C	\
93	01011101	5D]
94	01011110	5E	^
95	01011111	5F	_
96	01100000	60	`
97	01100001	61	a
98	01100010	62	b
99	01100011	63	c
100	01100100	64	d
101	01100101	65	e
102	01100110	66	f
103	01100111	67	g
104	01101000	68	h
105	01101001	69	i
106	01101010	6A	j
107	01101011	6B	k
108	01101100	6C	l
109	01101101	6D	m
110	01101110	6E	n
111	01101111	6F	o
112	01110000	70	p
113	01110001	71	q
114	01110010	72	r
115	01110011	73	s
116	01110100	74	t
117	01110101	75	u
118	01110110	76	v
119	01110111	77	w
120	01111000	78	x
121	01111001	79	y
122	01111010	7A	z
123	01111011	7B	{
124	01111100	7C	
125	01111101	7D	}
126	01111110	7E	~
127	01111111	7F	(DEL)
128	10000000	80	
129	10000001	81	
130	10000010	82	
131	10000011	83	
132	10000100	84	
133	10000101	85	
134	10000110	86	
135	10000111	87	
136	10001000	88	
137	10001001	89	
138	10001010	8A	

ASCII character code (cont.)

Decimal	Binary	Hexadecimal	ASCII character
139	10001011	8B	
140	10001100	8C	
141	10001101	8D	
142	10001110	8E	
143	10001111	8F	
144	10010000	90	
145	10010001	91	
146	10010010	92	
147	10010011	93	
148	10010100	94	
149	10010101	95	
150	10010110	96	
151	10010111	97	
152	10011000	98	
153	10011001	99	
154	10011010	9A	
155	10011011	9B	
156	10011100	9C	
157	10011101	9D	

Boolean algebra

$A = \bar{\bar{A}}$	
$AA = A$	
$A + A = A$	
$A \times 0 = 0$	
$A + 0 = A$	
$A \times 1 = A$	
$A + 1 = 1$	
$A \times \bar{A} = 0$	
$A + \bar{A} = 1$	
$AB = BA$	Commutative law for multiplication
$A + B = B + A$	Commutative law for addition
$(AB)C = A(BC)$	Associative law for multiplication
$A + (B + C) = (A + B) + C$	Associative law for addition
$A(B + C) = AB + AC$	Left distributive law
$(B + C)A = BA + CA$	Right distributive law
$\overline{A + B} = \bar{A}\bar{B}$	} De Morgan's laws
$\overline{AB} = \bar{A} + \bar{B}$	

$$\begin{aligned}
 AB + A\bar{B} &= A \\
 A + AB &= A \\
 (A + \bar{B})B &= AB \\
 (A + B)(A + \bar{B}) &= A \\
 (A + B)(A + C) &= A + BC \\
 A(A + B) &= A \\
 A\bar{B} + B &= A + B \\
 \bar{A}B + A\bar{B} &= A \oplus B
 \end{aligned}$$


Exclusive OR

Logic gates


Summary of the elementary positive and negative logic gates. (The positive logic gates in the left column are equivalent to the corresponding negative gates in the right column.)



OR




Negative AND




AND



Negative OR



NOR



Negative NAND



NAND



Negative NOR


Positive logic

Negative logic

The RS-232-C standard

The RS-232-C serial interface standard defines the binary state 1 as a voltage level between $+3\text{ V}$ and -15 V . The binary state 0 can range from $+3\text{ V}$ to $+15\text{ V}$. (Adapted from Libes, S. and Garetz, M. *Interfacing to S-100/IEEE Microcomputers*, Osborne/McGraw-Hill, Berkeley, CA.)

DB-25 type connector





Female Connector as Viewed from Wiring Side

RS-232 C

Pin Number	EIA Name	Common Mnemonic	Description
1	AA		Protective ground
2	BA	TxD	Data transmitted from terminal
3	BB	RxD	Data received from modem
4	CA	RTS	Request to send
5	CB	CTS	Clear to send
6	CC	DSR	Data set ready
7	AB		Signal ground
8	CF	DCD	Carrier detect
9	*		Reserved for Data Set Testing
10	*		Reserved for Data Set Testing
11	*		Unassigned
12	SCF		Secondary carrier detector
13	SCB		Secondary clear to send
14	SBA		Secondary transmitted data
15	SB		Transmitted bit clock from DCE
16	SBB		Secondary received data
17	DD		Received bit clock
18	*		Unassigned
19	SCA		Secondary request to send
20	CD	QTR	Data terminal ready
21	CG		Signal quality detector
22	CE		Ring indicator (used by auto answer equipment)
23	CA-CI		Data signal rate selector
24	DA		Transmitted bit clock from DTE
25	*		Unassigned

RS-232 C Connector and Pin definitions
 DTE: data terminal equipment
 DCE: data communications equipment




RS-232 Cable Wiring

RS-232-C Signal Level Conventions

Notation	Interchange Voltage	
	Negative	Positive
Signal Condition	Marking	Condition
Ready State (Data lines)	OFF	ON
Function (control lines)	OFF	ON


Parallel interface

Centronics interface connector pinouts as they appear on the IBM PC's 25-pin D-shell connector. The pin connections on the printer end are the same for lines 1–14 and 19–25 but differ somewhat on the other lines, since a 36-pin connector is used there. (From Sargent, M. & Shoemaker, R., *The IBM PC from the Inside Out*, Addison-Wesley, 1986, with permission.)



IEEE 488 interface

Example of an IEEE 488 interface bus (GPIB) configuration. (Intel Corp.)




DAV data valid
 NRFD not ready for data
 NDAC not data accepted
 IFC interface clear
 ATN attention
 SRQ service request
 REN remote unable
 EOI end or indentify

Data transmission

Data transmission nomogram.

Transmission time (s) – size (bytes); data rate (byte s⁻¹)

(1 byte = 8 bits; T1 and T3 are telephone transmission standards.)



Bibliography


- Techniques of Vacuum Ultraviolet Spectroscopy*, Samson, J. A. R., John Wiley and Sons, 1967.
- The Atomic Nucleus*, Evans, R. B., McGraw-Hill Book Company, 1955.
- Handbook of Chemistry and Physics*, CRC Press, 1979.
- Table of Isotopes*, C. M. Lederer & V. S. Shirley, eds., John Wiley and Sons, 1978.
- 'Low energy X-ray interaction coefficients: photoionization, scattering, and reflection', Henke, B. L. *et al.*, *Atomic Data and Nuclear Data Tables*, **27**, no. 1, 1982.
- Vacuum Technology and Space Simulation*, NASA SP-105, US Government Printing Office, 1966.
- 'X-ray wavelengths', Bearden, J. A., *Rev. Mod. Phys.*, **39**, 78, 1967.
- X-ray Data Booklet*, Lawrence Berkeley Laboratory, PUB-490 Rev., University of California, 1986.

Chapter 15

Aeronautics and astronautics

	<i>page</i>
Approximate lifetimes for Earth satellites	380
Space transportation system	381
Shuttle performance capability	382
Launch vehicles	384
<i>Current US space launch capabilities</i>	384
<i>US launch vehicles</i>	387
<i>French launch vehicles</i>	390
<i>Japanese launch vehicles</i>	393
<i>Soviet launch vehicles</i>	394
Bibliography	394

Approximate lifetimes for Earth satellites



$$\text{Lifetime} = NP = N \left[\frac{4\pi^2}{GM_E} \left(\frac{h_p + R_E}{1 - e} \right)^3 \right]^{1/2}$$

where

N = number of orbit revolutions in the satellite lifetime from the above diagram.

Ballistic coefficient = $m/(C_D A)$ (typically, 25–100 kg m⁻²),

m = mass of the satellite,

C_D = the drag coefficient (≈ 1.2),

A = satellite cross-sectional area perpendicular to the velocity vector.

G = gravitational constant,

h_p = perigee height,

e = eccentricity of the orbit,

P = satellite period,

M_E = mass of the Earth,

R_E = radius of the Earth.

$$\text{Lifetime (d)} \approx 1.15 \times 10^{-7} \times N \times \left(\frac{6378.14 + h_p \text{ (km)}}{1 - e} \right)^{3/2}$$


$$\text{Satellite period (hr)} = 1.41(a/R_E)^{3/2}$$

where a = semi-major axis.


(Adapted from Wertz, J., ed., *Spacecraft Attitude Determination and Control*, D. Reidel Publ. Co., 1980.)

Space transportation system (Space Shuttle)

Orbiter coordinate system and cargo bay envelope. The dynamic clearance allowed between the vehicle and the payload at each end is also illustrated.




Principal Orbiter interfaces with payloads.




Shuttle performance capability


Launch azimuth and inclination limits from KSC in Florida. The inset globe illustrates the extent of coverage possible when launches are made from KSC.




Cargo capability for a KSC (28.45 degree inclination) launch.



Launch azimuth and inclination limits from VAFB in California.
 Note added in proof: no longer to be used for shuttle launches.



Cargo capability for a KSC (57 degree inclination) launch.



Launch vehicles

Current US space launch capabilities

Vehicle	Stages	Thrust, kilonewtons (pounds × 1000)	Launch sites†	Orbit, kilometers (nautical miles)	Inclina- tion (degrees)	Payload weight, kilograms (pounds)
SCOUT	1. Algol III A	431.1(96.9)	SLC-5 WSMC	556(300)	0	181(400)
	2. Castor II A	285.2(64.1)				
	3. Antares III A	83.1(18.7)				
	4. Altair III A	25.6(5.8)				
Delta 3914	0. 9 Castor IVs	378.0(85.0) ea	LC-17 A&B	185 × 35 786 (100 × 19 323)	27	937(2065)
	1. Extended long tank	921.0(207.0)	FSMC			
	2. Standard 2nd stage	41.8(9.4)	SLC-2W			
	3. TE-M-364-4 solid rocket motor	65.8(14.8)	WSCS			
Delta 3910/ PAM	0. 9 Castor IVs	378.0(85.0) ea	LC-17 A&B	296 × 35 786 (160 × 19 323)	27	1111(2450)
	1. Extended long tank	921.0(207.0)	ESMC			
	2. Standard 2nd stage	41.8(9.4)	SLC-2W			
	3. Payload assist module (PAM)	84.5(19.0)	WSMC			
Delta 3920/ PAM	0. 9 Castor IVs	378.0(85.0) ea	LC-17 A&B	296 × 35 786 (160 × 19 323)	27	1270(2800)
	1. Extended long tank	921.0(207.0)	ESMC			
	2. Improved 2nd stage	41.8(9.4)	SLC-2W			
	3. Payload assist module (PAM)	84.5(19.0)	WSMC			


Atlas E	1. MA-3	SLC-3	157 × 20 198		
	2-Boosters 1-Sustainer	WSMC	(85 × 10 900)	6.3	1088(2398)
Atlas H	2. SGS II				
	1. MA-5	LC-36			
Atlas SLV-3D Centaur	2-Boosters	ESMC	157 × 20 198	55	1225(2700)
	1-Sustainer	SLC-3	(85 × 10 900)		
Atlas G Centaur	2. SGS II	WSMC			
	1. MA-5	LC-36	185 × 35 786	27	1996(4400)
Titan 34B Agena	2-Boosters	ESMC	(100 × 19 323)		
	1-Sustainer				
Titan 34D NUS	2. Centaur (RL-10)				
	1. MA-5	LC-36	185 × 35 786	27	2359(5200)
Titan 34D Transtage	2-Boosters	ESMC	(100 × 19 323)		
	1-Sustainer				
Titan 34D Transtage	2. Centaur (RL-10-3A)				
	1. LR-87	LC-36	185 × 35 786	27	2359(5200)
Titan 34D Transtage	2. LR-91	ESMC	(100 × 19 323)		
	3. Ascent Agena				
Titan 34D Transtage	0. Two 5½ segment SRMs				
	1. LR-87	SLC-4	185(100)	90	3583(7900)
Titan 34D Transtage	2. LR-91	WSMC	Circular		
	0. Two 5½ segment SRMs	SLC-4	185(100)	90	12 520(27 600)
Titan 34D Transtage	1. LR-87	WSMC	Circular		
	2. LR-91				
Titan 34D Transtage	0. Two 5½ segment SRMs				
	1. LR-87	LC-40	35 876(19 323)	0	843(1859)
Titan 34D Transtage	2. LR-91	ESMC	Circular		
	3. Transtage				

Current US space launch capabilities (cont.)

Vehicle	Stages	Thrust, kilonewtons (pounds × 1000)	Launch sites†	Orbit, kilometers (nautical miles)	Inclina- tion (degrees)	Payload weight, kilograms (pounds)
Titan 34D IUS	0. Two 5½ segment SRMs	12 454.4(2800.0)	LC-40	35 876(19 323) Circular	0	843(1859)
	1. LR-87	2353.0(529.0)	ESMC			
	2. LR-91	449.2(101.0)				
	3. IUS					
	Stage 1 Stage 2	186.8(42.0) 75.6(17.0)				
STS	0. 2 Solid rocket boosters	11 155.0(2508.0) ea	LC-39	278(150) Circular	28.5	29 500 (65 000)
	1. 3 Main engines	1668.0(375.0) ea	ESMC			
	2. Inertial upper stage (Two stage)	196.0(44.1) 74.7(16.8)				
	or 2. Inertial upper stage (Twin stage)					
	STS Parking orbit		35 786(19 323) Circular	0	2267(5000)	
	STS Parking orbit		278 × 35 786 Parking orbit (150 × 19 323)	Planetary missions	N/A	4993(11 009)
	or 2. PAM-D	196.2(44.1) 196.2(44.1) 84.5(19.0)	STS Parking orbit STS Parking orbit	278 × 35 786 Parking orbit (150 × 19 323)	27	1052(2320)


† Launch sites at the Eastern Space and Missile Center (ESMC) are designated as launch complexes (LC) and if served by the same launch control center, are further designated as A or B. Launch sites at the Western Space and Missile Center (WSMC) are designated as Space Launch complexes (SLC) and if served by the same launch control center are further designated as E(ast) or W(est).

US launch vehicles



Scout


(Ling-Temco-Vogt.)




Delta

(McDonnell Douglas Astronautics Co.)


US launch vehicles (cont.)




(General Dynamics Convair Division.)



Ascent Agena



TITAN 34B




TITAN 34D/
No Upper Stage

STAGE II			
Height	m (feet)	9.6 (31.3)	9.6 (31.3)
Diameter	m (feet)	3.1 (10.1)	3.1 (10.0)
Thrust	kn (pounds × 1,000)	449.2 (101.0)	449.2 (101.0)
STAGE I			
Height	m (feet)	23.8 (77.8)	23.8 (77.8)
Diameter	m (feet)	3.1 (10.0)	3.1 (10.0)
Thrust	kn (pounds × 1,000)	2,059.4 (463.0)	2,353.0 (529.0)
STAGE 0			
Height	m (feet)	N/A	27.6 (90.4)
Diameter	m (feet)		3.1 (10.2) ea
Thrust	kn (pounds × 1,000)		6,227.2 (1,400.0) ea


Titan

(Martin Marietta Denver Aerospace.)

French launch vehicles




(Diagrams courtesy Arianespace.)



The Ariane 4 configurations

(Diagrams courtesy Arianespace.)


ARIANE 5



(Aérospatiale.)

Japanese launch vehicles

Rockets of Japan and other typical rockets in the world





Rocket name	M-3SII (ISAS)	N-I (NASDA)	N-II (NASDA)	H-I (NASDA)	H-II (NASDA)	CZ-3 (CHINA)	A-2 (USSR)	ARIANE 4 (ESA)	ARIANE 5 (ESA)	ARIANE 5 Space Shuttle (USA)
Total weight (t)	61	90	135	140	258	-	-	460	500	2041
Payload capacity into LEO ¹ (kg)	670	1200	2000	3000	8000	5000	-	8000	15 000	29 500
Payload capacity into GEO ¹ (kg)	-	130	350	550	2000	-	-	2200	4000	2400 <small>(using upper stage)</small>

¹LEO Low Earth Orbit
²GEO Geostationary Earth Orbit

Reproduced from General Outline of H-II Rockets, published by NASDA, the National Space Development Agency of Japan. (ISAS refers to the Japanese Institute of Space and Astronomical Science, ESA to the European Space Agency.)

Soviet launch vehicles



US Department of Defense.

¹ Approximate
² In final stages of development

Bibliography

Spacecraft Attitude Determination and Control, J. R. Wertz, ed.,
 D. Reidel Publishing Co., 1980.

Space Transportation System User Handbook, National Aeronautics and
 Space Administration.


Chapter 16

Mathematics

	<i>page</i>
Coordinate transformations	396
Vector analysis	396
<i>Differential elements</i>	398
Definite integrals	399
The Fourier transform	399
<i>Useful definitions</i>	400
<i>Fourier transform theorems</i>	401
<i>Fourier transform pairs</i>	402
Special functions	403
<i>Spherical harmonics</i>	403
<i>Bessel functions</i>	404
Numerical analysis	404
<i>Taylor series</i>	404
<i>Quadrature</i>	404
<i>Six-point Gauss-Legendre</i>	405
<i>Linear interpolation</i>	405
<i>Approximations</i>	405
<i>Curve fitting (linear least-squares)</i>	406
Bibliography	406

Coordinate transformations

Components of a vector, \mathbf{r} , in Cartesian (x, y, z) , spherical (r, θ, ϕ) , and cylindrical (ρ, ϕ, z) coordinates.



The following relationships exist between the Cartesian, Spherical, and Cylindrical systems:

$$\begin{aligned}
 z &= r \cos \theta & &= z \\
 r &= (x^2 + y^2 + z^2)^{1/2} & &= (\rho^2 + z^2)^{1/2} \\
 \theta &= \arccos\{z/(x^2 + y^2 + z^2)^{1/2}\} = \arctan(\rho/z) & &0 \leq \theta \leq \pi \\
 \phi &= \arctan(y/x) & &= \phi \quad 0 \leq \phi < 2\pi \\
 \rho &= (x^2 + y^2)^{1/2} & &= r \sin \theta \\
 x &= r \sin \theta \cos \phi & &= \rho \cos \phi \\
 y &= r \sin \theta \sin \phi & &= \rho \sin \phi
 \end{aligned}$$

Vector analysis

Vectors \mathbf{a} and \mathbf{b} defined in terms of the unit vectors $\mathbf{i}, \mathbf{j}, \mathbf{k}$ having the directions of the positive $x, y,$ and z axes respectively:

$$\mathbf{a} = a_x \mathbf{i} + a_y \mathbf{j} + a_z \mathbf{k}$$

$$\mathbf{b} = b_x \mathbf{i} + b_y \mathbf{j} + b_z \mathbf{k}$$

The scalar product:

$$\mathbf{a} \cdot \mathbf{b} = a_x b_x + a_y b_y + a_z b_z$$

The vector product:

$$\begin{aligned}\mathbf{a} \times \mathbf{b} &= (a_y b_z - a_z b_y)\mathbf{i} + (a_z b_x - a_x b_z)\mathbf{j} + (a_x b_y - a_y b_x)\mathbf{k} \\ &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_x & a_y & a_z \\ b_x & b_y & b_z \end{vmatrix}.\end{aligned}$$

Vector identities:

$$\begin{aligned}\mathbf{a} \cdot \mathbf{b} \times \mathbf{c} &= (\mathbf{a} \times \mathbf{b}) \cdot \mathbf{c} = \mathbf{b} \cdot (\mathbf{c} \times \mathbf{a}) = (\mathbf{b} \times \mathbf{c}) \cdot \mathbf{a} \\ &= \mathbf{c} \cdot (\mathbf{a} \times \mathbf{b}) = (\mathbf{c} \times \mathbf{a}) \cdot \mathbf{b}, \\ \mathbf{a} \times (\mathbf{b} \times \mathbf{c}) &= (\mathbf{a} \cdot \mathbf{c})\mathbf{b} - (\mathbf{b} \cdot \mathbf{a})\mathbf{c}, \\ (\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) &= (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{a} \cdot \mathbf{d})(\mathbf{b} \cdot \mathbf{c}), \\ (\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d}) &= ((\mathbf{a} \times \mathbf{b}) \cdot \mathbf{d})\mathbf{c} - ((\mathbf{a} \times \mathbf{b}) \cdot \mathbf{c})\mathbf{d}.\end{aligned}$$

The vector operator ∇ :

$$\nabla = \mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} = \text{grad}$$

The divergence of a vector function \mathbf{F} :

$$\begin{aligned}\text{div } \mathbf{F} = \nabla \cdot \mathbf{F} &= \mathbf{i} \cdot \frac{\partial \mathbf{F}}{\partial x} + \mathbf{j} \cdot \frac{\partial \mathbf{F}}{\partial y} + \mathbf{k} \cdot \frac{\partial \mathbf{F}}{\partial z}, \\ \nabla \cdot \mathbf{F} &= \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} + \frac{\partial F_z}{\partial z}.\end{aligned}$$

The curl of a vector function \mathbf{F} :

$$\begin{aligned}\text{curl } \mathbf{F} = \nabla \times \mathbf{F} &= \mathbf{i} \times \frac{\partial \mathbf{F}}{\partial x} + \mathbf{j} \times \frac{\partial \mathbf{F}}{\partial y} + \mathbf{k} \times \frac{\partial \mathbf{F}}{\partial z} \\ &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_x & F_y & F_z \end{vmatrix}.\end{aligned}$$

Differentiation formulae:

$$\begin{aligned}\nabla \cdot \phi \mathbf{u} &= \phi \nabla \cdot \mathbf{u} + \mathbf{u} \cdot \nabla \phi, \\ \nabla \times \phi \mathbf{u} &= \phi \nabla \times \mathbf{u} + \nabla \phi \times \mathbf{u}, \\ \nabla \cdot (\mathbf{u} \times \mathbf{v}) &= \mathbf{v} \cdot \nabla \times \mathbf{u} - \mathbf{u} \cdot \nabla \times \mathbf{v}, \\ \nabla \times (\mathbf{u} \times \mathbf{v}) &= (\mathbf{v} \cdot \nabla)\mathbf{u} - (\mathbf{u} \cdot \nabla)\mathbf{v} + \mathbf{u}(\nabla \cdot \mathbf{v}) - \mathbf{v}(\nabla \cdot \mathbf{u}), \\ \nabla(\mathbf{u} \cdot \mathbf{v}) &= (\mathbf{u} \cdot \nabla)\mathbf{v} + (\mathbf{v} \cdot \nabla)\mathbf{u} + \mathbf{u} \times (\nabla \times \mathbf{v}) + \mathbf{v} \times (\nabla \times \mathbf{u}), \\ \nabla \times (\nabla \phi) &= \text{curl grad } \phi = 0, \\ \nabla \cdot (\nabla \times \mathbf{u}) &= \text{div curl } \mathbf{u} = 0,\end{aligned}$$

$$\begin{aligned}\nabla \times (\nabla \times \mathbf{u}) &= \text{curl curl } \mathbf{u} = \nabla(\nabla \cdot \mathbf{u}) - \nabla \cdot \nabla \mathbf{u} \\ &= \text{grad div } \mathbf{u} - \nabla^2 \mathbf{u}\end{aligned}$$

$$\nabla \cdot (\nabla \phi_1 \times \nabla \phi_2) = 0$$

In these formulas \mathbf{u} and \mathbf{v} are arbitrary vectors and ϕ , ϕ_1 , and ϕ_2 are arbitrary scalars for which the indicated derivatives exist.

Surface integrals

Divergence theorem:

$$\iiint_V \nabla \cdot \mathbf{V} \, dV = \oiint_S \mathbf{V} \cdot \mathbf{n} \, dS$$

Stokes' theorem:

$$\iint_S \mathbf{n} \cdot (\nabla \times \mathbf{V}) \, dS = \oint_C \mathbf{V} \cdot d\mathbf{r}$$

Green's theorem:

$$\iiint_V [\phi_1 \nabla^2 \phi_2 - \phi_2 \nabla^2 \phi_1] \, dV = \iint_S \mathbf{n} \cdot (\phi_1 \nabla \phi_2 - \phi_2 \nabla \phi_1) \, dS,$$

where \mathbf{n} is the surface outward unit normal.

Differential elements

Cylindrical coordinates

Line element:

$$ds = \sqrt{(dr^2 + r^2 d\theta^2 + dz^2)}$$

Area elements:

$$dS_r = r \, d\theta \, dz, \quad dS_\theta = dr \, dz, \quad dS_z = r \, d\theta \, dr$$

Volume element:

$$dV = r \, d\theta \, dr \, dz$$

Spherical coordinates

Line element:

$$ds = \sqrt{(dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2)}$$

Area elements:

$$dS_r = r^2 \sin \theta \, d\theta \, d\phi, \quad dS_\theta = r \sin \theta \, dr \, d\phi, \quad dS_\phi = r \, dr \, d\theta$$

Volume element:

$$dV = r^2 \sin \theta \, dr \, d\theta \, d\phi$$

Definite integrals

$$\int_0^{\infty} t^{2n-1} e^{-at} dt = \frac{n!}{2a^{n+1}} \quad (n = 0, 1, 2, \dots)$$

$$\int_0^{\infty} t^{2n} e^{-at} dt = \frac{1 \cdot 3 \cdot \dots \cdot (2n-1)}{2^{n+1} a^{n+1}} \sqrt{\frac{\pi}{a}} \quad (n = 0, 1, 2, \dots)$$

$$\int_0^{\infty} t^n e^{-at} dt = \frac{n!}{a^{n+1}} \quad (n = 1, 2, \dots)$$

$$\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi} \quad \int_{-\infty}^{\infty} e^{-x^2} dx = \left(\frac{\pi}{4}\right)^{1/2}$$

$$\int_0^{\infty} xe^{-x^2} dx = \frac{1}{2} \quad \int_0^{\pi} \left(\frac{\sin x}{x}\right)^2 dx = \frac{\pi}{2}$$

$$\operatorname{erf} x = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt, \quad \text{error function}$$

$$C(x) = \int_0^x \cos\left(\frac{\pi}{2} t^2\right) dt, \quad S(x) = \int_0^x \sin\left(\frac{\pi}{2} t^2\right) dt, \quad \text{Fresnel integrals}$$

$$Ei(x) = \int_{-\infty}^x \frac{e^t}{t} dt, \quad \text{exponential integral}$$

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt, \quad \text{gamma function}$$

$$L\{F(t)\} = \int_0^{\infty} e^{-st} F(t) dt, \quad \text{Laplace transform}$$

The Fourier transform

The Fourier transform of $f(x)$ is:

$$F(s) = \int_{-\infty}^{\infty} f(x) e^{-i2\pi sx} dx$$

and

$$f(x) = \int_{-\infty}^{\infty} F(s) e^{i2\pi sx} ds.$$

There are two other equivalent versions:

$$F(s) = \int_{-\infty}^{\infty} f(x) e^{-i\pi sx} dx,$$

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(s) e^{i\pi sx} ds$$

and

$$F(s) = \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{\infty} f(x) e^{-ixs} dx,$$

$$f(x) = \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{\infty} F(s) e^{ixs} ds,$$

which are also used.

Useful definitions

The *convolution* of two functions $f(x)$ and $g(x)$ is:

$$\int_{-\infty}^{\infty} f(u)g(x-u) du = f * g = \int_{-\infty}^{\infty} g(u)f(x-u) du = g * f.$$

The *autocovariance* function of $f(x)$ is:

$$f(x) \otimes f(x) = \int_{-\infty}^{\infty} f^*(u)f(u+x) du$$

The *autocorrelation* function of $f(x)$ is:

$$\gamma(x) = \frac{\int_{-\infty}^{\infty} f^*(u)f(u+x) du}{\int_{-\infty}^{\infty} f(u)f^*(u) du},$$

$$\gamma(0) = 1.$$

The *cross-correlation* of two functions $g(x)$ and $h(x)$ is:

$$f(x) = \int_{-\infty}^{\infty} g(u-x)h(u) du$$

The *power spectrum* of a function $f(x)$ is:

$$\left| \int_{-\infty}^{\infty} f(x) e^{-i2\pi xs} dx \right|^2 = |F(s)|^2$$

The *normalized power spectrum* is the *power spectral density function*:

$$|\phi(s)|^2 = \frac{|F(s)|^2}{\int_{-\infty}^{\infty} |F(s)|^2 ds}$$

The *equivalent width* of a function $f(x)$ is:

$$W_f = \frac{\int_{-\infty}^{\infty} f(x) dx}{f(0)}$$

The *filtering or interpolation function*, *sinc* x :

$$\text{sinc } x = \frac{\sin \pi x}{\pi x}$$

Fourier transform theorems

Theorem	$f(x)$	$F(s) = \int_{-\infty}^{\infty} f(x)e^{-i2\pi sx} dx$
Similarity	$f(ax)$	$\frac{1}{ a } F(s/a)$
Addition	$f(x) + g(x)$	$F(s) + G(s)$
Shift	$f(x - a)$	$e^{-i2\pi as} F(s)$
Convolution	$f(x) * g(x)$	$F(s)G(s)$
Autocovariance	$f(x) \otimes f(x)$	$ F(s) ^2$
Derivative	$f'(x)$	$i2\pi sF(s)$

Derivative of convolution:

$$\frac{d}{dx} [f(x) * g(x)] = f'(x) * g(x) = f(x) * g'(x)$$

Parseval:

$$\int_{-\infty}^{\infty} |f(x)|^2 dx = \int_{-\infty}^{\infty} |F(s)|^2 ds$$

Multiplication:

$$\int_{-\infty}^{\infty} f^*(x)g(x) dx = \int_{-\infty}^{\infty} F^*(s)G(s) ds$$

Sampling theorem:

A function whose Fourier transform is zero for $|s| > s_c$ is fully specified by values spaced at equal intervals not exceeding $\frac{1}{2}s_c^{-1}$ save for any harmonic term with zeros at the sampling points.

(Adapted from Bracewell, R. N., *The Fourier Transform and its Applications*, McGraw-Hill Book Company, 1978.)

Fourier transform pairs

	$f(t) = \int_{-\infty}^{\infty} F_F(i\omega) e^{-i\omega t} \frac{d\omega}{2\pi}$	$F_F(i\omega) = \int_{-\infty}^{\infty} f(t) e^{i\omega t} dt$	
	$\text{rect } \frac{t}{T} = \begin{cases} 1 & (t < T/2) \\ 0 & (t > T/2) \end{cases}$	$T \text{ sinc } \frac{\omega T}{2\pi} \equiv T \frac{\sin \frac{\omega T}{2}}{\frac{\omega T}{2}}$	
	$\text{sinc } \frac{t}{T} \equiv \frac{\sin \frac{\pi t}{T}}{\frac{\pi t}{T}}$	$T \text{ rect } \frac{\omega T}{2\pi} = \begin{cases} 0 & (\omega < \frac{\pi}{T}) \\ T & (\omega > \frac{\pi}{T}) \end{cases}$	
	$\begin{cases} 1 - \frac{ t }{T} & (t < T) \\ 0 & (t \geq T) \end{cases}$	$T \text{ sinc}^2 \frac{\omega T}{2\pi} \equiv T \left(\frac{\sin \frac{\omega T}{2}}{\frac{\omega T}{2}} \right)^2$	
	$e^{-\frac{ t }{T}}$	$\frac{2T}{(\omega T)^2 + 1}$	
	$e^{-\frac{1}{2}(\frac{t}{T})^2}$	$\sqrt{2\pi} T e^{-\frac{1}{2}\omega T^2}$	
	$\delta(t-T)$	$e^{-i\omega T}$	(Complex)
	$\cos \omega_0 t$	$\pi [\delta(\omega - \omega_0) + \delta(\omega + \omega_0)]$	
	$\sin \omega_0 t$	$\frac{\pi}{i} [\delta(\omega - \omega_0) - \delta(\omega + \omega_0)]$	(Imaginary)
	$\sum_{k=-\infty}^{\infty} \delta(t-kT) \equiv \frac{1}{T} \sum_{j=-\infty}^{\infty} e^{2\pi i j t}$	$\frac{2\pi}{T} \sum_{j=-\infty}^{\infty} \delta(\omega - \frac{2\pi j}{T}) \equiv \sum_{k=-\infty}^{\infty} e^{ik\omega T} T$	

(From Korn, G. A., *Basic Tables in Electrical Engineering*, McGraw-Hill, New York, 1965, with permission.)

Special functions

Spherical harmonics

$$Y_l^m(\theta, \phi) = \sqrt{\left[\frac{(2l+1)(l-m)!}{4\pi(l+m)!} \right]} P_l^m(\cos \theta) e^{m\phi}$$

$$P_l^m(\cos \theta) = (-1)^m \sin^m \theta \left[\left(\frac{d}{d(\cos \theta)} \right)^m P_l(\cos \theta) \right] \quad (m \leq l)$$

$$P_l(\cos \theta) = \frac{1}{2^l l!} \left[\left(\frac{d}{d(\cos \theta)} \right)^l (-\sin^2 \theta)^l \right]$$

$$Y_l^{-m}(\theta, \phi) = (-1)^m [Y_l^m(\theta, \phi)]^*$$

$$P_l^{-m}(\cos \theta) = (-1)^m \frac{(l-m)!}{(l+m)!} P_l^m(\cos \theta)$$

$$l=0 \quad Y_0^0 = \frac{1}{\sqrt{4\pi}}$$

$$l=1 \quad Y_1^0 = \sqrt{\left(\frac{3}{4\pi} \right)} \cos \theta$$

$$Y_1^1 = -\sqrt{\left(\frac{3}{8\pi} \right)} \sin \theta e^{i\phi}$$

$$l=2 \quad Y_2^0 = \sqrt{\left(\frac{5}{16\pi} \right)} (3 \cos^2 \theta - 1)$$

$$Y_2^1 = -\sqrt{\left(\frac{15}{8\pi} \right)} \sin \theta \cos \theta e^{i\phi}$$

$$Y_2^2 = \sqrt{\left(\frac{15}{32\pi} \right)} \sin^2 \theta e^{2i\phi}$$


$$l=3 \quad Y_3^0 = \sqrt{\left(\frac{7}{16\pi} \right)} (5 \cos^3 \theta - 3 \cos \theta)$$

$$Y_3^1 = -\sqrt{\left(\frac{21}{64\pi} \right)} \sin \theta (5 \cos^2 \theta - 1) e^{i\phi}$$

$$Y_3^2 = \sqrt{\left(\frac{105}{32\pi} \right)} \sin^2 \theta \cos \theta e^{2i\phi}$$

$$Y_3^3 = -\sqrt{\left(\frac{35}{64\pi} \right)} \sin^3 \theta e^{3i\phi}$$

Bessel functions



$$J_n(x) = \sum_{s=0}^{\infty} \frac{(-1)^s}{s!(n+s)!} \left(\frac{x}{2}\right)^{n+2s} \quad J_{n-1}(x) + J_{n+1}(x) = (2n/x)J_n(x)$$

$$J_0(x) = (2/\pi x)^{1/2} \cos(x - \pi/4)$$

(Adapted from Chantry, G. U. J., *Long-wave Optics*, Academic Press, New York, 1984.)

Numerical analysis

Taylor series

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + f''(x_0) \frac{(x - x_0)^2}{2!} + \dots$$

$$+ f^{(n)}(x_0) \frac{(x - x_0)^n}{n!} + R_n$$

$$R_n = f^{(n+1)}(\zeta) \frac{(x - x_0)^{n+1}}{(n+1)!} \quad (x_0 < \zeta < x)$$

Quadrature

Trapezoidal rule

$$\int_a^b f(x) dx = h \left(\frac{y_0}{2} + y_1 + y_2 + \dots + y_{n-1} + \frac{y_n}{2} \right) - R_n$$

$$R_n = \frac{1}{12}(b-a)h^2 f''(x_1) \quad (a < x_1 < b)$$

$$h = (b-a)/n, \quad y_k = f(a + kh), \quad k = 0, 1, \dots, n.$$

Simpson's rule (n even)

$$\int_a^b f(x) dx = \frac{h}{3} [y_0 + 4(y_1 + y_3 + \cdots + y_{n-1}) + 2(y_2 + y_4 + \cdots + y_{n-2}) + y_n] + R_n$$

$$R_n = \frac{1}{90}(b-a)h^4 f^{(4)}(x_1) \quad (a < x_1 < b), \quad h = (b-a)/n.$$

Six-point Gauss Legendre

$$\int_a^b f(x) dx \approx \frac{b-a}{2} \sum_{i=1}^6 w_i f\left(z_i \frac{b-a}{2} + \frac{b+a}{2}\right)$$

$$\int_a^1 f(x) dx \approx 2 \sum_{i=1}^6 \frac{w_i}{(1+z_i)^2} f\left(\frac{2}{1+z_i} + a - 1\right)$$

where

$$z_1 = -z_2 = 0.238\ 619\ 186\ 1$$

$$z_3 = -z_4 = 0.661\ 209\ 386\ 5$$

$$z_5 = -z_6 = 0.932\ 469\ 514\ 2$$

$$w_1 = w_2 = 0.467\ 913\ 934\ 6$$

$$w_3 = w_4 = 0.360\ 761\ 573$$

$$w_5 = w_6 = 0.171\ 324\ 492\ 4.$$

Linear interpolation

$$y = \frac{(x_{k+1} - x)y_k + (x - x_k)y_{k+1}}{x_{k+1} - x_k} \quad \text{for } x_k < x < x_{k+1}.$$

Approximations

$f(x)$	Approximation	Parameters	Maximum absolute error
$\log_{10} x$ ($1/\sqrt{10} \leq x \leq \sqrt{10}$)	$a_1 t + a_3 t^3$ $t = (x-1)/(x+1)$	$a_1 = 0.863\ 04$ $a_3 = 0.364\ 15$	6×10^{-4}
$\arctan x$ ($-1 \leq x \leq 1$)	$\frac{x}{1 + 0.28x^2}$		5×10^{-3}
$\operatorname{erf} x$ ($0 \leq x < \infty$)	$1 - (a_1 t + a_2 t^2 + a_3 t^3)e^{-x^2}$ $t = 1/(1 + px)$	$p = 0.47047$ $a_1 = 0.348\ 024\ 2$ $a_2 = -0.095\ 879\ 8$ $a_3 = 0.747\ 855\ 6$	2.5×10^{-5}

(Adapted from Hastings, C., *Approximations for Digital Computers*, Princeton, New Jersey, 1955.)

Curve fitting (linear least-squares)

$$y(x) = \sum_{j=0}^n a_j \phi_j(x), \quad \text{approximating function}$$

$$\chi^2 = \sum_{i=1}^N \frac{1}{\sigma_i^2} [y_i - y(x_i)]^2,$$

where

σ_i = standard deviation of the i th observation y_i ,

N = number of observations.

Minimizing χ^2 , $\frac{\partial}{\partial a_k} \chi^2 = 0$, yields the following set of normal equations:

$$\sum_{i=1}^N \left[\frac{1}{\sigma_i^2} y_i \phi_k(x_i) \right] = \sum_{j=0}^n \left(a_j \sum_{i=1}^N \left[\frac{1}{\sigma_i^2} \phi_j(x_i) \phi_k(x_i) \right] \right), \quad \text{for } k = 0 \text{ to } n.$$

(For the case of a least-squares fit to a straight line, see Chapter 17.)

Bibliography

- Handbook of Mathematical Functions*, Abramowitz, M. & Stegun, I. A.,
Dover Publications, Inc., New York, 1972.
- The Fourier Transform and Its Applications*, Bracewell, R. N., McGraw-
Hill, New York, 1978.
- The Fast Fourier Transform*, Brigham, E. O., Prentice-Hall, Inc., New
Jersey, 1974.
- Table of Integrals, Series, and Products*, Gradshteyn, I. S. &
Ryzhik, I. M., Academic Press, New York, 1980.
- A Table of Series and Products*, Hansen, E. R., Prentice-Hall, Inc., New
Jersey, 1975.
- Tables of Higher Functions*, Jahnke-Emde-Loesch, McGraw-Hill, New
York, 1960.
- Mathematical Methods for Digital Computers*, A. Ralston & H. S. Wilf,
eds., John Wiley and Sons, New York, 1960.
- Numerical Recipes: The Art of Scientific Computing*, Press, W. H.,
Flannery, B., Teukolsky, S. & Vetterling, W., Cambridge University
Press, 1985.

Chapter 17

Statistics

	<i>page</i>
Probability distributions	408
<i>Gaussian distribution</i>	408
<i>Binomial distribution</i>	410
<i>Poisson distribution</i>	410
<i>Chi-square distribution, confidence levels</i>	413
χ^2 test of distributions	414
Propagation of errors	415
Least-squares fit to a straight line	415
Bibliography	416

Probability distributions

Gaussian distribution

$$P_G(x, \mu, \sigma) = (\sigma\sqrt{2\pi})^{-1} \exp\left[-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2\right],$$

where

x = value of random observation,

μ = mean value of parent distribution,

σ = standard deviation of parent distribution,

σ^2 = variance.


FWHM:

$$\Gamma = 2.354\sigma.$$


Probable error:

$$PE = 0.6745\sigma.$$

Gaussian probability distribution $P_G(x, \mu, \sigma)$ vs. $x - \mu$; $\Gamma = 2.345 \sigma$; probable error (PE) = 0.6745σ . (Adapted from Bevington, P. R., *Data Reduction and Error Analysis for the Physical Sciences*, McGraw-Hill Book Company, 1969.)



The Gaussian probability distribution $P_G(x, \mu, \sigma)$ vs. $z = (x - \mu) / \sigma$.




The integral of the Gaussian probability distribution

$A_G(x, \mu, \sigma)$ vs. $z = (x - \mu) / \sigma$, where

$$A_G(x, \mu, \sigma) = \int_{\mu - z\sigma}^{\mu + z\sigma} P_G(x, \mu, \sigma) dx$$

(Adapted from Bevington, P. R., *Data Reduction and Error Analysis for the Physical Sciences*, McGraw-Hill Book Company, 1969.)



Binomial distribution

$$P_B(x, n, p) = \frac{n!}{x!(n-x)!} p^x (1-p)^{n-x}$$

If p is the probability that an event will occur, then in a random group of n independent trials, P_B is the probability that the event will occur x times.

$$\mu = \text{mean} = np$$

$$\sigma^2 = \text{variance} = np(1-p)$$

Poisson distribution

$$P_p(x, \mu) = \frac{\mu^x}{x!} e^{-\mu}$$


P_p is the probability of observing x events when the average for a large number of trials is μ events.

$$\sigma^2 = \text{variance} = \mu = \text{mean}$$

The probability of observing at least s events is:


$$P_p(x \geq s, \mu) = \sum_{x=s}^{\infty} \frac{\mu^x}{x!} e^{-\mu}$$

Probability of n_t or more random events with Poisson distribution when the expected or mean number of events is \bar{n} as a function of the threshold number n_t . (From RCA Electro-Optics Handbook, 1974.)



Poisson distribution (cont.)

Probability of n_1 or more random events with Poisson distribution when the expected or mean number of events is \bar{n} as a function of the threshold number n_1 . Curves for $10^{-7} < \bar{n} < 10^{-4}$ are approximate. (From *RCA Electro-Optics Handbook*, 1974.)



Chi-square distribution, confidence levels

The chi-square distribution for n_D degrees of freedom is given by:


$$P_{n_D}(\chi^2) d\chi^2 = \frac{1}{2^h \Gamma(h)} (\chi^2)^{h-1} e^{-\chi^2/2} d\chi^2$$

where h (for "half") = $n_D/2$.

The confidence level CL associated with a given value of n_D and an observed χ^2 is the probability of χ^2 exceeding the observed value:

$$CL = \int_{\chi^2}^{\infty} d\chi^2 P_{n_D}(\chi^2)$$

χ^2 confidence level vs. χ^2 for n_D degrees of freedom. (From *Review of Particle Properties*, Lawrence Berkeley Laboratory, University of California, Berkeley, CA, 1982.)



χ^2 test of distributions

$$\chi^2 = \sum_{j=1}^n \frac{[f(x_j) - NP(x_j)]^2}{NP(x_j)}$$

where

$f(x_j)$ = observed frequency distribution of possible observation x_j ,

n = number of different values of x_j observed,

N = total number of measurements,

$P(x_j)$ = theoretical probability distribution.

$\chi_v^2 = \chi^2/v$ (for χ^2 tests, χ_v^2 should be $\cong 1$),


v = degrees of freedom = n - number of parameters

calculated from the data to describe the distribution.

The probability $P_\nu(\chi^2, \nu)$ of exceeding χ^2 vs. the reduced chi-square $\chi_v^2 = \chi^2/v$ and the number of degrees of freedom ν .

$$P_\nu(\chi^2, \nu) = \frac{1}{2^{\nu/2} \Gamma(\nu/2)} \int_{\chi^2}^{\infty} (x^2)^{\nu/2 - 2} e^{-x^2/2} dx^2$$

(Adapted from Bevington, P. R., *Data Reduction and Error Analysis for the Physical Sciences*, McGraw-Hill Book Company, 1969.)



Propagation of errors

Sample mean:

$$\bar{x} = \frac{1}{N} \sum x_i \approx \mu, \quad \text{parent mean.}$$

Sample variance:

$$s^2 = \frac{1}{N-1} \sum (x_i - \bar{x})^2 \approx \sigma^2, \quad \text{parent variance.}$$

For $x = f(u, v, \dots)$

$x = f(\bar{u}, \bar{v}, \dots)$, the most probable value for x , and

$$\sigma_x^2 \approx \sigma_u^2 \left(\frac{\partial x}{\partial u} \right)^2 + \sigma_v^2 \left(\frac{\partial x}{\partial v} \right)^2 + 2\sigma_{uv} \left(\frac{\partial x}{\partial u} \right) \left(\frac{\partial x}{\partial v} \right) + \dots,$$

where

$$\sigma_u^2 = \lim_{N \rightarrow \infty} \frac{1}{N} \sum (u_i - \bar{u})^2, \quad \sigma_v^2 = \lim_{N \rightarrow \infty} \frac{1}{N} \sum (v_i - \bar{v})^2$$

and

$$\sigma_{uv} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum [(u_i - \bar{u})(v_i - \bar{v})], \quad \text{the covariance.}$$

For u and v uncorrelated, $\sigma_{uv}^2 = 0$.

Least-squares fit to a straight line

Linear function:

$$y(x) = a + bx.$$

Chi-square:

$$\chi^2 = \sum \left[\frac{1}{\sigma_i^2} (y_i - a - bx_i)^2 \right].$$

σ_i = standard deviation of the observation y_i .

Least-squares fitting procedure:

minimize χ^2 with respect to each of the coefficients a and b simultaneously.

Coefficients of least-squares fit:

$$a = \frac{1}{\Delta} \cdot \left(\sum \frac{x_i^2}{\sigma_i^2} \sum \frac{y_i}{\sigma_i^2} - \sum \frac{x_i}{\sigma_i^2} \sum \frac{x_i y_i}{\sigma_i^2} \right)$$

$$b = \frac{1}{\Delta} \cdot \left(\sum \frac{1}{\sigma_i^2} \sum \frac{x_i y_i}{\sigma_i^2} - \sum \frac{x_i}{\sigma_i^2} \sum \frac{y_i}{\sigma_i^2} \right)$$

$$\Delta = \sum \frac{1}{\sigma_i^2} \sum \frac{x_i^2}{\sigma_i^2} - \left(\sum \frac{x_i}{\sigma_i^2} \right)^2$$

Estimated variance:

$$\sigma^2 \simeq s^2 = \frac{1}{N-2} \sum (y_i - a - bx_i)^2$$

s^2 = sample variance

Uncertainties in coefficients:

$$\sigma_a^2 \simeq \frac{1}{\Delta} \sum \frac{x_i^2}{\sigma_i^2} \quad \sigma_b^2 \simeq \frac{1}{\Delta} \sum \frac{1}{\sigma_i^2}$$

Bibliography

Statistical Methods in Experimental Physics, Eadie, W. T. et al., North-Holland Publishing Co., 1971.

Data Reduction and Error Analysis for the Physical Sciences, Bevington, P. R., McGraw-Hill Book Company, 1969.

Data Analysis for Scientists and Engineers, Meyer, S. L., Peer Management Consultants, Ltd., Evanston, IL, 1986.

Chapter 18

Radiation safety

	<i>page</i>
Radiation physics	418
Maximum permissible flux	419
Clinical effects of acute exposure to ionizing radiation	420
Bibliography	421

Radiation physics

Unit of activity = Curie:

$$1 \text{ Ci} = 3.7 \times 10^{10} \text{ disintegration s}^{-1}.$$

Unit of exposure dose for X- and γ -radiation = Roentgen:

$$1 \text{ R} = 1 \text{ esu cm}^{-3} = 87.8 \text{ erg g}^{-1} (5.49 \times 10^7 \text{ MeV g}^{-1}) \text{ of air.}$$

Unit of absorbed dose = rad:

$$1 \text{ rad} = 100 \text{ erg g}^{-1} (6.25 \times 10^7 \text{ MeV g}^{-1}) \text{ in any material.}$$

Unit dose equivalent (for protection) = rem:

$$\text{rems (Roentgen equivalent for man)} = \text{rad} \times QF.$$

where QF (quality factor) depends upon the type of radiation and other factors. For γ -rays and HE protons, $QF \simeq 1$; for thermal neutrons, $QF \simeq 3$; for fast neutrons, QF ranges up to 10; and for α particles and heavy ions, QF ranges up to 20.

Maximum *permissible occupational dose* for the whole body:

$$5 \text{ rem yr}^{-1} \text{ (or } \sim 100 \text{ mrem wk}^{-1}\text{).}$$

Fluxes (per cm^2) to liberate 1 rad in carbon:

$$3.5 \times 10^7 \text{ minimum ionizing singly charged particles,}$$

$$1.0 \times 10^9 \text{ photons of 1 MeV energy}$$

(These fluxes are correct to within a factor of 2 for all materials.)

Radiation exposure (mrem yr^{-1}) of a typical person in the US:

<i>Natural sources</i>		<i>Artificial sources</i>	
Cosmic radiation	28	Environmental	8
Terrestrial radiation	26	Medical	92
Internal isotopes	26	Occupational	1
		Nuclear power	0.3
		Miscellaneous	5

(This table has been adapted from *Particle Properties Data Booklet*, Lawrence Berkeley Laboratory, Berkeley, CA, 1980, and Upton, A. C., 'The biological effects of low-level ionizing radiation', *Sci. Amer.*, **246**, 41, Feb. 1982.)

Maximum permissible flux for occupational exposure to various types of ionizing radiations

Type of radiation	QF or RBE ^(a)	Average exposure rate ^(b) (mrad wk ⁻¹)	Approximate flux to give a maximum permissible exposure in an 8-hr day ^(c)
X- and gamma-rays	1	100	$[1400/E]$ photons cm ⁻² s ⁻¹ in free air at 0°C (error <13% for E = 0.07 to 2 MeV)
Beta-rays and electrons	1	100	$[(4.3 \times 10^{-7})/(RBE)P]$ electrons or beta-rays cm ⁻² s ⁻¹ ^(d)
Thermal neutrons	2.5	40	700 thermal neutrons cm ⁻² s ⁻¹ ^(d)
Fast neutrons	10	10	19 neutrons of 2-MeV energy cm ⁻² s ⁻¹ ^(d)
Alpha particles	10	10	$[(4.3 \times 10^{-7})/(RBE)P]$ alpha particles cm ⁻² s ⁻¹ ^(d)
Protons	10	10	$[(4.3 \times 10^{-7})/(RBE)P]$ protons cm ⁻² s ⁻¹ ^(d)
Heavy ions	20	5	$[(4.3 \times 10^{-7})/((RBE)P)]$ heavy ions cm ⁻² s ⁻¹ ^(d)

^(a) RBE, relative biological effectiveness.

^(b) Average occupational exposure rate permissible to blood-forming organs (essentially total body exposure), gonads, and eyes of persons of age 18 or over. These values may be averaged over a year, provided the dose in any thirteen weeks does not exceed 3 rem (rem = RBE × rad). All rates may be increased by a factor of 6 if the exposure is primarily to the skin, thyroid, or bone. They may be increased by a factor of 3 if the exposure is limited to organs other than blood-forming organs, gonads, or eyes.


^(c) Maximum permissible exposure rate based on a 20-mrem dose delivered to tissue in an 8-hr day. P is the stopping power in units of electron volts per g cm⁻² of soft tissue.

^(d) Incident on tissue.

(Adapted from *Nuclear Instruments and Their Uses*, A. H. Snell, ed., John Wiley and Sons, 1962.)

Maximum permissible flux for occupational exposure to various types of ionizing radiations (cont.)

Stopping power as a function of energy. (Adapted from *Nuclear Instruments and Their Uses*, A. H. Snell, ed., John Wiley and Sons, 1962.)



Clinical effects of acute exposure to ionizing radiation

Dose (rem)	Probable effect
0-100	No illness
100-200	No or slight illness
200-600	0-90% deaths, 2-12 weeks
600-1000	90-100% deaths, 1-6 weeks
1000-5000	100% deaths, 2-14 days
> 5000	100% deaths, 0-2 days

(Adapted from Glasstone, S. & Dolan, P. J., *The Effects of Nuclear Weapons*, US Government Printing Office, 1977.)

Bibliography

A Handbook of Radioactivity Measurement Procedures, NCRP Report No. 58, National Council on Radiation Protection and Measurements, Washington, DC 20014, 1978.

Chapter 19

Astronomical catalogs

	<i>page</i>
Astronomical catalogs	424
Selected astronomical catalog prefixes	428

Astronomical catalogs

X-ray sources

- Amuel, P. R., Cuseinov, O. H. & Rakhamimov, Sh. Yu., 1979, *Ap. J. Suppl.*, **41**, 327.
 Bradt, H. V., Doxsey, R. E. & Jernigan, J. G., 1978, *Adv. Space Exploration*, **3**.
 Forman, W. *et al.*, 1978, *Ap. J. Suppl.*, **38**, 357.
 Cooke, B. A. *et al.*, 1978, *M.N.R.A.S.*, **182**, 489.
 Markert, T. H. *et al.*, 1979, *Ap. J. Suppl.*, **39**, 573.
 Marshall, F. E. *et al.*, 1978, *NASA Tech Mem.* 79694.

Radio sources

- Dixon, R. S., 1970, *Ap. J. Suppl.*, **20**, 1.
 Finlay, E. A. & Jones, B. B., 1977, *Aust. J. Phys.*, **26**, 389.

NGC/IC objects

- Sulentic, J. W. & Tift, W. G., 1973, *The Revised New General Catalog of Non-Stellar Astronomical Objects*, University of Arizona Press.
 Schmidtke, P. C., Dixon, R. S. & Gearhart, M. R., 1979, *Palomar Sky Survey Overlays*, Ohio State University Radio Observatory.

Optical (non-stellar)

- Dixon, R. S. & Sonneborn, G., 1979, *Master Optical List*, Ohio State University Radio Observatory.

Infrared sources

- Schmitz, M. *et al.*, 1978, *Merged Infrared Catalog*, NASA Tech Mem. 79683, Goddard Space Flight Center, Greenbelt, Maryland.
 Price, S. & Walker, R., 1976, *The AFGL Four Color Infrared Sky Survey*, AFGL-TR-76-0208, Hanscom AFB, Massachusetts (Supplement: AFGL-TR-77-0160, 1977).
 Gezari, D. Y. *et al.*, 1984, *Catalog of Infrared Observations*, NASA ref. pub. 1118.

Gamma-ray sources

- Wills, R. D. *et al.*, 1980 *Adv. in Space Exploration*, Vol. 7, Pergamon Press.

Quasars

- de Veny, J. B., Osborn, W. H. & Hanes, K., 1972, *Pub. A.S.P.*, **83**, 611.
 Burbidge, G. R., Crowne, A. H. & Smith, H. E., 1977, *Ap. J. Suppl.*, **33**, 113.
 Hewitt, A. & Burbidge, A., 1987, *Ap. J. Suppl.*, **63**, 1.
 Hewitt, A. & Burbidge, G. R., 1980, *A Revised Optical Catalogue of Quasi-Stellar Objects*, *Ap. J. Suppl.*, **43**, 57.

Clusters of galaxies

- Abell, G., 1958, *Ap. J. Suppl.*, **3**, 211.
 Klemola, A. R., 1969, *A.J.*, **74**, 804.

Seyfert galaxies

- Weedman, D. W., 1977, *Ann. Rev. Astr. Ap.*, **15**, 69.
 Adams, T. F., 1977, *Ap. J. Suppl.*, **33**, 19.

Markarian galaxies

- Peterson, S. D., 1973, *A.J.*, **78**, 811.

Galaxies

- Sandage, A., 1961, *The Hubble Atlas of Galaxies*, Carnegie Institute, Washington, DC.

Zwicky, F. *et al.*, 1961–68, *Catalog of Galaxies and Clusters of Galaxies*, 6 vols., Calif. Inst. Tech. (Pasadena).
 Vorantsov, Velyaminov, 1964, *Morphological Catalog of Galaxies*.

Bright galaxies

de Vaucouleurs, G. & de Vaucouleurs, A., 1964, *Reference Catalog of Bright Galaxies*, University of Texas Press (Austin).
 Dressel, L. L. & Condon, J. J., 1976, *Ap. J. Suppl.*, **31**, 187.

Peculiar galaxies

Arp, H., 1966, *Ap. J. Suppl.*, **123**, 1.

H II regions in galaxies

Hodge, P. W., 1969, *Ap. J. Suppl.*, **157**, 73.
 Lynds, B. T., 1974, *Ap. J. Suppl.*, **267**, 391.

Infrared galaxies

Reike, G. H., 1978, *Ap. J.*, **226**, 550.
 Reike, G. H. & Lebofsky, M. J., 1978, *Ap. J.*, **220**, L37.
 Reike, G. H. & Low, F. J., 1972, *Ap. J.*, **176**, L95.
 Neugebauer, G., Becklin, E. E., Oke, J. B. & Searle, L., 1976, *Ap. J.*, **205**, 29.
 Kleinmann, D. E. & Low, F. J., 1970, *Ap. J.*, **159**, L165.

Stars

AGK3 Star Catalogue, 1975, Hamburger Sternwarte (Hamburg).
Smithsonian Astrophysical Observatory Star Catalog, 1966, Smithsonian Publ. 4652, US Government Printing Office, Washington, DC.
 Hoffleit, D., 1964, *Catalogue of Bright Stars*, Yale University Observatory (New Haven).
 Fricke, W. & Kopf, A., 1963, *Fourth Fundamental Catalog (FK4)*, Braun (Karlsruhe).
 Nagy, T. A. & Mead, J., 1978, *HD-SAO-DM Cross Index*, NASA Tech. Mem 79564, Goddard Space Flight Center, Greenbelt, Maryland.
 Schmidtke, P. C., Dixon, R. S. & Gearhart, M. R., 1979, *Palomar Sky Survey Overlays*, Ohio State Radio Observatory.
 Gottlieb, D. M., 1978, *Ap. J. Suppl.*, **38**, 287.
 Hirshfeld, A. & Sinnott, R. W., 1981, *Sky Catalogue 2000.0*, Sky Publishing Corp. (Cambridge, MA).
 Rufener, F., 1980, *Third Catalogue of Stars Measured in the Geneva Observatory Photometric System*, Observatoire de Genève, Switzerland.

Star clusters and associations

Alter, G., Balázs, B. & Ruprecht, J., 1970, *Catalogue of Star Clusters and Associations*, Akadémiai Kiado (Budapest).
 Fenkart, R. P. & Binggeli, B., 1979, *Astron. Ap. Suppl.*, **35**, 271.
 Becker, W. & Fenkart, R., 1971, *Astron. Ap. Suppl.*, **4**, 241.

Supergiants, O stars, and OB associations

Humphreys, R. M., 1978, *Ap. J. Suppl.*, **38**, 309.
 Cruz-Gonzales, C., Recillas-Cruz, E., Costero, R., Peimbert, M. & Torres-Peimbert, S., 1974, *Revista Mexicana de Astronomia y Astrofisica*, **1**, 211.
Luminous Stars in the Northern Milky Way, 6 vols., 1959–65, Hamburger Sternwarte and Warner and Swasey Observatories (Hamburg-Bergedorf).

Variable stars

Kukarkin, B. V. *et al.*, 1969, *Catalogue of Variable Stars*, 3rd edn. Astron. Council of the Academy of Sciences in the USSR (Moscow) (plus *Supplements*).

Ultraviolet

Catalogue of Stellar Ultraviolet Fluxes, 1978, The Science Research Council.

White dwarfs

McCook, G. P. & Scion, E. M., 1977, *Obs. Contrib.*, No. 2, Villanova Univ.

Nearby stars

Gliese, W., 1969, *Catalogue of Nearby Stars*, Veröffentlichungen des Astronom., Rechen-Inst., Heidelberg, No. 22, Verlag G. Braun (Karlsruhe).

Woolley, R., Epps, E. A., Penston, M. J. & Pocock, S. B., 1970, *Roy. Obs. Ann.*, No. 5.

Halliwell, M. J., 1979, *Ap. J. Suppl.*, **41**, 173.

Gliese, W. & Jahreiss, H., 1979, *Astron. Astrophys. Suppl.*, **38**, 423.

Proper motion and halo stars

Eggen, O. J., 1979, *Ap. J.*, **230**, 786.

Eggen, O. J., 1979, *Ap. J. Suppl.*, **39**, 89.

Eggen, O. J., 1980, *Ap. J.*, **43**, 457.

Luyten, W. J., 1979, *NLTT Catalogue*, 4 parts, Univ. of Minnesota (Minneapolis).

Luyten, W. J., 1963, *Bruce Proper Motion Survey*, 2 vols., Univ. of Minnesota (Minneapolis).

Luyten, W. J., 1961, *A Catalog of 7127 Stars in the Northern Hemisphere with Proper Motions Exceeding 0.2 arcsec Annually*, Lund Press (Minn.).

Luyten, W. J., 1976, *A Catalog of Stars with Proper Motions Exceeding 0.5 arcsec Annually*, University of Minnesota (Minneapolis).

Giclas, H., Burnham, R. Jr. & Thomas, N. G., 1971, *Lowell Proper Motion Survey (The G Numbered Stars)*, *Lowell Obs. Bull.*: Nos. 118, 125, 141, 153, 163, Lowell Obs., Flagstaff (Arizona).

Double stars

Aitken, R. G., 1932, *New General Catalog of Double Stars within 120' of the North Pole*, Publ. 417, Carnegie Inst., Washington.

Radial velocities

Abt, H. A. & Biggs, E. S., 1972, *A Bibliography of Stellar Radial Velocities*, Kitt Peak National Observatory.

Evans, D. S., 1978, *Catalog of Stellar Radial Velocities* (microfiche).

Moore, J. H., 1932, *A General Catalog of the Radial Velocities of Stars, Nebulae, and Clusters*, Publ. of Lick Obs., Vol. 18, Univ. of Calif. Press (Berkeley).

Wilson, R. E., 1953, *General Catalog of Stellar Radial Velocities Prepared at Mount Wilson Observatory*, Publ. 601, Carnegie Inst., Washington.

Stellar spectra

Houk, N. & Cowley, A. P., 1975, *Univ. of Michigan Catalog of Two-Dimensional Spectral Types for the HD Stars*, Vol. I (Ann Arbor).

Buscombe, W., 1977, *MK Spectral Classifications, Third General Catalog*, Northwestern Univ. (Evanston, Illinois).

Kennedy, P. M. & Buscombe, W., 1974, *MK Spectral Classifications Published Since Jaschek's La Plata Catalog*, Northwestern Univ. (Evanston, Illinois).

Jaschek, C., Conde, H. & de Sierra, A. C., 1964, *Catalog of Stellar Spectra Classified by the Morgan-Keenan System*, Obs. Astron. de la Univ. Nacional de la Plata (Argentina).

Seitter, W. C., 1970, *Atlas für Objektiv-Prismen-Spektren*, Dümmler (Bonn).

Breakiron, L. A. & Upgren, A. R., 1979, *Ap. J. Suppl.*, **41**, 709.

Globular clusters

Harris, W. E., 1976, *A.J.*, **81**, 1095.

Arp, H., 1966, in *Galactic Structure*, eds. A. Blaauw & M. Schmidt, Univ. of Chicago Press, p. 401.

Planetary nebula

Perek, I. & Kohoutek, L., 1967, *Catalogue of Galactic Planetary Nebula* (Czechoslovak Acad. of Sciences (Prague)).

Reflection nebulae

van den Bergh, S., 1966, *A.J.*, **71**, 990.

Supernova remnants

Downes, D., 1971, *A.J.*, **76**, 305.

Clark, D. and Caswell, J., 1976, *M.N.R.A.S.*, **174**, 267.

Pulsars

Manchester, R. N., Lyne, A. G., Taylor, J. H., Durdin, J. M., Large, M. I. & Little, A. G., 1978, *M.N.R.A.S.*, **185**, 409.

Damashek, M., Taylor, J. H. & Hulse, R. A., 1978, *Ap. J.*, **225**, L31.

OH masers

Turner, B. E., 1979, *Astron. Ap. Suppl.*, **37**, 1.

CO clouds

Kutner, M. L., Machnik, D. E., Tucker, K. D. & Dickman, R. L., 1980, *Ap. J.*, **237**, 734.

H II regions and spiral structure

Georgelin, Y. M. & Georgelin, Y. P., 1976, *Astron. Ap.*, **49**, 57.

Maršáková, P., 1974, *Ap. Space Sc.*, **27**, 3.

5 GHz continuum surveys (galactic plane)

Altenhoff, W. J., Downes, D., Pauls, T. & Schraml, J., 1978, *Astron. Ap. Suppl.*, **35**, 23.

Haynes, R. F., Caswell, J. L. & Simons, L. W. J., 1978, *Australian J. Phys.*, *Ap. Suppl.*, **45**, 1.

Haynes, R. F., Caswell, J. L. & Simons, L. W. J., 1979, *Australian J. Phys.*, *Ap. Suppl.*, **48**, 1.

Radio recombination line surveys

Downes, D. *et al.*, 1980, *Astron. Ap. Suppl.* **40**, 379.

Reifenstein, E. C. III, Wilson, T. L., Burke, B. F., Mezger, P. G. & Altenhoff, W. J., 1970, *Astron. Ap.*, **4**, 357.

H I (21 cm) surveys

Heiles, C. & Habing, H., 1974, *Astron. Ap. Suppl.*, **14**.

Heiles, C. & Jenkins, E. B., 1976, *Astr. Ap.*, **46**, 333.

Sun

Solar-Geophysical Data (monthly, 2 parts), NOAA, National Geophys. and Solar-Terrestrial Data Center (Boulder); *Solar-Geophysical Data, Descriptive Text*, 1974, No. 354 (Suppl.).

Planets

American Ephemeris and Nautical Almanac (yearly), US Government Printing Office, Washington, DC; *Explanatory Suppl.*, 1975, HM Stationery Office, London.

Stellar rotational velocities

- Bernacca, P. L. & Perinotto, M., *A Catalog of Stellar Rotational Velocities*, Contrib. Oss. Astrofis. Asiago, Univ. Padova, No. 239, 1970; No. 250, 1971; No. 294, 1973.
- Boyarchuk, A. A. & Kopylov, I. M., 1964, *A General Catalog of Rotational Velocities of 2558 Stars*, Publ. Crimean Astrophys. Observ., 31, 44.
- Uesugi, A. & Fukuda, I., 1970, *A Catalog of Rotational Velocities of the Stars*, Contrib. Instit. Astrophys. and Kwasan Obser. Univ. Tokyo, No. 189.

Radio galaxies

- Burbidge, G. & Crowne, A. H., 1979, *Ap. J. Suppl.*, 40, 583.

Dark clouds

- Lynds, B. T., 1962, *Ap. J. Suppl.*, 7, 1.

Binaries

- Batten, A. H., Fletcher, J. M. & Mann, P. J., 1978, *Seventh catalogue of the orbital elements of spectroscopic binary systems*, Pub. Dominion Astrophys. Obs., vol. XV, No. 5.
- Wood, F. B., Oliver, J. P., Florkowski, D. R. & Koch, R. H., 1980, *Finding list for observers of interacting binary stars*, University of Pennsylvania Press.

Machine-readable astronomical catalogues

- The Astronomical Data Center of the NASA-Goddard Space Flight Center, Greenbelt, MD 20771, maintains a large number of machine-readable astronomical catalogues. See *Astronomical Data Center Bulletin*, NSSDC/WDC NASA-Goddard Space Flight Center.

Selected astronomical catalog prefixes (see references in previous section)

X-ray and gamma-ray

IE	HEAO-2 (Einstein).
H	HEAO-1, A-2 experiment (GSFC).
XRS	Amnuel <i>et al.</i> compilation.
4U, 3U, etc.	Uhuru catalogs.
IM	OSO-7 catalog (MIT).
2A, A	Ariel catalogs.
2S	SAS-3 source (MIT).
CGS	Bradt <i>et al.</i> galactic sources.
MXB	MIT burst source (Bradt <i>et al.</i>).
CG	(cosmic gamma-ray), usually COS-B source.

Radio

G	(galactic coordinates), various sources – usually continuum surveys.
3C	(3rd Cambridge) 1959, <i>Mem. R.A.S.</i> , 68, 37; 1962, 68, 163.
4C	(4th Cambridge) 1965, <i>Mem. R.A.S.</i> , 69, 183; 1967, 47, 49.
W	(Dwingeloo) Westerhout 1958, <i>B.A.N.</i> , 14, 215.

CTA, CTB, CTD	(Cal Tech) 1960, <i>Publ. A.S.P.</i> , 72 , 237; <i>Cal. Tech. Radio. Obs. Reports</i> (# 2) 1960-65, 1963, <i>A.J.</i> , 68 , 181
NRAO	(Green Bank) 1966, <i>Ap. J. Suppl.</i> , 116 .
PKS	(Parkes) 1964, <i>Australian J. Phys.</i> , 17 , 340; 1965, 18 , 329; 1966, 19 , 35; 1966, 19 , 837; 1968, 21 , 377.
MSH	(Sydney) Mills, Slee & Hill, 1958, <i>Australian J. Phys.</i> , 11 , 360; 1960, 13 , 676; 1961, 14 , 497.
OA OZ	(Ohio State) 1966, <i>Ap. J.</i> , 144 ; 1967, <i>A.J.</i> , 72 , 536; 1968, 73 , 381; 1969, 74 , 612; 1970, 75 , 351; 1971, 76 , 777.
AMWW	(Bonn) Altenhoff, Mezger, Wendkar & Westerhout, 1960, <i>Publ. Univ. Obs. Bonn.</i> , No. 59.

Optical-stars general

HD	<i>Henry Draper Catalog</i> 1918–25, <i>Harvard Ann.</i> , 91–100.
AGK #	<i>Astronomische Gesellschaft Katalog</i> .
FK #	<i>Fundamental Katalog</i> .
SAO or #####	<i>Smithsonian Astrophysical Observatory Catalog</i> .
GC	<i>General Catalog</i> , Boss, 1936, Carnegie Inst. Wash. Publ. 468.
BD	<i>Bonner Durchmusterung</i> , 1860, <i>Beob. Bonn. Obs.</i> , 3 ; 4 ; 5 .
SD	<i>Southern Durchmusterung</i> , 1886, <i>Beob. Bonn. Obs.</i> , 8 .
CD (or CoD)	<i>Cordoba Durchmusterung</i> , 1892, <i>Result. Natl. Obs. Argentina</i> , 16 ; 17 ; 18 ; 21a ; 21b .
CPD	<i>Cape Photographic Durchmusterung</i> , 1896, <i>Ann. Cape Obs.</i> , 3 ; 4 ; 5 .
DM	BD, CP, CPD combined.
± # #' ...	Usually DM catalogs.
HR	(Harvard revised) <i>Harvard Ann.</i> , 1908, 50 .
BS	(Bright star) <i>Yale Bright Star Catalog</i> . Follows HR numbering system (BS = HR #).

Optical-stars proper motion

G ### – ###	Lowell P.M. Surveys.
(or GD, HG)	
BPM (or L)	Bruce P.M. Survey.
LP	(Luyten-Palomar) 1969a, 1969b, Luyten 1963, <i>P.M. Survey with the 48-Inch Schmidt</i> , Univ. Minn., Minneapolis.
LHS	(Luyten Half-Second) Luyten, 0.5" yr ⁻¹ , P.M. Survey.
LTT	Luyten, 0.2" yr ⁻¹ , P.M. Survey.
NLTT	Luyten, new P.M. Catalog.
LB, etc.	other Luyten P.M. Catalogs.

Optical stars-miscellaneous

EG or GR	(White Dwarfs) Eggen and/or Greenstein; EG: <i>Ap. J.</i> , 1965, 141 , 83; 1965, 142 , 925; 1967, 150 , 927; 1969, 158 , 281. GR: <i>Ap. J.</i> , 1970, 162 , L55; 1974, 189 , L131; 1975, 196 , L117; 1976, 207 , L119; 1977, 218 , L21; 1979, 227 , 244. Also Greenstein, 1976, <i>A.J.</i> , 81 , 323; 1976, <i>Ap. J.</i> , 210 , 524.
GL	Gliese, W., 1969, <i>Catalog of Nearby Stars</i> , G. Braun, Karlsruhe.

- Y (Yale) Jenkins, L. F., 1952, *General Catalog of Trigonometric Stellar Parallaxes*, Yale Univ. Obs., New Haven. (Also 1963, *Suppl.*)
 (Yerkes) van Altena *et al.*, *A.J.*, 1969, **74**, 2; 1971, **76**, 932; 1973, **78**, 781; 1973, **78**, 201; 1975, **80**, 647.
- HZ Humason & Zwicky, 1946, *Ap. J.*, **105**, 85-91.
- Wolf (or W) Nearby star discovered by Max Wolf (see Gliese catalog for data).
- Ross (or R) Nearby star discovered by Frank Ross (see Gliese catalog for data).
- PHL (Ton, Tn, TS) (Palomar-Haro-Luyten) Haro and Luyten, 1962, *Bol. Obs. Tonantzintla y Tacubaya*, **3**, 37. (Faint blue stars.)
- VB Van Biesbroeck, G., 1961, *A.J.*, **66**, 528.
- Feige (or F) Feige, 1958, *Ap. J.*, **128**, 267.

Optical-stars-variable

Naming convention (if no standard name):

Constellation preceded by the following combinations in order of variability discovery:

R, S, T, . . . , Z, RR, RS, . . . , RZ, SS, . . . , SZ, . . . , ZZ, AA, AB, . . . , AZ, BB, . . . , BZ, . . . , QQ, . . . , QZ, V335, V336, . . . (Note: the letter J is not used.)

Optical-miscellaneous galactic

- TR Trumpler, R., 1930, *Lick Obs. Bull.*, No. 420 (associations).
- Coll Collinder, P., 1931, *Ann. Obs. Lund.*, No. 2 (associations).
- RCW Rodgers, Campbell & Whiteoak, 1960, *M.N.R.A.S.*, **121**, 103 (H II regions).
- R Reflection nebula, preceded by constellation, as in Mon R2.
- S Sharpless, 1959, *Ap. J. Suppl.*, **4**, 257 (H II regions).
- SS Stevenson and Sanduleak object.
- HH Herbig-Haro object. Herbig, 1951, *Ap. J.*, **113**, 697; Haro, 1952, *Ap. J.*, **115**, 572; Herbig, 1974, *Lick Obs. Bull.*, **658**.

Optical-general-non-stellar

- NGC Dreyer's *New General Catalog*.
- IC Dreyer's *Index Catalog*.

Optical-extragalactic

- Mrk (or Mkn) Markarian, *Astrofizika* (in Russian), 1967, **3**, 55; 1969, **5**, 443; 1969, **5**, 581; 1971, **7**, 511; 1971, **8**, 155; 1973, **9**, 488; 1974, **10**, 307; 1976, **12**, 390; 1976, **12**, 657; 1977, **13**, 225.
- Zw Zwicky.
- MCG *Morphological Catalog of Galaxies*.

Infrared

- IRC (or TMSS) (Infrared Catalog) Neugebauer, G. & Leighton, R. B., 1969, *Two-Micron Sky Survey*, Cal. Tech., NASA SP-3047.
- AFGL Air Force Geophys. Lab.
- GMS Gillett, Merrill & Stein, 1971, *Ap. J.*, **164**, 83.
- Hall Hall, R. J., 1974, *A Catalog of 10- μ m Celestial Objects*, Space and Missile Systems Org., SAMSO-TR-74-212.
- MIRC *Merged Infrared Catalog*.
- BN Becklin-Neugebauer object (in Orion Nebula), 1967, *Ap. J.*, **147**, 799.
- KL Kleinmann-Low object (in Orion Nebula), 1967, *Ap. J.*, **149**, L1.

(The material in this chapter was prepared by C. W. Maxson of the Harvard-Smithsonian Center for Astrophysics.)

Diskettes for PCs

A template library is available for IBM PCs and compatibles running DOS 2.0 or higher versions. This library is for version 1A or higher of LOTUS® 1-2-3 and all compatible spreadsheet programs. The library is available on diskettes and contains most of the complex formulae and much of the tabular data of this handbook and additional material. It is menu-driven and will yield graphical and tabular output for 'what-if' problems in spherical astronomy, including coordinate transformations and rigorous precession and nutation corrections, planetary positions, binary star orbits, UV, X-ray, and gamma-ray absorption coefficients, relativity, unit conversions, plasma physics parameters, synchrotron radiation, least-squares fitting, statistics, electron ranges, X-ray reflectivity, blackbody radiation, vacuum physics, interstellar photoelectric absorption, and atmospheric absorption.

Requests for information should be directed to SCIENCE.wks, P.O. Box 426, Winchester, MA 01890, USA.

Index

- Aberration, 124-5
 - constant, 10
- Absolute magnitude, 105
 - of galaxies, 85
 - of stars, 45-9, 105
- Absorption cross-sections, 177, 198-202, 298
 - oscillator strength, 257
- Abundances, 30, 221-3
- AGNs, 87
- Alfvén speed, 284
- Alpha particle sources, 312
 - range-energy, 308-9
 - specific energy loss, 309
- Angstrom unit, 11
- Angular density vs. redshift, 253
 - size vs. redshift, 250-51
- Apparent magnitude, 98
 - place, 125
 - solar time, 107, 125
- Approximations, 405
- ASCII character set, 370-3
- Astronomical catalogs, 424-8
 - catalog prefixes, 428-30
 - coordinate transformations, 111
 - facilities, 114-19
 - photometry, 98-102
 - unit, 10, 126
- Astrophysical plasma parameters, 286
- Atmosphere, attenuation of photons in, 203, 240
 - composition, 239
 - constituents, 239
 - density vs. altitude, 238
 - electron density vs. altitude, 239
 - International Reference, 238
 - pressure vs. altitude, 241
 - standard, 8
 - structure, 243
 - temperature vs. altitude, 238, 242
 - transmission, 164
 - US standard, 241-2
- Atmospheric composition, 239
 - constituents, 239
 - depth, 229
 - opacity, 155, 240
 - transmission, 164
- Atomic energy levels, 258-60
 - mass unit, 8, 10, 11
 - number, 20-1
 - physics, 256-7
 - scattering coefficient, 257
 - scattering factor, 322
 - time, 107-8
 - weight, 20-1
- Attenuation coefficients, 291-303
 - of electromagnetic radiation, 291-303
 - lengths, 295
 - of photons in the atmosphere, 79, 203, 240
 - of photons in the interstellar medium, 198-202, 204
- Autocorrelation function, 400
- Autocovariance function, 400
- Avogadro's constant, 6
- Background, EUV, 172
 - gamma-ray, 216
 - infrared, 162-3, 165
 - microwave, 143-4
 - X-ray, 197
- Ballistic coefficient, 380

- Bar, 13
- Barn, 11
- Bessel function, 314, 404
- Besselian year, 107, 139
- Beta-ray attenuation, 308
 - range-energy, 311
- Binary number system, 16
- Binary star elements, 61
- Binomial distribution, 410
- Blackbody, flux densities, 165
 - radiation, 268-9
- BL Lac objects, 88
- Bohr magneton, 3, 10
 - radius, 3, 10
- Bolometric correction, 68 70, 102
 - magnitude, 102
- Boltzmann constant, 7, 10
 - formula, 257
- Boolean algebra, 373
- Bragg-Kleeman rule, 308
- Bremsstrahlung radiation, 206 8, 276
- Brighter galaxies, 85
- Brightest stars, 44-9
- Brightness, Moon, 105
 - night sky, 106
 - planetary, 106
 - Sun, 106
 - temperature, 165
- BTU, 13, 19

- Calorie, 11, 19
- Celestial coordinates, 110
 - sphere, 109
- Cepheid variable, 62
- Characteristic lines, 320
- Chart, equatorial to galactic coordinates, 112
 - precession, 113
 - star, 38-9
- Chi-square distribution, 413
 - test, 414
- Chromosphere, 27
- Civil day, 107
- Classification, spectral, 64-5
 - of variable stars, 62-3
 - star, 71
- Clinical effects of radiation, 420
- Clusters of galaxies, 91-2, 193
- Color excess, 103
 - index, 99, 103
 - temperature, 165
- Comets, 36

- Compton scattering, 272-5
 - inverse, 274
 - shift, 272
 - wavelength, 4, 10
- Constellations, 40-3
- Constants, astronomical, 10-11
 - numerical, 16
 - physical, 2 10
- Conversions, 11, 12-15, 19, 278, 343
- Convolution, 400
- Coordinate transformations, 396
- Corona, 27
- COS-B catalog, 212
- Cosmic background radiation, 143-4
 - temperature, 11
- Cosmic ray abundances, 221 3
 - altitude variation, 228
 - composition, 220
 - energy density, 93
 - magnetic rigidity, 225
 - spectra, 224
- Cosmological constant, 248
 - data, 11
- Cosmology, 248-54
- Coloumb gauge, 278
 - logarithm, 285
- Crab nebula, 64, 192, 212, 215
- Critical density, 11, 249
- Cross-correlation, 400
- Cryogenic fluids, 334
- Crystal properties, 319
 - rocking curve, 318
 - scintillators, 366-7
 - solid-state, 366-7
 - spectroscopy, 318
- Curie, 11, 418

- Data transmission nomogram, 378
- Day, civil, 107
 - sidereal, 107
 - solar, 107
- Debye length, 284, 286
- Definite integrals, 399
- Density, critical, 11, 249
 - of crystals, 319
 - data for astronomical objects, 94
 - of elements, 20-1
 - of hydrogen, 180 2
 - of detector materials, 366-8
- Detectors, visible and ultraviolet, 348-58
 - X-ray and gamma-ray, 358-69
- Deuteron magnetic moment, 6
 - mass, 6

- Diffraction, 337–8
- Dispersion measure, 55
- Distance, luminosity, 250
 - modulus, 105
 - vs. redshift, 252
- Doppler effect, 246
 - shift, 257
 - width, 257
- Dynamical form factor, Earth, 10

- Earth's atmosphere, 232–43
 - dynamical form factor, 10
 - environment, 232–44
 - equatorial radius, 10
 - equatorial surface gravity, 31
 - magnetic dipole moment, 10
 - magnetosphere, 244
 - mass, 10, 31
 - mean surface gravity, 10
 - radiation environment, 232–35
 - radius, 10, 31
 - rotational energy, 19
 - rotation rate, 10
 - satellite lifetime, 380
 - total heat content, 19
- Ecliptic, 109
 - obliquity of, 111
 - system, 111
- Effective temperature, 68–70, 103, 165
- Einstein coefficient, 257
 - field equations, 248
- Electromagnetic conversion table, 278
 - definitions, 279
 - field strength tensor, 247
 - radiation attenuation, 291–303
 - spectrum nomogram, 281
- Electron charge, 2, 4, 10, 17
 - classical radius, 4, 10
 - Compton wavelength, 4, 10
 - cosmic rays, 224
 - density, 27, 156, 286
 - drift velocity, 284
 - energy loss, 304–7
 - energy standards, 313
 - flux contours, 235
 - gyrofrequency, 284
 - gyroradius, 285
 - ion collision frequency, 285
 - magnetic moment, 4, 17
 - mass, 3, 10, 17
 - radius, 4, 10
 - range-energy, 307, 311
 - resistivity, 285
 - rest energy, 3
 - spin, 17
 - storage ring, 316–17
 - temperature, 286
 - thermal velocity, 284
 - volt, 8
- Elementary particles, 17–18
- Elements, abundance, 30
 - atomic numbers, 20–1
 - atomic weights, 20–1
 - boiling points, 20–1
 - densities, 20–1
 - melting points, 20–1
 - periodic table, 20–1
- Emission coefficient, 279
 - of hot plasma, 276–7
 - line widths, 71
 - lines, 173, 206, 276
- Emissivity, 279
- Energy content of space, 93
 - conversions, 19
 - density of the Galaxy, 82
 - levels, 258–61
 - loss, 304–10
- Ephemeris day, 10
 - second, 107
 - time, 107–8, 129
- Equatorial coordinates, 111
 - system, 111
- Erg, 19
- EUV absorption cross-sections, 177
 - attenuation, 179
 - background, 172
 - sources, 171
 - spectra, 173–4
 - spectral features, 175
 - strong lines, 176
- Excitation data for gases, 365

- Faraday constant, 6
 - rotation, 154
- Feigenbaum's number, 16
- Filter materials for X-ray detectors, 368
- Fine structure constant, 3, 10
- Fluorescent converters, 358
 - yields, 304
- Flux unit, 11
- Fourier theorems, 401
 - transforms, 388–402
- Fraunhofer diffraction, 338
- Fresnel's equations, 323
- Friedmann universes, 248

- Fundamental physical constants, 2-10
- Galactic coordinates, 111
- Galaxies, classification, 84
 - clusters, 91-2
 - liners, 89
 - local group, 83
 - luminous emission, 11
 - mean sky brightness, 11
 - named, 86-7
 - nearby, 84
 - radio, 89
 - selected brighter, 85
 - Seyfert, 88
 - space density, 11
 - X-ray emission, 193
- Galaxy, age, 82
 - diameter, 82
 - energy density, 82
 - luminosity, 82
 - mass, 82
 - nucleus, 82
 - period of rotation, 82
 - properties, 82
 - type, 82
- Gamma-ray absorption, 291-303
 - attenuation, 291-303
 - background, 216
 - Crab Nebula spectrum, 215
 - detectors, 369
 - downward flux, 218
 - energy standards, 312
 - production in the atmosphere, 227
 - source catalogs, 212-13
 - source map, 212
 - sources, 212, 214
 - spectral features, 217
- Gaunt factor, 276
- Gaussian distribution, 408
- Gaussian gravitational constant, 130
- Gauss-Legendre quadrature, 405
- Geocoronal emission, 172
- GMT, 109
- Grating spectroscopy, 321
- Gravitational constant, 2, 10
- Gravity, Earth surface, 10
 - planetary surface, 31
 - solar surface, 25
- Gregorian calendar, 131
- Greek alphabet, 22
- Gyrofrequency, 284
- Gyroradius, 285
- HEAO A1 All Sky Map, 185
- Henry Draper spectral classification, 64
- Hexadecimal number system, 16
- Hertzsprung-Russell (HR) diagram, 66
- Hour angle, 109, 111, 131
- H-R diagram, 66
- HST, 120-4
- Hubble, classification of galaxies, 84
 - constant, 11
 - distance, 11
 - radius, 249
 - Space Telescope, 120-4
 - time, 11
- Hydrogen densities, 81, 180-2
- IEEE interface, 377
- Image intensifier, 350
- Index of refraction, 322, 325
- Infrared background, 162-3
 - diffuse fluxes, 163
 - source temperatures, 165
 - sources, 160-1
 - transmission of the atmosphere, 164
- Ionospheric electron density, 156
- Intensity, 11, 279
- International atomic time, 108, 132
- International Reference Atmosphere, 238
- Interplanetary medium, 153
- Interstellar abundances, 199-202
 - EUV attenuation, 179
 - extinction, 178
 - gas parameters, 81
 - hydrogen densities, 81, 180-2
 - magnetic field, 81
 - medium, 154-5
 - medium cross-section, 198-202
 - photoabsorption cross-sections, 198-202
 - reddening, 103
 - reddening law, 104
 - scintillation, 155
- Ion gyro-frequency, 284
 - gyroradius, 285
 - sound velocity, 284
- Ionization data for gases, 365
 - loss, 304-7
- Ionosphere, 156, 243, 286
- Jansky, 11
- JHKLM system, 101
- Joule, 11, 19
- Julian calendar, 132
 - data, 107, 132

- K alpha, 262-3
 - beta, 262-3
 - edge, 292
 - lines, 262-3
 - series, 262-3
- Kepler's third law, 61
- keV, 11
- Klein-Nishina cross-section, 272-4

- L alpha, 264-5
 - beta, 264-5
 - lines, 264-5
 - series, 264-5
- Launch vehicles, 384-94
- Least-squares fit, 406, 415
- Light, speed, 2, 10
 - year, 11, 133
- Linear interpolation, 405
- Linear least-squares, 406, 415
- Liners, 89
- Local group of galaxies, 83
- Logic gates, 374
- Look-back time, 254, 249
- Lorentz factor, 271
 - force, 247
 - gauge, 278
 - transformation, 246
- Loschmidt constant, 7
- LS coupling, 256
- Luminosity, absolute, 102
 - class, 65
 - distance, 250, 252
 - functions, 74-5, 80-1
 - stellar, 72-3
 - vs. surface temperature, 66
- Magnetic dipole moment of Earth, 10
 - field, 81-2, 93, 286
 - rigidity, 225
- Magnetosphere, 244
- Magnitude, stellar, 98
- Main sequence, 66, 68-9
- Mass, atomic unit, 8, 10, 11
 - attenuation coefficients, 291-303
 - data for astronomical objects, 94
 - function, 74-5
 - loss rate, 170
 - relativistic, 246
 - rest, 246
- Mathematical formulae, 17
- Maxwell's equations, 247, 277, 280
- Maxwell velocity distribution, 285
- Mean free path, 333
 - life of elementary particles, 17-18
 - magnetic field in a plasma, 285
 - plane, 134
- Mesons, 17-18
- Messier catalog, 95-7
- Meteor streams, 37
- Microwave background, 143-4
- MK luminosity class, 65
- Mile, nautical, 11
 - statute, 11
- Moon, apparent magnitude, 105
 - phase law, 105
 - physical elements, 31
- MTF, 335-7
- Muon charge, 17
 - magnetic moment, 4, 17
 - mass, 4, 17-18
 - mean life, 17-18
 - spin, 17-18

- Natural satellites, 33-5
- Neutrino charge, 17
 - energy density, 93
 - magnetic moment, 17
 - mass, 17
 - spin, 17
- Neutron charge, 17
 - Compton wavelength, 5
 - magnetic moment, 6, 17
 - mass, 5, 6, 10, 17-18
 - mean life, 17-18
 - spin, 17-18
- Night sky brightness, 106
- Novae, 63
- Number system conversions, 16
- Numerical analysis, 404-6
 - constants, 16
- Nutation, 134
 - constant, 10

- OB associations, 59-60
- Obliquity of the ecliptic, 10, 135
- Octal number system, 16
- Optical constants, 325
 - depth, 179
 - telescopes, 114-18, 120-4, 339-40
- Oscillator strength, 257

- Parallax, 135
 - solar, 10
- Parallel interface, 376
- Parsec, 11, 135

- Particle production, 226
- Pascal, 13
- PDMF, 74-5
- Periodic table of the elements, 20
- Permeability of vacuum, 2
- Permittivity of vacuum, 2
- Photocathode, 349, 353-7
- Photodiode, 348-9
- Photometry, 341 7
 - astronomical, 98 9
 - filter bands, 100
 - lunar, 105
 - response curves, 100-1
 - standard system, 100
- Photomultiplier, 351-2
- Photon, spin, 17
- Photosphere, 27
- Physical constants, 2-10
- Pion, charge, 17
 - magnetic moment, 17
 - mass, 17-18
 - mean life, 17-18
 - spin, 17-18
- Planck constant, 2, 10
 - functions, 268
 - length, 2
 - mass, 2
 - radiation curves, 269
 - time, 2
- Planets, brightness, 106
 - orbital elements, 32
 - physical elements, 31
- Plasma emission, 173-4, 206-8, 276-7
 - oscillation frequency, 284
 - parameters, 284-6
- Poisson distribution, 410-12
- Point spread function, 334-8
- Positron, 17
- Power spectrum, 400
- Precession, general, 10, 136
- Prefixes, 19
- Probability distributions, 408 13
- Propagation of errors, 415
- Proper motion, 53, 136
- Proportional counter, 360 1
 - counter gases, 367
- Proton charge, 17
 - Compton wavelength, 5
 - electron mass ratio, 5
 - energy loss, 309
 - magnetic moment, 5, 17
 - mass, 5, 10, 17 18
 - range-energy, 308-9
 - spin, 17 18
- Pulsars, 54 8, 189-90
- Quadrature, 136, 404
- Quasars, catalogs, 424
 - infrared fluxes, 161, 163
 - radio fluxes, 145-50
 - selected, 87
 - X-ray luminosities, 195
 - 3C272, 87, 145, 148, 150, 193-4, 21
- QF, 418
- Rad, 418
- Radiation environment, 232
 - exposure, 418
 - length, 306
 - physics, 418 20
 - trapped, 233 5
- Radiative loss, 305-7
- Radio, brightest sources, 148-9
 - flux calibrators, 151-2
 - galaxies, 89
 - propagation effects, 153-5
 - selected sources, 145 6
 - source catalogs, 424
 - spectra, 150
 - spectral features, 157
 - surveys, 142
 - telescopes, 118-19
- Radioactive sources, 311-13
- Range-energy for particles, 307-11
- Rayleigh, 11
- Rayleigh-Jeans law, 268
- RBE, 419
- Recombination radiation, 277
- Redshift, angular density vs., 253
 - angular size vs., 251
 - correction, 250
 - functions, 251-4
 - look-back time vs., 254
 - luminosity distance vs., 252
 - magnitude relationship, 250
 - objects with large, 90
- Relativity, special, 246
- Rem, 418
- Robertson Walker line element, 248
- Roentgen, 418
- RS232 interface, 375
- Rydberg constant, 2, 10
- Sakur-Tetrode constant, 7
- Satellite lifetimes, 380
- Scintillation materials, 366-7

- Sidereal day, 107
 - local time, 109
 - period of planets, 31
 - rotation of Sun, 25
 - time, 107
- Seyfert galaxies, 88
- Simpson's rule, 405
- Solar (*see* Sun) chromosphere, 27
 - constant, 25
 - corona, 27
 - day, 107
 - electron density, 27
 - effective temperature, 25
 - EUV flux, 29
 - flare, 236-7
 - high energy particle radiation, 232
 - gravity, 25
 - irradiance, 236
 - luminosity, 25
 - magnetic field strength, 25
 - magnitude, 25, 28, 106
 - mass, 25
 - mean density, 25
 - neighborhood density, 82
 - parallax, 10
 - photosphere, 27
 - radius, 25
 - spectral irradiance, 28, 237
 - spectrum, 237
 - system abundances, 30
 - temperature, 25, 27
 - time, 107
 - ultraviolet radiation, 236-7
 - X-ray radiation, 236-7
- Solid-state materials, 366-7
- Space Shuttle, 381-3
- Space Telescope, 120-4
- Special relativity, 246
- Spectra, blackbody, 268-9
 - EUV, 173
 - radio source, 150
 - UV 168-9
 - X-ray, 205
- Spectral features, gamma-ray, 217
 - infrared and visible, 67
 - EUV, 175
 - irradiance of the Sun, 28
 - radio, 157
 - type, 65
 - ultraviolet, 175
 - X-ray, 204-5
- Spectrograph, 321
 - grazing incidence, 321
- Spectroscopic terminology, 256
- Spectroscopy, crystal, 318
 - grating, 320
- Spherical astronomy, 107-13
- Stars (*see* stellar), absolute magnitude, 71-2
 - binary, 61
 - brightest, 44-9
 - catalogs, 425-6
 - charts, 38-9
 - classification, 71
 - counts, 80
 - giant, 70
 - infrared emission, 160
 - integrated light, 78
 - large proper motion, 53
 - nearest, 50-2
 - number densities, 76-9
 - relative numbers, 78
 - ultraviolet spectra, 168
 - variable, 62-3
 - X-ray emission, 188
- Stefan-Boltzmann constant, 7, 10
 - law, 268
- Stellar (*see* stars) density, 72-3
 - EUV sources, 171
 - luminosity, 72-3
 - luminosity function, 80-1
 - magnitude, 71, 98
 - mass, 72-3
 - mass function, 74-5
 - mass loss rates, 170
 - radius, 72-3
 - spectral irradiance, 98, 103
 - surface flux, 170
 - temperatures, 66
 - UV spectra, 168-9
 - winds, 170
 - X-ray flux, 188
- Steradian, 15-16
- Stopping power, 420
- Sun (*see* solar) - Earth system constants, 10
 - distance from Galactic center, 82
 - height above Galactic disk, 82
 - magnetic field, 25
- Sunspot, magnetic field, 25
 - numbers, 26
- Supernovae remnants, 64
 - X-ray emitting, 191
- Synchrotron radiation, 270-1, 314-17
- Synodic period of planets, 32
 - rotation of Sun, 25
- Taylor series, 404

- Telescopes, configurations, 339–40
 ground-based, 114–19
 radio, 118
 reflecting, 114–16
 refracting, 116–18
 Schmidt, 115–16
 Space, 120–4
 X-ray, 332
- Temperature, brightness, 165
 colour, 165
 effective, 68–70, 165
 electron, 286
 of the interstellar gas, 81
 -luminosity diagram, 66
 solar distribution, 27
 surface, 66
- Tesla, 11
- Thomson cross-section, 4, 10
- Thomson limit, 274
- Time, 107
 barycentric dynamical, 108, 126
 sidereal, 107
 solar, 107
 standard, 108
 terrestrial dynamical, 108, 138
 universal, 108, 139
- Torr, 11, 13
- Trapezoidal rule, 404
- Tropical year, 10, 107, 140
- UBV system, 100
- Uhuru map, 184
 unit, 11
- Ultraviolet absorption, 177
 background fluxes, 172
 fluorescent converters, 358
 source catalog, 426
 spectra, 168–9
 transmission, 357
- Unit conversions, 11–15
- Universal time, 139
- UV extinction, 178
 stellar spectra, 168–9
 window transmission, 357
- Vacuum technology, 333
- Variable stars, 62–3
- Vector analysis, 396–8
- Water vapor in the atmosphere, 239
- Weak coupling constant, 11
- Wien's displacement law, 268
 law, 268
- Wilson-Bappu, 71
- Window materials, UV detectors, 357
 X-ray detectors, 368
- Wolter mirror, 332
- X-ray absorption, 198–204
 atomic energy levels, 258–61
 attenuation, 203
 background, 197
 characteristic lines, 262–5, 320
 detectors, 358–64
 emission from galaxies, 193
 energy level diagram, 261
 line emission, 206–8
 material properties, 367–8
 mirror, 332
 optical parameters, 325
 pulsars, 189–90
 reflection of, 322–4
 reflectivity, 326–31
 source catalogs, 424
 source nomogram, 196
 sources, 184–95, 214, 311
 spectra, 205–8
 spectroscopy, 318
 supernova remnants, 191
 telescope, 332
 wavelengths, 262–5
- XU unit, 8, 11
- XUV absorption cross-sections, 198–9
 attenuation, 179
 spectrum, 173–4
- Year, tropical, 10, 107, 140
- Zenith, 140, 153
- Zodiac, 110

50.000

The Second Edition of this widely acclaimed handbook has been completely revised, re-designed and re-set. Many additions have been made to the data and its method of presentation. In response to numerous suggestions from users, many entirely new features have been added. The First Edition quickly established itself as an essential reference for space astronomy and astrophysics. Now this revision brings even more information to the desktop of every space astronomer. It saves valuable time by presenting the latest information in tables, graphs and formulas. Much of this information is presently scattered throughout the primary literature, and is here assembled in one place for the first time. The material is indispensable to graduate students, technicians and researchers working in space astronomy.

Cambridge University Press

ISBN 0-521-34787-4



9 780521 347877